



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

TX 512.Z .M659ac
Milne, William J.
Academic algebra /

Stanford University Libraries



3 6105 04932 7070



SCHOOL OF EDUCATION
LIBRARY

GIFT OF
THE PUBLISHERS



STANFORD UNIVERSITY
LIBRARIES

Gift of
American Book Company



ACADEMIC ALGEBRA

BY

WILLIAM J. MILNE, PH.D., LL.D.

PRESIDENT OF NEW YORK STATE NORMAL COLLEGE, ALBANY, N.Y.



NEW YORK: AMERICAN BOOK COMPANY, 1904.

CHICAGO: AMERICAN BOOK COMPANY, 1904.

CINCINNATI: AMERICAN BOOK COMPANY, 1904.

NEW YORK · CINCINNATI · CHICAGO
AMERICAN BOOK COMPANY

34

COPYRIGHT, 1901, BY
WILLIAM J. MILNE.

ENTERED AT STATIONERS' HALL, LONDON.

ACADEMIC ALGEBRA.

E-P 3

C

THE
ROYAL CANADIAN MOUNTED POLICE
ALGEBRA

PURPOSE AND PLAN OF THIS BOOK

THE Academic Algebra has been prepared to meet the requirements of the most exacting entrance examination of any College or University in the United States.

The book contains a thorough treatment of the science, so far as it is taught in the secondary schools. A full development of each subject, and a clear statement of its principles and laws, precedes the proofs of the principles—an arrangement that makes it possible for a teacher, without hindrance to the progress of the student, to postpone, if he sees fit, the rigorous proofs. The examples and problems are sufficiently numerous and complex to test the student's skill in applying all the principles that are developed. They are carefully graded, increasing in difficulty in each subject, so that, if desired, a brief and easier course may be conveniently provided by omitting the more difficult problems at the end of each list.

In several respects, the order of the topics deviates from that which is usually followed. These innovations, made in accordance with sound pedagogical principles, will arouse and sustain a greater interest in the science. The method of presentation also is unique. The principles are developed by appropriate questions designed to lead the student to infer and apprehend clearly the truth that is presented; these are followed, first, by a brief, yet clear and complete statement of the principles, and then by full and rigorous proofs of the principles. Thus the natural method of mathematical teaching has been followed, the student being led, first, to make proper inferences; second, to express the inferences briefly and accurately; and, third, to prove their truth by the method of deductive reasoning.

The acknowledgments of the author are due to Prof. J. H. Tanner, of the Department of Mathematics, Cornell University, for many valuable suggestions in connection with the preparation of this book.

WILLIAM J. MILNE.

STATE NORMAL COLLEGE, Albany, N.Y.

CONTENTS

	PAGE
ALGEBRAIC SOLUTIONS	7
DEFINITIONS AND NOTATION	12
ADDITION	21
SUBTRACTION	32
MULTIPLICATION	45
DIVISION	71
REVIEW	87
FACTORING	90
REVIEW OF FACTORING	108
HIGHEST COMMON DIVISOR	115
LOWEST COMMON MULTIPLE	126
FRACTIONS	133
REVIEW OF FRACTIONS	160
SIMPLE EQUATIONS	163
SIMULTANEOUS SIMPLE EQUATIONS	186
INVOLUTION	214
EVOLUTION	221
THEORY OF EXPONENTS	239
RADICALS	250
REVIEW	274
QUADRATIC EQUATIONS	278
GENERAL REVIEW	316
RATIO AND PROPORTION	325
VARIATION	338

	PAGE
PROGRESSIONS	344
IMAGINARY AND COMPLEX NUMBERS	363
INEQUALITIES	372
VARIABLES AND LIMITS	378
INTERPRETATION OF RESULTS	387
INDETERMINATE EQUATIONS	394
THE BINOMIAL THEOREM	398
LOGARITHMS	405
UNDETERMINED COEFFICIENTS	422
PERMUTATIONS AND COMBINATIONS	435

ACADEMIC ALGEBRA



ALGEBRAIC SOLUTIONS

1. PROBLEM 1. A man had 400 acres of corn and oats. If there were 3 times as many acres of corn as of oats, how many acres were there of each?

ARITHMETICAL SOLUTION

A certain number = the number of acres of oats.
Then, 3 times that number = the number of acres of corn,
and 4 times that number = the number of acres of both ;
therefore, 4 times that number = 400.
Hence, the number = 100, the number of acres of oats,
and 3 times the number = 300, the number of acres of corn.

ALGEBRAIC SOLUTION

Let x = the number of acres of oats.
Then, $3x$ = the number of acres of corn,
and $4x$ = the number of acres of both ;
therefore, $4x = 400$.
Hence, $x = 100$, the number of acres of oats,
and $3x = 300$, the number of acres of corn.

2. An expression of equality between two numbers or quantities is called an **Equation**.

$5x = 30$ is an equation.

3. A question that can be answered only after a course of reasoning is called a **Problem**.

4. The process of finding the result sought is called the **Solution** of the problem.

5. The expression in algebraic language of the conditions of a problem is called the **Statement** of the problem.

Solve algebraically the following problems:

2. A horse and saddle cost \$50. If the horse cost 4 times as much as the saddle, what was the cost of each?

3. A bicycle and suit cost \$90. How much did each cost, if the bicycle cost 5 times as much as the suit?

4. Of 240 stamps that Harry and his sister collected, Harry collected 3 times as many as his sister. How many did each collect?

5. If Mr. Brown and his son together had \$220, and Mr. Brown had 10 times as much as his son, how much money had each?

6. In a room containing 45 students there were twice as many girls as boys. How many were there of each?

7. A had 7 times as many sheep as B, and both together had 608. How many sheep had each?

8. A and B began business with a capital of \$7500. If A furnished half as much capital as B, how much did each furnish?

SUGGESTION. — Let x = the number of dollars A furnished.

9. A man bought a cow and a calf for \$36, paying 8 times as much for the cow as for the calf. What was the cost of each?

10. James sold his pony and a saddle for \$60. If the saddle sold for $\frac{1}{3}$ as much as the pony, what was the selling price of each?

11. A certain number added to twice itself equals 96. What is the number?

12. A farmer raised a certain number of bushels of wheat, 4 times as much corn, and 3 times as much barley. If there were in all 4000 bushels of grain, how many bushels of each kind did he raise?

13. A boy bought a bat, a ball, and a glove for \$2.25. If the bat cost twice as much as the ball, and the glove cost 3 times as much as the bat, what was the cost of each?

14. In a fire B lost twice as much as A, and C lost 3 times as much as A. If their combined loss was \$ 300, what was the loss of each ?

15. A house and lot cost \$ 3000. If the house cost 4 times as much as the lot, what was the cost of each ?

16. In a business enterprise the joint capital of A, B, and C was \$ 2100. If A's capital was twice B's, and B's was twice C's, what was the capital of each ?

17. John, William, and George together had 120 marbles. If William had twice as many as John, and George had 3 times as many as John, how many had each ?

18. In an orchard of apple, pear, and cherry trees, containing 1690 trees in all, there were 4 times as many cherry trees as pear trees, and twice as many apple trees as cherry trees. How many trees were there of each kind ?

19. A number plus itself, plus twice itself, plus 4 times itself, is equal to 72. What is the number ?

20. Charles is twice as old as his younger brother, and half as old as his older brother. If the sum of the ages of the three brothers is 28 years, what is the age of each ?

21. A farmer had twice as many sheep as horses, and twice as many hogs as sheep and horses together. If there were in all 360 animals, how many were there of each kind ?

22. A tract of land containing 640 acres was divided into three farms, such that the first was 3 times as large as the second, and the third 4 times as large as the first. How many acres did each farm contain ?

23. Three boys divided 160 marbles among themselves so that one of them received twice as many as each of the others. How many did each receive ?

24. Divide 30 into two parts, one of which is 14 times the other.

25. Divide 18 into three parts, such that the first is twice the third, and the second is 3 times the third.

26. Divide 21 into three parts, such that the first is twice the second, and the second is twice the third.

27. Divide 36 into three parts, such that the first is twice the second, and the third is equal to twice the sum of the first and second.

28. Three newsboys sold 60 papers. If the first sold twice as many as the second, and the third sold 3 times as many as the second, how many did each sell?

29. Henry earned a certain number of dollars per week. With 4 weeks' earnings he purchased a rifle, and with 20 weeks' earnings, a bicycle. If both together cost \$72, how much did he earn per week? How much did the rifle cost? the bicycle?

30. A man sold some ducks for 50 cents each, and the same number of geese for 75 cents each. If for all he received \$12.50, how many of each did he sell?

31. John has 5 times as much money as James. James has 24 cents less than John. How much has each?

32. A man had 675 sheep in three fields. If there were twice as many in the first field as in the second, and twice as many in the third field as in both of the others, how many sheep were there in each field?

33. A man bequeathed to his daughter twice as much money as to his son, and to his wife 3 times as much as to his daughter. If all together received \$9000, how much did each receive?

34. A plumber and two helpers together earned \$7.50 per day. How much did each earn per day, if the plumber earned 4 times as much as each helper?

35. What number added to $\frac{2}{3}$ of itself equals 20?

SOLUTION

Let

x = the number.

Then,

$$x + \frac{2}{3}x = 20,$$

$$\frac{5}{3}x = 20,$$

$$\frac{1}{3}x = 4.$$

Therefore,

$$x = 12, \text{ the number.}$$

36. If $\frac{1}{2}$ of a number is added to the number, the sum is 12. What is the number?

37. If $\frac{1}{3}$ of a number is added to twice the number, the sum is 35. What is the number?

38. The difference between 4 times a certain number and $\frac{1}{4}$ of the number is 30. What is the number?

39. The difference between $\frac{3}{4}$ of a certain number and $\frac{2}{3}$ of it is 16. What is the number?

40. After spending $\frac{1}{2}$ of my money and losing $\frac{1}{3}$ of it, I had \$30. How much had I at first?

41. The difference between twice a certain number and $\frac{3}{4}$ of it is 20. What is the number?

42. The number 150 can be divided into two parts, one of which is $\frac{2}{3}$ of the other. What are the parts?

43. One part of 45 is $\frac{5}{4}$ of the other. What are the parts?

44. Find two parts of 30 such that one is $\frac{7}{8}$ of the other.

45. To A, B, and C I owe in all \$93. If I owe A $\frac{2}{3}$ as much as C, and B $\frac{2}{3}$ as much as C, how much do I owe each?

46. The length of a field is $1\frac{3}{4}$ times its width, and the distance around the field is 120 rods. If the field is rectangular, what are its dimensions?

47. A, B, C, and D buy \$16,000 worth of railroad stock. How much does A take, if B takes 3 times as much as A, C twice as much as A and B together, and D $\frac{1}{3}$ as much as A, B, and C together?

48. In one season an orchard bore 650 bushels of fruit, consisting of $\frac{3}{4}$ as many bushels of pears as of peaches, and twice as many bushels of apples as of pears. How many bushels were there of each?

49. A horse, harness, and carriage cost \$340. If the horse cost 3 times as much as the harness, and the carriage cost $1\frac{1}{2}$ times as much as the horse, what was the cost of each?

DEFINITIONS AND NOTATION

6. The ideas of number and the knowledge of the processes with abstract numbers that the student has gained in arithmetic are a proper and necessary introduction to his work in algebra; but since number is discussed in a more *general* way in algebra than in arithmetic, many arithmetical processes, terms, and symbols, as 'addition,' 'subtraction,' 'greater,' 'less,' 'exponent,' '+,' '—,' etc., must now be *extended* in meaning and application.

For example, in an arithmetical sense 8 cannot be subtracted from 5, nor does $8^{\frac{1}{2}}$ have any meaning; but in an algebraic sense, as will be shown hereafter, 8 can be subtracted from 5 and $8^{\frac{1}{2}}$ is as intelligible as 8^2 .

Indeed, the processes and principles of arithmetic are but special cases of the more fundamental processes and principles of algebra.

7. A unit or an aggregate of units is called a **Whole Number**, or an **Integer**.

One of the equal parts of a unit or an aggregate of equal parts of a unit is called a **Fractional Number**.

Such numbers are called *Arithmetical*, or *Absolute Numbers*.

8. Arithmetical numbers have fixed and known values, and are represented by symbols called **numerals**; as 1, 2, 3, etc., Arabic *figures*, and I, V, X, etc., Roman *letters*.

9. It is often convenient, in solving a problem, to employ letters, such as x , y , z , to represent the numbers whose values are sought; and, in stating a rule, to employ letters to represent the numbers that must be given whenever the rule is applied.

Numbers represented by letters are called **Liter al Numbers**.

For example, the volume of any rectangular prism is equal to the area of the base multiplied by the height. By using v for volume, a for area of base, and h for height, this rule is abbreviated to

$$v = a \times h.$$

When $a = 60$ and $h = 5$, $v = 60 \times 5 = 300$;
 when $a = 36$ and $h = 10$, $v = 36 \times 10 = 360$; etc.

In each problem to which this rule applies a and h represent *fixed, known* values, but in consequence of being used for all problems of this class, a and h represent numbers to which *any arithmetical values whatever* may be assigned. Hence, the arithmetical idea of number is extended as follows.

10. A literal number to which any value can be assigned at pleasure is called a **General Number**.

11. A number whose value is known or a number to which any value can be assigned is called a **Known Number**.

The numerals, 3 and $4\frac{1}{2}$, and the general numbers a and h in $v = a \times h$, in § 9, are known numbers.

Known literal numbers are generally represented by the *first letters* of the alphabet.

12. A number whose value is to be found is called an **Unknown Number**.

Unknown numbers are usually represented by the *last letters* of the alphabet.

ALGEBRAIC SIGNS

13. The **Sign of Addition** is $+$, read '*plus*.'

It indicates that the number following it is to be added to the number preceding it.

$a + b$, read '*a plus b*,' indicates that b is to be added to a .

14. The **Sign of Subtraction** is $-$, read '*minus*.'

It indicates that the number following it is to be subtracted from the number preceding it.

$a - b$, read '*a minus b*,' indicates that b is to be subtracted from a .

15. The Sign of Multiplication is \times or \cdot , read '*multiplied by.*'

It indicates that the number preceding it is to be multiplied by the number following it.

$a \times b$, or $a \cdot b$, indicates that a is to be multiplied by b .

The sign of multiplication is usually omitted in algebra, except between figures.

$a \times b$, or $a \cdot b$, may be abbreviated to ab , $x \times y$ to xy , $4 \times b$ to $4b$, etc. But 3×5 cannot be written 35, because 35 means $30 + 5$.

16. The Sign of Division is \div , read '*divided by.*'

It indicates that the number preceding it is to be divided by the number following it.

$a \div b$ indicates that a is to be divided by b .

Division may be indicated also by writing the dividend above the divisor with a line between them.

Such indicated divisions are called *Fractions*. (Cf. § 158.)

$\frac{a}{b}$ indicates that a is to be divided by b .

17. The Sign of Equality is $=$, read '*is equal to*' or '*equals.*'

18. The Sign of Inequality is $>$ or $<$.

When used between two numbers, it signifies that they are unequal, the greater number being at the opening of the sign.

$a > b$ is read '*a is greater than b.*'

$x < 5$ is read '*x is less than 5.*'

19. The Signs of Aggregation are: the *Parenthesis*, $()$; the *Vinculum*, $-$; the *Brackets*, $[\]$; the *Braces*, $\{ \}$; and the *Vertical Bar*, $|$.

They show that the expressions included by them are to be treated as single numbers.

Thus, each of the forms $(a + b)c$, $\overline{a + b} \cdot c$, $[a + b]c$, $\{a + b\}c$, and $\left. \begin{array}{l} a \\ + b \end{array} \right| c$, signifies that the sum of a and b is to be multiplied by c .

When numbers are included by any of the signs of aggregation, they are commonly said to be *in parenthesis*.

20. The Sign of Continuation is \dots or $----$, read '*and so on,*' or '*and so on to.*'

$2, 4, 6, 8, ----$ is read '*2, 4, 6, 8, and so on.*'

21. The Sign of Deduction is \therefore . It signifies *therefore* or *hence*.

FACTORS, POWERS, AND ROOTS

22. Each of two or more numbers which multiplied together produce a given number is called a **Factor** of the number.

Since $12 = 2 \times 6$, or 4×3 , each of these numbers is a factor of 12.

Since $3ab = 3 \times a \times b$, each of these numbers is a factor of $3ab$.

23. When a factor of a number is considered as the multiplier of the remaining factor, it is called a **Coefficient** of that factor.

In $7x$, $5ax$, bxy , and $(a-b)x$, the coefficients of x are 7, $5a$, by , and $(a-b)$; in bxy , bx is the coefficient of y .

Coefficients are *Numerical*, *Literal*, or *Mixed*, according as they are composed of *figures*, *letters*, or both *figures and letters*.

When no numerical coefficient is expressed, the coefficient may be considered to be 1.

24. When a number is used a certain number of times as a factor, the product is called a **Power** of the number.

Powers are *named* from the number of times the number is used as a factor.

When a is used *twice* as a factor, the product is the *second* power of a ; when a is used *three* times as a factor, the product is the *third* power of a ; four times, the *fourth* power of a ; n times, that is, any number of times, the *nth* power of a .

The *second* power is also called the *square*, and the *third* power the *cube*.

The product indicated by $a \times a \times a \times a \times a$ may be more briefly indicated by a^5 . Likewise, if a is to be used n times as a factor, the product may be indicated by a^n .

25. A figure or letter placed a little above and to the right of a number is called an **Index** or an **Exponent** of the power thus indicated.

The integers that the student has been using in arithmetic have been positive integers.

When the exponent is a positive integer, it indicates the number of times that the number is to be used as a factor.

5^2 indicates that 5 is to be used twice as a factor; a^3 indicates that a is to be used 3 times as a factor.

When no exponent is written, the exponent is regarded as 1.

5 is regarded as the first power of 5, and a^1 is usually written a .

The terms coefficient and exponent should be carefully distinguished.

Thus, $5a = a + a + a + a + a$, but $a^5 = a \times a \times a \times a \times a$.

26. One of the equal factors of a number is called a **Root** of the number.

5 is a root of 25; a is a root of a^4 ; $4x$ is a root of $64x^3$.

Roots are *named* from the number of equal factors into which the number is separated.

One of the *two* equal factors of a number is its *second* root; one of the *three* equal factors of a number is its *third* root; one of the *four* equal factors, the *fourth* root; one of the n equal factors, the n th root.

The *second* root of a number is also called its *square* root, and its *third* root is called its *cube* root.

27. The symbol which denotes that a root of a number is sought is $\sqrt{}$, written before the number.

It is called the **Root Sign**, or the **Radical Sign**.

A figure or letter written in the opening of the radical sign indicates what root of the number is sought.

It is called the **Index** of the root.

When no index is written, the second, or square root is meant.

$\sqrt[3]{8}$ indicates that the third, or cube root of 8 is sought.

\sqrt{ax} and $\sqrt{a-b}$ indicate the square roots of ax and $a-b$, respectively.

The horizontal line used in connection with the radical sign is a vinculum.

ALGEBRAIC EXPRESSIONS

28. A number expressed by algebraic symbols is called an **Algebraic Expression**.

29. When signs of operation are employed in algebraic expressions, the sequence of operations is determined by the following conventional law:

A series of additions and subtractions or of multiplications and divisions are performed in order from left to right.

$$3 + 4 - 2 + 3 = 7 - 2 + 3 = 5 + 3 = 8.$$

$$3 \times 4 \div 2 \times 3 = 12 \div 2 \times 3 = 6 \times 3 = 18.$$

$a + b - c + d$ indicates that b is to be added to a , then from this result c is to be subtracted, and to the result just obtained d is to be added.

30. When a particular number takes the place of a letter or general number, the process is called **Substitution**.

NUMERICAL SUBSTITUTIONS

1. When $a = 2$, $b = 3$, and $c = 5$, what are the numerical values of $3c$, c^3 , $\sqrt{8ab^2}$, $a^2 + b^2$, and $(a + b)^2$, respectively?

SOLUTIONS

$$3c = 3 \cdot 5 = 15.$$

$$c^3 = 5 \cdot 5 \cdot 5 = 125.$$

$$\sqrt{8ab^2} = \sqrt{8 \cdot 2 \cdot 3 \cdot 3} = \sqrt{2 \cdot 2 \times 2 \cdot 2 \times 3 \cdot 3} = 2 \times 2 \times 3 = 12.$$

$$a^2 + b^2 = 2 \cdot 2 + 3 \cdot 3 = 4 + 9 = 13.$$

$$(a + b)^2 = (a + b)(a + b) = (2 + 3)(2 + 3) = 5 \cdot 5 = 25.$$

Find the numerical value of each of the following algebraic expressions, when $a = 5$, $b = 3$, $c = 10$, $m = 4$, $n = 1$:

2. $10a$.

11. $(ab)^2$.

19. $\sqrt[3]{4ac^2m}$.

3. $2ab$.

12. a^2b^2 .

20. $\frac{c + 2m}{c - 2m}$.

4. $3cm$.

13. $\sqrt{2acn}$.

5. $6bc$.

14. $3b^2cn^2$.

21. $c + \frac{2m}{c - 2m}$.

6. $5cm^2$.

15. $a^2 - b^2$.

7. $2a^2b$.

16. $(a - b)^2$.

22. a^3c .

8. $3bm^2$.

17. $(n + 1)^4$.

23. m^{a-b} .

10. am^4 .

18. $n^5 + 1$.

24. $(bm)^{a-b}$.

31. An algebraic expression whose parts are not separated by + or - is called a **Term**; as $2x^2$, $-5xyz$, and $\frac{xy}{z}$.

In the expression $2x^2 - 5xyz + \frac{xy}{z}$ there are three terms.

The expression $m(a + b)$ is a term, the parts being m and $(a + b)$.

32. Terms that contain the same letters with the same exponents are called **Similar Terms**.

$3x^2$ and $12x^2$ are similar terms; also $3(a + b)^2$ and $12(a + b)^2$; also ax and bz , regarding a and b as the coefficients of x .

33. Terms that contain different letters, or the same letters with different exponents, are called **Dissimilar Terms**.

$5a$ and $3by$ are dissimilar terms; also $3a^2b$ and $3ab^2$.

34. Each literal factor of a term is called a **Dimension** of the term.

The number of literal factors or dimensions of a term indicates its *Degree*.

$abcd$ is a term of the fourth degree, because it is composed of four literal factors, or has four dimensions. x^3 is a term of the third degree, since x^3 is a convenient way of indicating that x is taken three times as a factor. The expressions $4x^2yz^3$ and $5xyz^4$ are each of the sixth degree.

35. The term of highest degree in an expression determines the **Degree of the Expression**.

$x^3 + 3x^2 + x + 2$ and $abc + b^2c + ac - b$ are expressions of the third degree.

36. When all the terms of an expression are of the same degree, the expression is called a **Homogeneous Expression**.

$x^3 + 3x^2y + xy^2 + 2y^3$ and $abc + b^2c + ac^2$ are each homogeneous expressions.

37. An algebraic expression of one term only is called a **Monomial**, or a **Simple Expression**.

xy and $3ab$ are monomials.

38. An algebraic expression of more than one term is called a **Polynomial**, or a **Compound Expression**.

$3a + 2b$, $xy + yz + zx$, and $a^2 + b^2 - c^2 + 2ab$ are polynomials.

39. A polynomial of two terms is called a **Binomial**.

$3a + 2b$ and $x^2 - y^2$ are binomials.

40. A polynomial of three terms is called a **Trinomial**.

$a + b + c$ and $3x - 2y - z$ are trinomials.

41. An expression, any term of which is a fraction, is called a **Fractional Expression**.

$\frac{2x^2}{a^2} - 3x + \frac{a}{x}$ is a fractional expression.

42. An expression that contains no fraction is called an **Integral Expression**.

$5a^2 - 2a$ and $6x$ are integral expressions.

Expressions like $x^2 + \frac{1}{4}x^2 + \frac{1}{4}x + 1$ are sometimes regarded as integral, since the literal numbers are not in fractional form.

43. An expression that can be written without using a root sign is called a **Rational Expression**.

1, 2, 3, ..., $a + \frac{1}{2}$, $\frac{x^2}{x-y}$, and $(a-b)^2$ are rational expressions.

$\sqrt{25}$ is rational, since it can be written 5 without a root sign.


44. An expression that cannot be written without using a root sign is called an **Irrational Expression**.

$a + \sqrt{b}$, $a + 2\sqrt{a} + 1$, and $\sqrt[3]{4}$ are irrational expressions.

$\sqrt{a^2}$ is not irrational, however, since it may be written a .

POSITIVE AND NEGATIVE NUMBERS

45. For convenience, arithmetical numbers may be arranged in an ascending scale:



The operations of addition and subtraction are thus reduced to counting along a scale of numbers. 2 is added to 3 by beginning at 3 in the scale and counting 2 units in the ascending, or additive direction; and consequently, 2 is subtracted from 3 by beginning at 3 and counting 2 units in the descending, or subtractive direction. In the same way 3 is subtracted from 3. But if we attempt to subtract 4 from 3, we discover that the operation of subtraction is restricted in arithmetic, inasmuch as a greater number cannot be subtracted from a less. If this restriction held in algebra, it would be impossible to subtract one literal number from another without taking into account their arithmetical values. Therefore, this restriction must be removed in order to proceed with the discussion of numbers.

To subtract 4 from 3 we begin at 3 and count 4 units in the descending direction, arriving at 1 on the opposite, or subtractive side of 0. It now becomes necessary to extend the scale 1 unit in the subtractive direction from 0.

To subtract 5 from 3 we begin at 3 and count 5 units in the descending direction, arriving at 2 on the opposite, or subtractive side of 0. The scale is again extended, and may be extended indefinitely in the subtractive direction in a similar way.

For convenience, numbers on opposite sides of 0 are distinguished by means of the small signs + and -, called *signs of quality*, or *direction signs*, + being prefixed to those which stand in the additive direction from 0 and - to those which stand in the subtractive direction from 0.

The former are called **Positive Numbers**, the latter **Negative Numbers**.

Hence, the scale of algebraic numbers may be written :

..., -5, -4, -3, -2, -1, 0, +1, +2, +3, +4, +5, ...

46. By repeating +1 as a unit any positive number may be obtained, and by repeating -1 as a unit any negative number may be obtained. Hence, positive numbers are measured by the **positive unit**, +1, and negative numbers by the **negative unit**, -1, or by parts of these units.

47. If +1 and -1, or +2 and -2, or any two numbers numerically equal but opposite in quality are taken together, they cancel each other. For counting any number of units from 0 in either direction and then counting an equal number of units from the result in the opposite direction, we arrive at 0. Hence,

If a positive and a negative number are united into one number, any number of units or parts of units of one cancels an equal number of units or parts of units of the other.

48. Two concrete quantities of the same kind are sometimes opposed to each other in some sense so that, if united, any number of units of one cancels an equal number of units of the other. For convenience, such quantities are often distinguished as *positive* and *negative*.

If money *gained* is *positive*, money *lost* is *negative*, for any sum gained is canceled by an equal sum lost. If a *rise* in temperature is *positive*, a *fall* in temperature is *negative*. If distances *north* or *west* or *upstream* are *positive*, distances *south* or *east* or *downstream* are *negative*.

ADDITION

49. 1. If a man has 10 dollars in one pocket and 15 dollars in another, how much money has he?

2. If in algebra money in hand is considered a positive quantity, indicate his financial condition algebraically. What is the sum of 10 positive units and 15 positive units, that is, of +10 and +15? of +4 and +8? of $+a$ and $+b$?

3. If a person owes one man 10 dollars and another 15 dollars, how much does he owe both? Indicate his financial condition algebraically, regarding a debt as a negative quantity.

4. What is the sum of 10 negative units and 15 negative units, that is, of -10 and -15? of -6 and -14? of $-a$ and $-b$?

5. What sign has the sum of two algebraic numbers that have like signs?

6. If a man has 25 dollars and owes 15 dollars, how much of his money will be required to cancel the debt? How many dollars will he have after settlement?

7. What is the result when -15 is united with +25, that is, what is the *algebraic sum* of -15 and +25? of -20 and +10? of +8 and -3? of +6 and -10?

50. The aggregate value of two or more algebraic numbers is called their **Algebraic Sum**.

The process of finding the simplest expression for the algebraic sum of two or more numbers is called **Addition**.

51. PRINCIPLES. — 1. *The algebraic sum of two numbers with like signs is equal to the sum of their absolute values with the common sign prefixed.*

2. *The algebraic sum of two numbers with unlike signs is equal to the difference between their absolute values with the sign of the numerically greater prefixed.*

By successive applications of the above principles any number of numbers may be added.

Only similar terms can be united into a single term.

Principle 1 may be established as follows :

The sum of 5 positive units and 3 positive units is evidently $(5 + 3)$ positive units, or 8 positive units ; that is,

$$+5 + +3 = +(5 + 3) = +8.$$

Similarly, whatever absolute values a and b represent, since a times the unit $+1$ plus b times the unit $+1$ is equal to $(a + b)$ times the unit $+1$,

§ 46, $+a + +b = +(a + b).$

Again, the sum of 5 negative units and 3 negative units is $(5 + 3)$ negative units, or 8 negative units ; that is,

$$-5 + -3 = -(5 + 3) = -8.$$

Similarly, whatever absolute values a and b represent, since a times the unit -1 plus b times the unit -1 is equal to $(a + b)$ times the unit -1 ,

§ 46, $-a + -b = -(a + b).$

Principle 2 may be established as follows :

The sum of 5 positive units and 3 negative units is 2 positive units, since, § 47, the 3 negative units cancel 3 of the positive units and leave 2 positive units ; that is,

$$+5 + -3 = +(5 - 3) = +2.$$

The sum of 5 positive units and 7 negative units is 2 negative units, since, § 47, the 5 positive units cancel 5 of the negative units and leave 2 negative units ; that is,

$$+5 + -7 = -(7 - 5) = -2.$$

Similarly, whatever absolute values a and b represent,

if $a > b$, $+a + -b = +(a - b),$

for, § 47, the b negative units will cancel b of the a positive units and leave $(a - b)$ positive units ;

but if $b > a$, $+a + -b = -(b - a),$

for, § 47, the a positive units will cancel a of the b negative units and leave $(b - a)$ negative units.

EXAMPLES

Find the value of

- | | | |
|---------------|---------------|------------------------------|
| 1. $+7 + +3.$ | 3. $+7 + -3.$ | 5. $-3 + +4 + -2 + +8 + -9.$ |
| 2. $-7 + -3.$ | 4. $-7 + +3.$ | 6. $+m + -n$, if $m > n.$ |

52. To conform with the ideas already presented, the terms 'greater' and 'less' must be interpreted as follows:

An algebraic number is increased, or made *greater*, when a positive number is added to it, and decreased, or made *less*, when a negative number is added to it.

Since, by § 51, $-3 + +1 = -2$, $-2 + +1 = -1$, $-1 + +1 = 0$, $+1 + +1 = +2$, etc., in the scale of algebraic numbers

$$\dots, -5, -4, -3, -2, -1, 0, +1, +2, +3, +4, +5, \dots,$$

each number is greater than the number on its left and less than the number on its right; that is,

$$\dots, -3 < -2, -2 < -1, -1 < 0, 0 < +1, +1 < +2, +2 < +3, \dots$$

NOTE. $-3 < -2$ may be read '3 is less than 2' or '-2 is greater than -3.'

Hence, it follows that:

1. *Any positive number is greater than zero and any negative number is less than zero.*

2. *Of two positive numbers that which has the greater absolute value is the greater, and of two negative numbers that which has the less absolute value is the greater.*

53. Abbreviated notation for addition.

Referring to the scale of algebraic numbers, it is evident that adding positive units to any number is equivalent to counting them in the positive direction from that number, and adding negative units to any number is equivalent to counting them in the negative direction from that number. Hence, in addition, the signs $+$ and $-$ denoting quality have primarily the same meanings as the signs $+$ and $-$ denoting arithmetical addition and subtraction. For example, by the definition of positive and negative numbers,

$$+1 \text{ means } 0 + 1 \text{ and } -1 \text{ means } 0 - 1;$$

also $+5 \text{ means } 0 + 5 \text{ and } -5 \text{ means } 0 - 5; \text{ etc.}$

Hence, in addition, but one set of signs, $+$ and $-$, is necessary, and in finding the sum of any given numbers, the signs $+$ and $-$ may be regarded either as signs of quality or as signs of operation, though it is commonly preferable to regard them as signs of operation.

For brevity, it is customary to omit the sign $+$ before a monomial or before the first term of a polynomial. But the sign $-$ cannot be omitted.

$+5 + +3 + -6$ is written $5 + 3 - 6$; $-4 + +8 + -2$ is written $-4 + 8 - 2$.

When there is need of distinguishing between the signs of quality $+$ and $-$ and the signs of operation $+$ and $-$, the numbers and their signs of quality may be inclosed in parentheses.

Thus, if $a = +5$, $b = -3$, and $c = -2$, then $a + b + c = (+5) + (-3) + (-2)$; $a - b - c = (+5) - (-3) - (-2)$; $abc = (+5)(-3)(-2)$; etc.

54. A term preceded by $+$, expressed or understood, is called a **Positive Term**, and a term preceded by $-$, a **Negative Term**.

Thus, in the polynomial $3a + 2b - 5c$ the first and second terms are positive and the third term is negative.

EXAMPLES

Write the following with one set of signs:

- | | | |
|----------------|----------------------|---------------------|
| 1. $+7 + +8$. | 4. $+10 + -2 + -4$. | 7. $+a + -b$. |
| 2. $+6 + -5$. | 5. $-6 + -3 + +16$. | 8. $-a + +b + -c$. |
| 3. $-3 + -7$. | 6. $+8 + +4 + -5$. | 9. $-x + -y + -z$. |

55. 1. How does $5 + 3 - 2$ compare in value with $5 - 2 + 3$, or with $3 - 2 + 5$, or with $-2 + 3 + 5$?

2. How does $a + b - c$ compare in value with $b - c + a$, or with $b + a - c$, or with $a - c + b$?

3. In what order may numbers be added?

Law of Order, or Commutative Law for Addition. — *Algebraic numbers may be added in any order.*

The Law of Order may be established as follows:

We know from arithmetic that arithmetical numbers may be added in any order. Since, § 51, Prin. 1, algebraic numbers having like signs are added by prefixing their common sign to the sum of their absolute, or arithmetical values, and since in finding this sum the absolute values may be added in any order, it follows that algebraic numbers having like signs may be added in any order.

If some of the numbers are positive and some are negative, § 47, the same number of positive and negative units will cancel each other, and the same number of one or the other will be left, in whatever order the numbers are added.

Hence, whether the numbers have like or unlike signs, they may be added in any order; that is,

$$a + b + c = b + c + a = c + a + b = c + b + a, \text{ etc.},$$

for all values of a , b , and c .

56. 1. How are the numbers 4, $\frac{1}{2}$, 2, and $\frac{3}{4}$ grouped in adding? the numbers 25 and 32, or $20 + 5$ and $30 + 2$?

2. In what manner may the terms of an expression be grouped in addition?

Law of Grouping, or Associative Law for Addition. — *The sum of three or more algebraic numbers is the same in whatever manner the numbers are grouped.*

The Law of Grouping may be established as follows:

By the Law of Order, $a + b + c = b + c + a$

$$\S 29, \quad \quad \quad = (b + c) + a$$

$$\text{by the Law of Order,} \quad \quad \quad = a + (b + c).$$

Other cases, as $a + b + c = (a + c) + b$, etc., may be proved similarly.

Hence, for all values of a , b , and c ,

$$a + b + c = a + (b + c) = (a + c) + b = c + (a + b), \text{ etc.}$$

57. To add similar monomials.

EXAMPLES

1. Add $4a$ and $3a$.

PROCESS **EXPLANATION.** — Just as $4 = 1 + 1 + 1 + 1$, so $4a = a + a + a + a$; just as $3 = 1 + 1 + 1$, so $3a = a + a + a$. Therefore, $4a + 3a = a + a + a + a + a + a + a$, the symbol for which is $7a$.

$$\begin{array}{r} 4a \\ 3a \\ \hline 7a \end{array}$$
 Or, the sum may be obtained by adding the numerical coefficients and annexing to their sum the common literal part.

2. Add $4a$, $\frac{3}{2}a$, $-3a$, and $\frac{1}{2}a$.

PROCESS

$$4a + \frac{3}{2}a - 3a + \frac{1}{2}a = 4a - 3a + (\frac{3}{2}a + \frac{1}{2}a) = a + 2a = 3a.$$

EXPLANATION. — By the Law of Grouping the sum of $\frac{3}{2}a$ and $\frac{1}{2}a$ may be added to the sum of $4a$ and $-3a$. Just as $4 = 1 + 1 + 1 + 1$, so $4a = a + a + a + a$; just as $-3 = -1 - 1 - 1$, so $-3a = -a - a - a$. Therefore, $4a - 3a = a + a + a + a - a - a - a = 0 + a = a$. Just as $\frac{3}{2} = \frac{1}{2} + \frac{1}{2} + \frac{1}{2}$, so $\frac{3}{2}a = \frac{1}{2}a + \frac{1}{2}a + \frac{1}{2}a$. Therefore, $\frac{3}{2}a + \frac{1}{2}a = \frac{1}{2}a + \frac{1}{2}a + \frac{1}{2}a + \frac{1}{2}a = \frac{4}{2}a = 2a$. Adding the two groups, $a + 2a = a + a + a = 3a$.

Or, the sum may be obtained by adding in any order or manner the numerical coefficients, and annexing to their sum the common literal part.

Simplify the following:

3. $2y - 7y - 5y - y + 10y - 6y + 8y$.
4. $5a - 3a + 8a - 10a - 5a - 11a + 24a$.
5. $3by - 5by - 10by - 14by + 48by$.
6. $8a^3b + 6a^3b - 11a^3b - 2a^3b + 9a^3b$.
7. $1\frac{4}{5}x^2y^2 - \frac{1}{2}x^2y^2 - 1\frac{7}{10}x^2y^2 + 3\frac{1}{2}x^2y^2 + x^2y^2$.
8. $5(xy)^2 - 3(xy)^2 - 15(xy)^2 + 4(xy)^2 + 13(xy)^2$.
9. $(a - x) + 5(a - x) + 7(a - x) - 3(a - x) - 2(a - x)$.
10. $3(a + b)^2 + 6(a + b)^2 - 10(a + b)^2 - (a + b)^2 + 12(a + b)^2$.
11. $20\sqrt{x-3} - 8\sqrt{x-3} - 12\sqrt{x-3} + \sqrt{x-3} + 7\sqrt{x-3}$.
12. $3x(x^2 - 2x + 3) - x(x^2 - 2x + 3) + 2x(x^2 - 2x + 3)$.
13. $2(x - 1) - 13(x - 1) + 5(x - 1) + 10(x - 1) + 6(x - 1)$.
14. $\frac{1}{2}(a + b - c) - \frac{4}{3}(a + b - c) + \frac{1}{2}\frac{1}{6}(a + b - c)$.

Since only similar terms can be united into a single term, in algebra dissimilar terms are considered to have been added when they have been written in succession with their proper signs.

In algebra many indicated operations are regarded as performed.

Since $5a$, $-3b$, and $2c$ cannot be united into a single term, their sum is written $5a - 3b + 2c$.

15. Add $6a$, $-5b$, $-3a$, $3b$, $2c$, and $-a$.

SOLUTION. $6a - 5b - 3a + 3b + 2c - a = 2a - 2b + 2c$.

Add the following:

16. $2xy$, $4ab$, $3xy$, and ab .
17. mn , $-3cd$, $-6mn$, and $4cd$.
18. a , $-b$, $2c$, $-2a$, $3b$, and $-4c$.
19. $6x$, $3y$, $-2x$, y , $-3x$, z , and $-3y$.
20. $2a$, $2b$, $2c$, $2d$, $-a$, $-3b$, $-c$, and $-3d$.
21. a , $-4a$, $2b$, cd , $-2ab$, $5b$, and $-3cd$.

58. To add polynomials.

EXAMPLES

1. Add $3a - 3b + 5c$, $-3a + 2b$, and $c - 4b + 2a$.

PROCESS	EXPLANATION.
$3a - 3b + 5c$	For convenience in adding, similar terms are written in the same column.
$-3a + 2b$	The algebraic sum of the first column is $2a$, of the second $-5b$, and of the third $+6c$; and these
$2a - 4b + c$	numbers written in succession express in its simplest form the sum sought.
<hr/>	
$2a - 5b + 6c$	

2. Simplify $11a^2b - 7ab^2 + 2ac^2 + 10ab - 4ac^2 + 5a^2b - 4ab^2 + 5ac^2 + b^3 + 9ab^2 - 7a^2b - 2b^3 + 2ab^2 - 8ab - 6a^2b$.

PROCESS

$$\begin{array}{r}
 11a^2b - 7ab^2 + 2ac^2 + 10ab + b^3 \\
 + 5a^2b - 4ab^2 - 4ac^2 \quad - 2b^3 \\
 - 7a^2b + 9ab^2 + 5ac^2 \\
 - 6a^2b + 2ab^2 \quad - 8ab \\
 \hline
 3a^2b \quad + 3ac^2 + 2ab - b^3
 \end{array}$$

RULE. — Arrange the terms so that similar terms stand in the same column.

Find the algebraic sum of each column, and write the results in succession with their proper signs.

3. Add $2a - 3b$, $2b - 3c$, $5c - 4a$, $10a - 5b$, and $7b - 3c$.

4. Add $x + y + z$, $x - y + z$, $y - z - x$, $z - x - y$, and $x - z$.

Simplify the following polynomials:

5. $7x - 11y + 4z - 7z + 11x - 4y + 7y - 11z - 4x + y - x - z$.

6. $a + 3b + 5c - 6a + d + 4b - 2c - 2b + 5a - d + a - b$.

7. $4x^2 - 3xy + 5y^2 + 10xy - 17y^2 - 11x^2 - 5xy + 12x^2 - 2xy$.

8. $2xy - 5y^2 + x^2y^2 - 7xy + 3y^2 - 4x^2y^2 + 5xy + 4y^2 + x^2y^2$.

9. $2ay - 3ac - 4ay + 4ac - 6ay + 5ac + 11ay - 4ac - ay$.

10. $5am - 3a^2m^2 + 4 - 4am + a^2m^2 - 2 + 5 + a^2m^2 - 6 + 3am$.
11. $6\sqrt{x} - 5\sqrt{xy} + 3\sqrt{y} - 4\sqrt{x} + 6\sqrt{xy} - \sqrt{x} - \sqrt{y} + 3\sqrt{y} - 2\sqrt{xy}$.

Add the following polynomials:

12. $7a - 3b + 5c - 10d$, $2b + d - 3c - 4e$, $5c - 6a + 2d - 4e$, $8b - 7a - 8c - e$, $a - 5c + 5d + 11e$, $a - b + c + 2d + e$, and $5a - 4b + 2c$.

13. $5x - 3y - 2z$, $4y - 2x + 6z$, $3a - 2x - 4y$, $4b - 2z - 5y$, $a - 5b$, $5y - 6x$, $8x + 2y - 5a - 2b$, and $6x - y - 2z + 4b$.

14. $m + n - \sqrt{mn}$, $\sqrt{mn} - 2m - 3n$, $3m + 2n$, $4n - \sqrt{mn} - 3m$, $5\sqrt{mn} - n$, $4m - n - 2\sqrt{mn}$, $5n + 2m - 3\sqrt{mn}$, and $n - 6m$.

15. $2c - 7d + 6n$, $11m - 3c - 5n$, $7n - 2d - 8c$, $8d - 3m + 10c$, $4d - 3n - 8m$, $m - 6n$, and $2m - 3d$.

16. $4x^3 - 2x^2 - 7x + 1$, $x^3 + 3x^2 + 5x - 6$, $4x^2 - 8x^3 + 2 - 6x$, $2x^3 - 2x^2 + 8x + 4$, and $2x^3 - 3x^2 - 2x + 1$.

17. $a^5 + 5a^4b + 5ab^4 + b^5$, $a^4b - 2a^5 + a^3b^2 - 2b^5$, $a^3b^2 - 3a^2b^3 - 4a^4b - a^5$, and $2a^5 + a^4b - 2a^3b^2 + 2a^2b^3 - 3ab^4 + b^5$.

18. $a^4 - 2a^3b + 3a^2b^2$, $3ab^3 - 4b^4 - 2a^2b^2$, $3a^3b + 4a^4 - 3ab^3 + 4b^4$, $5a^3b + 7a^2b^2 - 4ab^3 - 8b^4$, and $a^4 - 6a^3b - 8a^2b^2$.

19. $5x^5 - x^3 + 7x - 9$, $4x^5 - 3x^5 + 6x^3 + 12$, $x^5 - 5x^5 - x - 7$, $4 - x^3 - x^5$, $4x^4 - 10x^3 + 3x^5 + 4$, and $x^5 + x^3 - 3x^2 - 4x - 5$.

20. $3(a + b) + 6(b + c)$, $5(a + b) - 10(b + c)$, $2(a + b) + (b + c)$, $3(b + c) - (a + b)$, $2(b + c) - 10(a + b)$, and $3(a + b) - 3(b + c)$.

21. $x + 3(a + 1) - y$, $-(a + 1) - 2x + 4y$, and $3x - 4(a + 1)$.

22. $a^3 - 3a^2bc - 6ab^2c$, $a^2b - b^3 - c^3 - 3abc$, $ab^2 + b^2c + bc^2$, $5a^2bc + 4ab^2c + c^3$, $b^3 - a^2b - ab^2$, $a^3 + b^2c + bc^2$, and $2ab^3c - 2bc^2$.

23. $.12x^3 - 4x^2 + x + 2$, $.4x^2 - 4x + .4 - x^3$, $3\frac{1}{2}x - .6 + 3x^2 + 2x^3$, and $1 - \frac{1}{2}x + 1.2x^2 + \frac{2}{3}x^3$.

24. $ax - \frac{3}{2}ax^2 - \frac{1}{4}ax^3$, $\frac{3}{2}ax^2 - \frac{1}{2}ax^3 - \frac{1}{8}bxy$, $\frac{1}{2}bxy - \frac{3}{4}ax^3 - \frac{1}{8}ab$, $\frac{2}{3}bxy - \frac{5}{8}ab + \frac{1}{4}ax$, and $2ab - \frac{3}{2}ax + \frac{5}{4}ax^2$.

EXERCISES

59. 1. If a boy has n marbles and buys 10, how many will he then have? If he gives away m of these, how many will be left?

2. Mary has 25 cents. How many cents will she have after spending 10 cents and earning a cents? If she has c cents, spends b cents, and earns a cents, how many cents will she have?

3. A boy who has p marbles loses q marbles, and then buys r marbles. How many does he then have?

4. James is 15 years old. In how many years will he be 21 years old? In how many years will he be x years old? Harry is y years old. How many years older is he than James? In how many years will Harry be x years old?

5. Edith is 14 years old. How old was she 4 years ago? a years ago? How old will she be 3 years hence? b years hence?

6. William is x years old. How old was he a year ago? How old will he be in 5 years? in a years? After how many years will he be 21 years old? m years old? How old will he be when he is twice as old as he is now?

7. In a certain family there are five children each of whom is 2 years older than the one next younger. If the youngest is x years old, what are the ages of the others?

8. A woman sold some eggs, and with the money bought 8 pounds of sugar and 5 pounds of coffee. If the sugar cost a cents a pound, and the coffee cost b cents a pound, how much did she receive for the eggs?

9. What two whole numbers are nearest to 50? to x ? to $x + 5$? If y is an even number, what are the nearest even numbers?

10. George is a years younger than Henry, and b years younger than John. If John is 16 years old, how old is Henry?

11. A man paid two men, whom he owed, in the following manner: To the first he gave an a -dollar bill, and received change amounting to b dollars; and to the second he gave a b -dollar bill, and received change amounting to c dollars. How much did he owe both?

EQUATIONS AND PROBLEMS

60. 1. Simplify the equation $2x - 3x + 6x + 5x - x = 27$, and find the value of x .

SOLUTION

$$2x - 3x + 6x + 5x - x = 27.$$

Uniting terms,

$$9x = 27.$$

Hence,

$$x = 3.$$

Simplify, and find the value of x :

2. $10x - 7x + 4x - 6x + 11x - 20x + 12x = 40.$

3. $13x - 6x - 4x + 7x + 11x - 16x + 15x = 20.$

4. $25x - 5x - 7x - 2x + 14x - 10x - 12x = 36.$

5. $17x + 2x - 6x + 4x - 12x - 30x + 40x = 75.$

6. $10x + 2x + 3x + 4x + 11x + 12x + 18x = 60.$

7. $16x - 3x - 5x - 8x + 10x + 15x - 15x = 50.$

8. $12x + 10x - 20x + 16x - 3x - 2x + 2x = 75.$

9. $14x - 11x + 26x - 35x - 4x + 7x + 4x = 16.$

10. $75x - 37x - 40x + 10x - 8x - 6x + 9x = 21.$

11. $4x + 10x - 60x + 48x + 12x + 5x + 2x = 63.$

12. $7x + 11x - 13x + 15x - 17x - 3x + 5x = 25.$

13. $5x - 15x + 25x - 30x + 10x + 3x + 6x = 56.$

14. $x + 2x + 3x + 4x + 5x + 6x + 7x + 8x = 144.$

15. $4x + 12x - 17x - 10x + 15x + x + 15x = 400.$

16. $3x + 10x - 20x - 4x + 12x + 3x + 11x = 300.$

Solve the following problems:

17. A man bequeathed \$10,000 to 3 sons and 4 daughters, so that a son received twice as much as a daughter. What was the share of each daughter, and of each son?

18. John had twice as many marbles as Henry, and $\frac{1}{3}$ as many as Charles. If they had 225 marbles in all, how many had each?

19. A had twice as much money as B, who had 3 times as much money as C. If all together had \$2000, how much money had each?

20. A merchant owes A a certain sum of money, B $\frac{1}{2}$ as much, and C twice as much as A. Various persons owe him in all 12 times as much as he owes B. If all these debts were paid, he would have \$10,000. What are the amounts he owes?

21. Mr. Jones succeeded in doubling his capital once every 5 years. If his capital at the end of 20 years was \$150,000, with what capital did he begin?

22. The distance around a rectangular field 4 times as long as it is wide is 200 rods. What are the dimensions of the field?

23. What are the dimensions of a rectangular field whose length is twice its width, if 240 rods of fence are required to inclose it?

24. Two boys caught the same number of fish, another caught 10 more, and another 10 less. If they caught in all 120 fish, how many did each catch?

25. The sum of 3 consecutive whole numbers is 84. What are the numbers?

SUGGESTION. — Let x represent the middle number. Then, the other two numbers will be represented by $x - 1$ and $x + 1$.

26. Of what 3 consecutive even numbers is 150 the sum?

27. A, B, C, and D together have \$1500. If A had \$50 more and B \$50 less, they would each have the same sum as C and $\frac{1}{2}$ as much as D. How much money has each?

28. The ages of 4 brothers differ successively by 2 years. If the sum of their ages is 56 years, what is the age of each?

29. Three newsboys sold 270 papers in an evening. If the second sold 5 less than twice as many as the first, and the third 5 more than 3 times as many as the first, how many papers did each sell?

SUBTRACTION

61. 1. What is left when $5xy$ is taken from $12xy$? What is the sum of $-5xy$ and $+12xy$?

2. What is left when $+3mn$ is subtracted from $+10mn$? What is the sum of $-3mn$ and $+10mn$?

3. Instead of subtracting a positive number, what may be done to obtain the same result?

4. What is the result when $8a$ is subtracted from $10a$? When $(8a - 5a)$ is subtracted from $10a$? How does the second result compare with the first? What effect upon the result has the subtraction of the negative number $-5a$?

5. How does the result of subtracting $(5x - 2x)$ from $12x$ compare with the result of subtracting $5x$ from $12x$? What effect upon the result has the subtraction of the negative number $-2x$?

6. Instead of subtracting a negative number, what may be done to obtain the same result?

62. In addition two numbers are given, and their algebraic sum is required; in subtraction the algebraic sum, called the *minuend*, and one of the numbers, called the *subtrahend*, are given, and the other number, called the *remainder*, or *difference*, is required.

Subtraction is, therefore, the *inverse of addition*.

The **Difference** is the algebraic number that *added to the subtrahend gives the minuend*.

63. PRINCIPLES. — 1. *Subtracting a positive number is equivalent to adding a numerically equal negative number.*

2. *Subtracting a negative number is equivalent to adding a numerically equal positive number.*

The difference of similar terms, only, can be expressed in one term.

Principle 1 may be established as follows :

Let m represent any minuend and $+s$ any positive subtrahend.

It is to be proved that $m - +s = m + -s$.

§ 62, to find the remainder when $+s$ is subtracted from m is to find the algebraic number that added to $+s$ will give m .

Since the algebraic sum of $+s$ and $-s$ is 0, by the Associative Law for Addition the algebraic sum of $+s$ and $m + -s$ is $m + 0$, or m .

Hence, the algebraic number that added to $+s$ gives m is $m + -s$.

$$\therefore m - +s = m + -s.$$

Principle 2 may be established as follows :

Let m represent any minuend and $-s$ any negative subtrahend.

It is to be proved that $m - -s = m + +s$.

§ 62, to find the remainder when $-s$ is subtracted from m is to find the algebraic number that added to $-s$ will give m .

Since the algebraic sum of $-s$ and $+s$ is 0, by the Associative Law for Addition the algebraic sum of $-s$ and $m + +s$ is $m + 0$, or m .

Hence, the algebraic number that added to $-s$ gives m is $m + +s$.

$$\therefore m - -s = m + +s.$$

64. Since, from the above principles, subtracting algebraic numbers is equivalent to adding them to the minuend with their signs changed, it follows that the Laws of Order and Grouping for Addition hold in the subtraction of algebraic numbers; and that when one or more subtrahends with their signs changed are added to the minuend to form the algebraic sum called the difference, one set of signs, $+$ and $-$, suffices to denote either quality or operation.

65. To subtract when the terms are positive.

EXAMPLES

1. From $10x$ subtract $4x$.

PROCESS

$$\begin{array}{r} 10x \\ 4x \\ - \\ \hline 6x \end{array}$$

EXPLANATION. — Since subtracting a positive term is equivalent to adding a numerically equal negative term (Prin. 1), $4x$ may be subtracted from $10x$ by changing the sign of $4x$, and adding $10x$ and $-4x$.

2. From $10x$ subtract $15x$.

PROCESS

$$\begin{array}{r} 10x \\ 15x \\ \hline - \\ \hline - 5x \end{array}$$

EXPLANATION. — Since subtracting a positive term is equivalent to adding a numerically equal negative term (Prin. 1), $15x$ may be subtracted from $10x$ by changing the sign of $15x$ and adding $10x$ and $-15x$.

	3.	4.	5.	6.	7.	8.
From	$12a$	$9am$	$8x^2y^2$	$24mn^2$	$6\sqrt{ax}$	$11(a+b)$
Take	<u>$5a$</u>	<u>$21am$</u>	<u>$18x^2y^2$</u>	<u>$12mn^2$</u>	<u>$15\sqrt{ax}$</u>	<u>$21(a+b)$</u>

	9.	10.	11.	12.
From	$9a + 7b$	$5a + 10b$	$10x + 2y$	$3m + 3n$
Take	<u>$2a + 3b$</u>	<u>$7a + 4b$</u>	<u>$6x + 4y$</u>	<u>$2m + 5n$</u>

	13.	14.	15.	16.
From	$15m + n$	$7x + 2y$	$4x + 4y$	$8p + 3q$
Take	<u>$12m + 2n$</u>	<u>$4x + 4y$</u>	<u>$7x + 2y$</u>	<u>$10p + 2q$</u>

17. From $8p + 3z$ subtract $10p + z$.
18. From $15m + n$ subtract $5m + 3n$.
19. From $3ax + 5by$ subtract $4ax + 6by$.
20. From $8abc + 19mx$ subtract $20abc + 7mx$.
21. From $a + 3b + c$ subtract $a + b + 3c$.
22. From $12a^2 + 2b^2 + 14c^2$ subtract $3a^2 + 13b^2 + 3c^2$.
23. From $6ax + by + 7cz$ subtract $2ax + by + 2cz$.
24. From $7ax + by + 2cz$ subtract $4ax + 3by + cz$.
25. From $4ab + c$ subtract $a^2 + b^2 + abc + 2ab + 2c$.
26. From $5xy$ subtract $x^3 + 2x^2y + 2xy^2 + 3xy + y^3$.
27. From 1 subtract $x^4 + 13x^3 + 15x^2 + 16x + 25$.
28. From $7x^2 - 4y^2$ subtract $6x^2 + 3xy - 6y^2$.

66. To subtract when some terms are negative.

EXAMPLES

1. From $8x - 3y$ subtract $5x - 7y$.

PROCESS	EXPLANATION.
$8x - 3y$	Since subtracting a positive term is equivalent to adding a numerically equal negative term, subtracting $5x$ from $8x$ is equivalent to adding $-5x$ to $8x$ (Prin. 1).
$5x - 7y$	Since subtracting a negative term is equivalent to adding a numerically equal positive term, subtracting $-7y$ from $-3y$ is equivalent to adding $+7y$ to $-3y$ (Prin. 2).
$\begin{array}{r} - \quad + \\ \hline 3x + 4y \end{array}$	

RULE. — *Change the sign of each term of the subtrahend, or conceive it to be changed, and add the result to the minuend.*

	2.	3.	4.	5.	6.
From	$5a$	$6xy$	$-9mn$	$-13\sqrt{x}$	$-3(a+b)$
Take	<u>$-2a$</u>	<u>$-3xy$</u>	<u>$-4mn$</u>	<u>$-5\sqrt{x}$</u>	<u>$-10(a+b)$</u>

	7.	8.	9.
From	$4m - 3n + 2p$	$8a - 10b + c$	$3x + 2y - z$
Take	<u>$2m - 5n - p$</u>	<u>$6a - 5b - c$</u>	<u>$5x - 4y - z$</u>

	10.	11.	12.
From	$a - b + c$	$8a^2b - 5ac^2 + 9a^2c$	$r - s + t$
Take	<u>$2a + b - c$</u>	<u>$3a^2b + 2ac^2 - 9a^2c$</u>	<u>$r + s - t$</u>

13. From $5x - 3y + z$ take $2x - y + 8z$.14. From $3a^2b + b^3 - a^3$ take $4a^2b - 8a^3 + 2b^3$.15. From $13a^3 + 5b^2 - 4c^2$ take $8a^2 + 9b^2 + 10c^2$.16. From $15x - 3y + 2z$ subtract $3x + 8y - 9z$.17. From $a^2 - ab - b^2$ subtract $ab - 2a^2 - 2b^2$.18. From $m^2 - mn + n^2$ subtract $2m^2 - 3mn + 2n^2$.19. From $5x^2 - 2xy - y^2$ subtract $2x^2 + 2xy - 3y^2$.20. From $2ax - by - 5xy$ subtract $2by - 2ax - 3xy$.

21. From $2a + c$ subtract $a - b + c$.
22. From $2m + n$ subtract $n - 2p$.
23. From $x + y$ subtract $3a - 4 + y$.
24. From $2x^2 + 2xy$ subtract $x^2 - xy - y^2$.
25. From $2a - 2d$ subtract $a - b + c - d$.
26. From $2b$ subtract $b - a - c - d$.
27. From $a^3 + x^3$ subtract $a^3 - 3a^2x + 3ax^2 - x^3$.
28. From $a^4 + 1$ subtract $1 - a + a^2 - a^3 + a^4$.
29. From the sum of $3a^2 - 2ab - b^2$ and $3ab - 2a^2$ subtract $a^2 - ab - b^2$.
30. From $3x - y + z$ subtract the sum of $x - 4y + z$ and $2x + 3y - 2z$.
31. From $a + b + c$ subtract the sum of $a - b - c$, $b - c - a$, and $c - a - b$.
32. Subtract the sum of $m^2n - 2mn^2$ and $2m^2n - m^3 - n^3 + 2mn^2$ from $m^3 - n^3$.
33. Subtract the sum of $2c - 9a - 3b$ and $3b - 5a - 5c$ from $b - 3c + a$.
34. From $3bx + 4ay$ subtract the sum of $3ay - 4bx$ and $bx + ay$.
35. From the sum of $1 + x$ and $1 - x^2$ subtract $1 - x + x^2 - x^3$.
36. From $\frac{2}{3}x^3 - \frac{2}{3}x^2 + 3x - 7$ subtract $\frac{1}{2}x^3 - \frac{4}{3}x^2 + \frac{5}{2}x - 10$.
37. From $\frac{1}{8}m^3 - \frac{1}{4}m^2n + \frac{1}{8}mn^2 - \frac{1}{27}n^3$ subtract $n^3 - m^3 + \frac{1}{8}mn^2 - \frac{1}{2}m^2n$.
38. From $5(a + b) - 3(x + y) + 4(m + n)$ subtract $4(a + b) + 2(x + y) + (m + n)$.
39. From $n^5 - m^5$ subtract the sum of $2m^2n^3 - 3mn^4$ and $m^5 + 4m^3n^2 - 2m^2n^3 + 5mn^4 - n^5$.
40. From the sum of $3x^3 - 2x + 1$ and $2x - 5$ subtract the sum of $x - x^2 + 1$ and $2x^2 - 4x + 3$.

PARENTHESES

67. The subtrahend is sometimes written within a sign of aggregation preceded by the sign $-$.

If $a - b$ is to be subtracted from $2a$, it may be written $2a - (a - b)$.

1. What change must be made in the signs of the terms of the subtrahend, when it is subtracted from the minuend?

2. When a number *in parenthesis* is preceded by the sign $-$, what change must be made in the signs of the terms, when the subtraction is performed, or *when the parenthesis is removed*?

68. PRINCIPLES. — 1. *A parenthesis preceded by the minus sign may be removed from an expression, if the signs of all the terms in parenthesis are changed.*

2. *A parenthesis preceded by the minus sign may be used to inclose an expression, if the signs of all the terms to be inclosed are changed.*

1. When numbers are inclosed in a parenthesis preceded by the *plus* sign, the parenthesis may be removed without changing the signs of the terms.

2. Any number of terms may be inclosed in a parenthesis preceded by a *plus* sign without changing the signs of the terms.

3. The student should remember that in an expression like $-(x - y)$, or $-x - y$, the sign of x is plus, and the expression is the same as if it were written $-(+x - y)$, or $-+x - y$.

EXAMPLES

Simplify the following:

1. $a + (b - c)$.

8. $a - b - (c - d)$.

2. $a - (b - c)$.

9. $a - b - (-c + a)$.

3. $x - (y - z)$.

10. $a - m - (n - m)$.

4. $x - (-y + z)$.

11. $5a - 2b - (a - 2b)$.

5. $m - n - (-a)$.

12. $a - (b - c + a) - (c - b)$.

6. $m - (n - 2a)$.

13. $2xy + 3y^2 - (x^2 + xy - y^2)$.

7. $5x - (2x + y)$.

14. $m + (3m - n) - (2n - m) + n$.

Collect in alphabetical order the coefficients of x and y in the following, giving each parenthesis the sign of the first coefficient to be inclosed therein :

$$15. ax - by - 3bx + 2cy - fx - gy.$$

PROCESS

$$ax - by - 3bx + 2cy - fx - gy = (a - 3b - f)x - (b - 2c + g)y$$

SUGGESTION. — The coefficient of y is $-b + 2c - g$, which is written $-(b - 2c + g)$ (Prin. 2).

- | | |
|---------------------------------------|----------------------------|
| 16. $ax - by - bx - cy + dx - ey.$ | 21. $bx - cy - 2ay + by.$ |
| 17. $mx - 2ny + nx - ry - px + qy.$ | 22. $mx - bx - 4y - my.$ |
| 18. $5ax + 3ay - 2dx + ny - 5x - y.$ | 23. $rx - ay - sx + 2cy.$ |
| 19. $cx - 2bx + 7ay + 3ax - lx - ty.$ | 24. $x^2 + ax - y^2 + ay.$ |
| 20. $bx + cy - 2ax + by - cx - dy.$ | 25. $x^2 - ay - ax - y^2.$ |

Group the same powers of x in the following :

$$26. ax^3 + bx^2 - cx + ex^2 - dx^2 - fx.$$

$$27. x^3 + 3x^2 + 3x - ax^2 - 3ax^2 + bx.$$

$$28. x^2 - abx - x^3 - bx^2 - cx - mnx^2 + dx.$$

$$29. ax^4 - x^4 - ax^2 + x^2 + ax - x - abx^2 + x^2.$$

$$30. \text{Simplify } 2a - [a - \{b - (3b - 2a - b)\}] - (b - a).$$

When an expression contains parentheses within parentheses, they may be removed *in succession*, beginning with either the outermost or the innermost, preferably the innermost.

SOLUTION

	$2a - [a - \{b - (3b - 2a - b)\}] - (b - a)$
Prin. 1,	$= 2a - [a - \{b - (3b - 2a + b)\}] - b + a$
Uniting terms,	$= 3a - [a - \{b - (4b - 2a)\}] - b$
Prin. 1,	$= 3a - [a - \{b - 4b + 2a\}] - b$
Uniting terms,	$= 3a - [a - \{-3b + 2a\}] - b$
Prin. 1,	$= 3a - [a + 3b - 2a] - b$
Uniting terms,	$= 3a - [-a + 3b] - b$
Prin. 1,	$= 3a + a - 3b - b$
Uniting terms,	$= 4a - 4b.$

Simplify the following:

31. $4a + b - \{x + 4a + b - 2y - (x + y)\}.$
32. $ab - \{ab + ac - a - (2a - ac) + (2a - 2ac)\}.$
33. $a + [y - \{5 + 4a - (6y + 3)\} - (7y - 4a - 1)].$
34. $4m - [p + 3n - (m + n) + 3 - (6p - 3n - 5m)].$
35. $a + 2b + (14a - 5b) - \{6a + 6b - (5a - \overline{4a - 4b})\}.$
36. $12a - \{[4 - 3b - (6b + 3c)] + b - 8 - (5a - 2b - 6)\}.$
37. $a + b - \{-[a + b - (c + x)] - [3a - (c - x + a) - b] + 4a\}.$
38. $x^3 - [x^2 - (1 - x)] - \{1 + [x^2 - (1 - x) + x^2]\}.$
39. $4 - \{[5y - (3 - 2x - 2)] - [x + (5y - x + 3)]\}.$
40. $ab - \{5 + x - (b + c - ab + x)\} + [x - (b - c - 7)].$
41. $a^3 - b^2 - \{ad + a^2 - (x + a^2 - b^2) - b^2\} + 5ad - (x + 3ad).$
42. $a - (b - c) - [a - \{b - c - (b + c - a) + (a - b) + (c - a)\}].$
43. $-\{3ax - [5xy - 3z] + z - (4xy + [6z + 7ax] + 3z)\}.$
44. $1 - x - \{1 - x - [1 - x - (1 - x) - (x - 1)] - x + 1\}.$
45. $1 - x - \{1 - [x - 1 + (x - 1) - (1 - x) - x] + 1 - x\}.$
46. $x - [-\{-(-x) + x\} - 2x].$
47. $(a - b) - \{-a - (b - a) + (a - b)\}.$
48. $a - 7 - [-\{-a - (-a - \overline{a - 3})\}].$
49. $a - x - [-\{a + (x - a) - (x - 4a)\}].$
50. $5xy - [-\{(y^2 - xy) - (xy - \overline{y^2 - 2xy})\}].$
51. $2a - [a - \{b - (3b - \overline{2a - b})\}] - (b - a).$
52. $a - [-\{(m - a) - \{a - (m - \overline{2m + 6a})\}].$
53. $a - \{-b - (c - d)\} + a - [-b + \{-2c - (d - e)\}].$
54. $a^2 + 5 - [2ab - \{- (7 - 3ab) - \overline{ab + 2a^2 - z}\} - (3a - z)].$
55. $2x + (3y - \{2x - [y + 4x - (3y - x)] - 2y\} - \overline{x - y}).$
56. $1 - (-\{-[-(-a - \overline{a - 1}) - 3] - 2\} - a) - [a - (a - 1)].$

TRANSPOSITION IN EQUATIONS

69. 1. What number diminished by 2 is equal to 8?

2. If a number increased by 2 is equal to 8, what is the number?

3. In the equation $x - 2 = 8$, what is done with the 2 in obtaining the value of x ? In the equation $x = 8 + 2$, how does the sign of 2 compare with its sign in the previous equation?

4. In the equation $x + 2 = 8$, what is done with the 2 in obtaining the value of x ? In the equation $x = 8 - 2$, how does the sign of 2 compare with its sign in the previous equation?

5. In changing the 2's from one side of the equation to the other, what change was made in the sign?

6. When a term is changed from one side, or *member* of an equation, to the other, what change must be made in its sign?

7. If 3 is added to one member of the equation $2 + 5 = 7$, what must be done to preserve the equality?

8. If 3 is subtracted from one member of the equation $2 + 8 = 10$, what must be done to preserve the equality?

9. If one member of the equation $2 + 5 = 7$ is multiplied by 5, what must be done to preserve the equality?

10. If one member of the equation $10 + 25 = 35$ is divided by 5, what must be done to preserve the equality?

11. If one member of the equation $x = 5$ is raised to the second power, what must be done to preserve the equality?

12. If the square root of one member of the equation $x^2 = 25$ is taken, what must be done to preserve the equality?

13. What, then, may be done to the members of an equation without destroying the equality?

70. The parts of an equation on each side of the sign of equality are called its *Members*.

The part on the left of the sign of equality is called the *First Member*, and the part on the right, the *Second Member*.

71. The process of changing a term from one member of an equation to the other is called **Transposition**.

72. **PRINCIPLE.** — *A term may be transposed from one member of an equation to the other, provided its sign is changed.*

73. A truth that does not need demonstration is called an **Axiom**.

74. **AXIOMS.** — 1. *Things that are equal to the same thing are equal to each other.*

2. *If equals are added to equals, the sums are equal.*
3. *If equals are subtracted from equals, the remainders are equal.*
4. *If equals are multiplied by equals, the products are equal.*
5. *If equals are divided by equals, the quotients are equal.*
6. *The same powers of equal numbers are equal.*
7. *The same roots of equal numbers are equal.*

EQUATIONS AND PROBLEMS

75. 1. If $5x - 2 = 3x + 6$, find the value of x .

PROCESS

$$\begin{array}{rcl}
 5x - 2 & = & 3x + 6 \\
 3x & = & 3x \\
 \hline
 2x - 2 & = & +6 \\
 2 & = & 2 \\
 \hline
 2x & = & 8 \\
 \therefore x & = & 4
 \end{array}$$

OR

$$\begin{array}{rcl}
 5x - 2 & = & 3x + 6 \\
 5x - 3x & = & 2 + 6 \\
 2x & = & 8 \\
 \therefore x & = & 4
 \end{array}$$

VERIFICATION

$$\begin{array}{rcl}
 20 - 2 & = & 12 + 6 \\
 18 & = & 18
 \end{array}$$

EXPLANATION. — Since the value of x is sought, all terms containing x must be collected in one member of the equation, and the remaining, or known terms in the other member.

$3x$ may be made to disappear from the second member by subtracting $3x$ from both members. The result (Axiom 3) is the equation $2x - 2 = 6$.

-2 may be made to disappear from the first member by adding 2 to both members. The result (Axiom 2) is the equation $2x = 8$.

Dividing both members of this equation by 2, the coefficient of x , the resulting equation (Axiom 5) is $x = 4$, the value of x sought.

Or, since a term may be transposed from one member to the other if its sign is changed (Prin.), $3x$ transposed to the first member becomes $-3x$ and -2 transposed to the second member becomes $+2$. Therefore, the resulting equation is

$$5x - 3x = 2 + 6.$$

Uniting terms, $2x = 8$. Dividing both members by 2, the result is $x = 4$.

The result is *verified* by substituting the value of x for x in the original equation. If the members are then *identical*, the value found for the unknown number is correct.

2. If $5x - 7 = 30 - 7$, find the value of x .

PROCESS

$$5x - 7 = 30 - 7.$$

$$5x = 30.$$

$$\therefore x = 6.$$

SUGGESTION. — Since, if -7 were transposed from the first member to the second, it would appear as $+7$ and cancel the term -7 in that member, the two equal terms may be canceled before the transposition.

RULE. — *Transpose terms so that the unknown terms stand in the first member of the equation and the known terms in the second.*

Unite similar terms, and divide both members of the equation by the coefficient of the unknown number.

VERIFICATION. — *Substitute in the original equation the value of the unknown number thus found. If the members of the equation are then identical, the value of the unknown number found is correct.*

1. The same term with the same sign in both members of an equation may be canceled (Ax. 2 or 3).

2. If the signs of all the terms of an equation are changed, the equality will not be destroyed; for (Ax. 3) both members may be subtracted from 0 without destroying the equality.

Find the value of x and verify:

3. $5x + 3 = 8.$

10. $7x - 10 = 60.$

4. $x + 5 = 11.$

11. $7 + 2x = 11.$

5. $x - 5 = 11.$

12. $3 + 2x = 15.$

6. $2x - 3 = 21.$

13. $1 + 12x = 85.$

7. $5x + 7 = 42.$

14. $5 + 3x = 11.$

8. $3x - 2 = 25.$

15. $7 + 5x = 47.$

9. $2x + 4 = 10.$

16. $2 + 9x = 74.$

- | | |
|--------------------------|-------------------------------|
| 17. $3 = 5 - x$. | 26. $7x - 12 = x + 13 + 5$. |
| 18. $9 - 5x = -1$. | 27. $4x - 20 = 5x - 50 + x$. |
| 19. $7x + 2 = x + 14$. | 28. $3x + 16 = 20 - 5x + 4$. |
| 20. $5x - 5 = 2x + 4$. | 29. $7x - 55 = 18 - 2x - 1$. |
| 21. $3x + 2 = x + 30$. | 30. $-x - 12 = 40 - 8x + 4$. |
| 22. $5x - 2 = 2x + 7$. | 31. $80 - 3x = 83 - 8x + 7$. |
| 23. $2 + 13x = 50 - 9$. | 32. $9x - 90 = 16 - x + 4$. |
| 24. $10 + x = 18 - x$. | 33. $50 - x = 20 + x$. |
| 25. $2x + 2 = 32 - x$. | 34. $7x + 25 = 30 + 6x - 3$. |

Solve the following problems:

35. What number increased by 10 is equal to 19?
36. What number diminished by 30 is equal to 20?
37. What number diminished by 111 is equal to -15 ?
38. What number exceeds $\frac{1}{3}$ of itself by 10?

SUGGESTION. — Let $3x =$ the number.

39. Five times a number exceeds 3 times the number by 14.
What is the number?

40. If 5 is subtracted from a certain number, and the difference is subtracted from 3 times the number, the result is 35.
What is the number?

41. The double of a number is 64 less than 10 times the number.
What is the number?

42. If 4 is subtracted from a certain number, and the difference is subtracted from 40, the result is 3 times the number.
What is the number?

43. Three times a certain number is as much less than 72 as 4 times the number exceeds 12. What is the number?

44. Twice a certain number exceeds $\frac{1}{3}$ of the number as much as 6 times the number exceeds 65. What is the number?

45. If 16 is added to a certain number, the result is 56 diminished by 7 times the number. What is the number?

46. If 6 times a certain number lacks as much of 62 as 3 times the number exceeds 19, what is the number?

47. Three times a certain number increased by a is equal to the number increased by $9a$. What is the number?

48. The sum of 4 numbers in a row is 58. If their common difference is 3, what are the numbers?

SUGGESTION. — Let x = the smallest number.

Then, $x + 3$ = the second number,

$x + 6$ = the third number,

and $x + 9$ = the fourth number.

49. A man distributed 1 dollar among 5 boys so that each boy except the youngest received 5 cents more than the boy next younger. If the boys were all of different ages, how much did each receive?

50. The common difference of 5 numbers is 2, and their sum is 100. What are the numbers?

51. John and James were comparing their earnings. John said, "I have earned 50 cents." James replied, "If I had earned half as much as I have, and 10 cents more, I should have earned the same as you." How much had James earned?

52. A drover, when asked how many cattle he had, replied, "If I had $\frac{1}{2}$ more than I have and 2 more, I should have 200." How many cattle had he?

53. The earnings of a mill for 4 years were \$46,000. If the books showed an annual increase of \$1000, what were the earnings for each year?

54. James had $\frac{2}{3}$ as much money as John, John 5 cents less than William, and Robert 5 cents more than 3 times as much as James. If they together had \$1.50, how much had each?

SUGGESTION. — Let $3x$ = the number of cents John had.

55. A speculator who doubled his money by a fortunate investment, afterward lost \$600, but he still had \$400 more than the original sum. How much had he at first?

MULTIPLICATION

76. 1. How many are $3x + 3x + 3x + 3x + 3x$?
2. How many are 5 times $3x$? 2 times $3x$? 7 times $2a$?
3. A man saves \$10 a month, indicated by $+10$. How many dollars will he save in a year? What sign should be placed before the result to indicate the number of dollars *saved*?
4. How many are 12 times $+10$? 12 times $+10a$? 5 times $+3x$? 8 times $+2m$?
5. When a positive number is multiplied by a positive number, what is the sign of the product?
6. If a man loses \$5 a month, indicated by -5 , how many dollars will he lose in a year? What sign should be placed before the result to indicate the number of dollars *lost*?
7. How many are 12 times -5 ? 12 times $-5b$? 7 a times $-3x$? 3 times $-2b$? 11 times $-3y$? 5 times $-5ab$?
8. When a negative number is multiplied by a positive number, what is the sign of the product?
9. If a man's gains in business are \$10 a month, indicated by $+10$, how many dollars less had he 3 months ago, indicated by -3 , than he has now? Indicate the result algebraically.
10. How many are $+10$ multiplied by -3 ? $+10$ multiplied by -5 ? $+2$ multiplied by -3 ? $+a$ multiplied by $-b$?
11. What is the sign of the product when a positive number is multiplied by a negative number? When a negative number is multiplied by a positive number?
12. What, then, is the sign of the product of two numbers having unlike signs?

13. If a man who is in debt is getting deeper in debt at the rate of \$10 a month, indicated by -10 , how much better off was he 3 months ago, indicated by -3 , than he is now? Indicate the result algebraically.

14. How many are -10 multiplied by -3 ? -10 multiplied by -5 ? -2 multiplied by -3 ? $-a$ multiplied by $-b$?

15. What is the sign of the product of two negative numbers? of two positive numbers?

16. What, then, is the sign of the product of two numbers having like signs?

17. What is the sign of -5×-2 ? of $-5 \times -2 \times -2$? of $-5 \times -2 \times +2 \times -2$? of $+5 \times -2 \times -2 \times -2 \times -2$? of $-2 \times -3 \times +2 \times +2 \times -2 \times -2 \times -2$?

18. What sign has a product, if the number of negative factors is *even*? What sign has a product, if the number of negative factors is *odd*?

19. What, then, determines the sign of a product?

20. In the expression a^3 , what is 3 called? What does it indicate? In a^7 how many times is a used as a factor?

21. When a^3 is multiplied by a^4 , how many times is a used as a factor in the product? when a^2 is multiplied by a^5 ?

When a^m is multiplied by a^n , what is the product, if m and n are positive integers?

22. How, then, is the exponent of a factor in the product determined?

77. When the multiplier is a positive integer, the process of taking the multiplicand additively as many times as there are units in the multiplier is called **Multiplication**.

When the multiplier is any number, multiplication may be defined as *the process of finding a number that has the same relation to the multiplicand as the multiplier has to 1*.

The multiplicand and multiplier are called the *factors* of the product.

78. PRINCIPLES. — 1. **Law of Signs.** — *The sign of the product of two factors is + when they have like signs, and — when they have unlike signs.*

2. Law of Coefficients. — *The coefficient of the product is equal to the product of the coefficients of the factors.*

3. Law of Exponents. — *The exponent of a number in the product is equal to the sum of its exponents in the factors.*

79. The Law of Signs may be established as follows :

$$+3 = +1 + +1 + +1, \quad (1)$$

$$\text{and} \quad -3 = -1 + -1 + -1 = -+1 -+1 -+1; \quad (2)$$

that is, +3 is obtained from +1 by taking +1 additively three times, and -3 by taking +1 subtractively three times.

Hence, § 77, multiplying any number by +3 is equivalent to taking that number additively three times, and multiplying any number by -3 is equivalent to taking that number subtractively three times.

$$\text{By (1),} \quad +5 \times +3 = +5 + +5 + +5 = +15, \quad (3)$$

$$\text{and} \quad -5 \times +3 = -5 + -5 + -5 = -15. \quad (4)$$

$$\text{By (2),} \quad +5 \times -3 = -+5 -+5 -+5 = -15, \quad (5)$$

$$\text{and} \quad -5 \times -3 = - -5 - -5 - -5 = +15. \quad (6)$$

$$\text{Similarly, since } +\left(\frac{1}{2}\right) = +\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right) + \left(\frac{1}{2}\right) \text{ and } -\left(\frac{1}{2}\right) = -+\left(\frac{1}{2}\right) -+\left(\frac{1}{2}\right) -+\left(\frac{1}{2}\right),$$

$$+5 \times +\left(\frac{1}{2}\right) = +\left(\frac{5}{2}\right) + \left(\frac{5}{2}\right) + \left(\frac{5}{2}\right) = +\left(\frac{15}{2}\right); \quad (7)$$

and so on, as in (4), (5), and (6).

In (3) and (6) the product of two algebraic numbers with like signs is positive.

In (4) and (5) the product of two algebraic numbers with unlike signs is negative.

(7) shows that like results are obtained when the multiplier is a fractional number (§ 7).

Passing to general symbols, let a and b be any absolute numbers.

First, when b is a whole number.

$$\text{Since} \quad +b = +1 + +1 + +1 + \dots \text{ to } b \text{ terms,}$$

$$\begin{aligned} \S 77, \quad +a \times +b &= +a + +a + +a + \dots \text{ to } b \text{ terms} \\ &= +ab, \end{aligned} \quad (8)$$

$$\begin{aligned} \text{and} \quad -a \times +b &= -a + -a + -a + \dots \text{ to } b \text{ terms} \\ &= -ab. \end{aligned} \quad (9)$$

$$\text{Since} \quad -b = -+1 -+1 -+1 - \dots \text{ to } b \text{ terms,}$$

$$\begin{aligned} \S 77, \quad +a \times -b &= -+a -+a -+a - \dots \text{ to } b \text{ terms} \\ &= -ab, \end{aligned} \quad (10)$$

$$\begin{aligned} \text{and} \quad -a \times -b &= - -a - -a - -a - \dots \text{ to } b \text{ terms} \\ &= +ab. \end{aligned} \quad (11)$$

Second, when b is a fractional number.

As in (7), the same reasoning applies when b is a fractional number.

Hence, from (8) and (11), the product of any two algebraic numbers with like signs is positive; and from (9) and (10), the product of any two algebraic numbers with unlike signs is negative.

When the multiplier is a positive or negative whole or fractional number, it appears from the above proofs that algebraic multiplication is only abbreviated algebraic addition. Hence, as in addition, but one set of signs $+$ and $-$ is required to denote both quality and operation.

Hence, the Law of Signs may be expressed as follows:

$+ a$ multiplied by $+ b = + ab$,

$- a$ multiplied by $- b = + ab$,

$- a$ multiplied by $+ b = - ab$,

and

$+ a$ multiplied by $- b = - ab$.

80. It follows from the Law of Signs, applied repeatedly, that the product of any number of algebraic numbers is $+$ when the number of negative factors is even, and $-$ when the number of negative factors is odd.

81. The Law of Exponents or the Index Law for multiplication may be established for *positive integral exponents* as follows:

Let m and n be any positive integers.

By the definition of a power, § 24,

$$a^m = a \times a \times a \dots \text{to } m \text{ factors,}$$

$$a^n = a \times a \times a \dots \text{to } n \text{ factors;}$$

$$\therefore a^m \times a^n = (a \times a \times a \dots \text{to } m \text{ factors})(a \times a \times a \dots \text{to } n \text{ factors}) \\ = a \times a \times a \dots \text{to } (m + n) \text{ factors.}$$

Hence, $a^m \times a^n = a^{m+n}$.

In like manner, $a^m \times a^n \times a^p = a^{m+n+p}$.

Thus, $a^2 \times a^4 = a^{2+4} = a^6$,

and $a^8 \times a^2 \times a^4 = a^{8+2+4} = a^{14}$.

82. 1. How does 2×5 compare with 5×2 in value? 3×7 with 7×3 ? $2 \times 5 \times 6$ with $2 \times 6 \times 5$?

2. What is the effect upon the value of a product of changing the order of its factors?

Law of Order, or Commutative Law for Multiplication. — *The factors of a product may be taken in any order.*

The Law of Order may be established as follows :

Since the number of negative factors will not be changed by taking the factors in any order, § 80, the *sign* of the product is the same in whatever order the factors are taken.

We know from arithmetic that arithmetical numbers may be multiplied in any order. Hence, the *absolute value* of the product is the same in whatever order the factors are taken.

Since neither the sign nor the absolute value of the product of algebraic numbers is changed by changing the order of the factors, the factors may be taken in any order.

In general symbols, $a \times b \times c \times \dots = b \times c \times a \times \dots = \text{etc.}$

83. 1. How does $2 \times 3 \times 5$, or 6×5 , compare in value with $2 \times (3 \times 5)$, or 2×15 ? with 5×6 ? $a \times b \times c$, or $(ab) \times c$, with $a \times (bc)$?

2. How may the factors of a product be grouped?

Law of Grouping, or Associative Law for Multiplication. — *The factors of a product may be grouped in any manner.*

The Law of Grouping may be established as follows :

By the notation of multiplication, abc denotes that a is to be multiplied by b and then the product ab is to be multiplied by c ; that is,

$$abc = (ab)c. \quad (1)$$

1. Let it be required to prove that $(ab)c = a(bc)$.

By the Law of Order,

$$abc = bca$$

by notation,

$$= (bc)a$$

by the Law of Order,

$$= a(bc).$$

(2)

From (1) and (2),

$$(ab)c = a(bc).$$

Similarly, it may be proved that $(ab)c = b(ac)$, etc.

2. Let it be required to prove that $abcd = (ab)(cd)$.

By notation,

$$abcd = ab \times c \times d$$

putting m for ab ,

$$= m \cdot c \cdot d, \text{ or } mcd.$$

By 1,

$$m \cdot c \cdot d = m(cd).$$

Putting ab for m ,

$$(ab) \cdot c \cdot d = (ab)(cd);$$

that is, by notation,

$$abcd = (ab)(cd).$$

Similarly, it can be shown that $(abc)d = a(bcd) = (bc)(ad) = (ac)(bd) = (abd)c = (adc)b = c(dba)$ = etc., the factors being grouped in any manner whatever.

3. In a similar way the law may be established for any number of factors, successively for 5, 6, 7, ... factors.

Hence, $abc \dots p = a(bc \dots p) = b(ac \dots p)$, etc., for all values of the letters.

84. To multiply a monomial by a monomial.

EXAMPLES

1. Multiply $5x^2y^3$ by $-3xy^2z$.

PROCESS

$$\begin{array}{r} 5x^2y^3 \\ - 3xy^2z \\ \hline -15x^3y^5z \end{array}$$

EXPLANATION. — Since the multiplier is composed of the factors -3 , x , y^2 , and z , the multiplicand may be multiplied by each successively. -3 times $5x^2y^3 = -15x^2y^3$ (Prin. 1 and 2); x times $-15x^2y^3 = -15x^3y^3$ (Prin. 3); y^2 times $-15x^3y^3 = -15x^3y^5$ (Prin. 3); and this multiplied by z is equal to $-15x^3y^5z$ (Prin. 3).

Or, since the signs of the numbers are unlike, the sign of their product is $-$; the coefficient of the product is the product of the coefficients 5 and 3; and the product of the literal numbers is expressed by affecting each with an exponent equal to the sum of its exponents in the factors.

RULE. — *To the product of the numerical coefficients annex the letters, each with an exponent equal to the sum of its exponents in both factors.*

Write the sign $+$ before the product when its factors have like signs, and $-$ when they have unlike signs.

	2.	3.	4.	5.	6.
Multiply	-2	6	-7	$2a$	$2m^3$
By	8	-2	-9	5	$6m^2$
	7.	8.	9.	10.	11.
Multiply	$10a^5$	x^2y^2	$-4abc$	$5a^2bc^3$	$-2xy^2$
By	$5a^3$	xy^3	$2a^2b$	$-7ab^2c$	$2x^2y$
	12.	13.	14.	15.	16.
Multiply	$-3a^2x^2$	$-5m^3n^2$	$-6a^2b^2c^2x$	$4abcd$	$-3x^2by^3$
By	$-2ax^3$	$3mn$	$-4a^3bny^2$	-1	-1
	17.	18.	19.	20.	21.
Multiply	$-2a^6x^2$	$-3n^3y$	$4a^2xb^3y^4$	-1	$-5m^3n^2a^2y^3$
By	$-4ax^4$	$6b^3y$	$3a^3x^4b^2y$	-1	$-2m^{10}n^6c^2y^2$
	22.	23.	24.	25.	26.
Multiply	$5pq^2x^3$	$10m^4n^3$	$-2a^2m^3n^4$	x^2yz	$-p^2d^2q^2$
By	$-2rq^4x$	$-3n^2m^4$	$8b^3n^6m^7$	$-x^2yz^2$	$-abc$

	27.	28.	29.	30.	31.
Multiply	$2a^{m+1}$	$-5x^n$	$-x^ny^n$	$-x^{n-1}y^{n-2}$	$4x^{n-1}$
By	$3a^2$	x	$3xy$	$-xy$	$-2x^{n+4}$
	32.	33.	34.	35.	36.
Multiply	$5y$	$-a^m$	a^{m-n}	$-x^{1-n}$	y^{n-m}
By	$-3y^{n-2}$	$-a^n$	a^n	$-x^n$	y^{m-n+1}
	37.		38.		39.
Multiply	$a^2b^3x^3y^{n-2}$		$a^{n-1}b^{n-2}c^3$		$m^an^2b^2y^a$
By	$a^nb^{n-3}y^2$		$a^{n+1}b^2c^{n-1}$		$m^bn^2b^2y^{b-a}$

85. How does 25×2 compare in value with 20×2 plus 5×2 ? How is 133, or $100 + 30 + 3$, multiplied by 2? How is the polynomial $a + b + c$ multiplied by the monomial m ?

Distributive Law for Multiplication.— *The product of a polynomial by a monomial is equal to the algebraic sum of the partial products formed by multiplying each term of the polynomial by the monomial.*

The Distributive Law may be established as follows:

Let $a + b$ be the multiplicand and m the multiplier, a , b , and m being positive or negative integral or fractional numbers.

By the Law of Order the multiplier may change places with the multiplicand. Hence, $(a + b) \times m$ may be written $m(a + b)$.

It is to be proved that $m(a + b) = ma + mb$.

First, when m is a positive integer.

Since $m = 1 + 1 + 1 + \dots$ to m terms,

§ 77, $m(a + b) = (a + b) + (a + b) + (a + b) + \dots$ to m terms

§ 56, $= (a + a + a + \dots$ to m terms) $+ (b + b + b + \dots$ to m terms)

§ 77, $= ma + mb.$ (1)

Second, when m is a fractional number.

Let $m = \frac{p}{q}$, in which p and q are absolute integers.

$$\frac{p}{q}(a + b) = p \text{ times one } q\text{th of } (a + b)$$

$$= p(a + b) \text{ qths}$$

$$\text{by (1),} \quad = pa \text{ qths} + pb \text{ qths}$$

$$= \frac{p}{q} \text{ of } a + \frac{p}{q} \text{ of } b$$

$$= \frac{p}{q}a + \frac{p}{q}b. \quad (2)$$

Third, when m is negative.

Let $m = -n$, n being any positive whole or fractional number.

It is to be proved that $(-n)(a+b) = -na - nb$.

By (1) and (2), $n(a+b) = na + nb$. (3)

Since, if $+n$ is positive, $-(-n)$ is also positive, substituting $-(-n)$ for $+n$ in (3),

$$\begin{aligned} -(-n)(a+b) &= -(-n)a - (-n)b \\ &= -(-na - nb). \end{aligned} \quad (4)$$

Since both $-(-n)(a+b)$ and $-(-na - nb)$ are now *monomial* in form, both members of (4) may be multiplied by the monomial -1 .

$$\therefore +(-n)(a+b) = +(-na - nb),$$

or $(-n)(a+b) = -na - nb$. (5)

By (1), (2), and (5), $m(a+b) = ma + mb$

for all positive or negative whole or fractional values of m and for all values of a and b .

86. To multiply a polynomial by a monomial.

EXAMPLES

1. Multiply $3x^2 - y^2$ by $-4y$.

PROCESS	EXPLANATION.
$3x^2 - y^2$	By § 85, each term of the multiplicand is to be multiplied by the multiplier.
$-4y$	The product of $3x^2$ and $-4y$ is $-12x^2y$. But since the entire multiplicand is $3x^2 - y^2$, $-4y$ times y^2 must be subtracted from $-12x^2y$. $-4y$ times $y^2 = -4y^3$, which subtracted from $-12x^2y$ gives $-12x^2y + 4y^3$.
$-12x^2y + 4y^3$	

Or, since a polynomial multiplied by a monomial is equal to the algebraic sum of the partial products formed by multiplying each term of the polynomial by the monomial, § 85, $3x^2 - y^2$ multiplied by $-4y$ is equal to $-12x^2y + 4y^3$.

RULE. — *Multiply each term of the polynomial by the monomial, and find the algebraic sum of the partial products.*

	2.	3.	4.
Multiply	$2a^2 - 2ab + 3b^2$	$5m^2 - 4n^2$	$5x^2 - 2xy - y^2$
By	$3ab$	$-2m^2n$	$-x^2y$

Multiply:

5. $3x^2 - 2xy$ by $5xy^2$.

8. $p^2q^2 - 2pq^3$ by $-pq$.

6. $m^2n^3 - 3mn^4$ by $2mn$.

9. $4a^2 - 5b^2c$ by abc^2 .

7. $3a^3 - 6a^2b$ by $-2b$.

10. $-2ac + 4ax$ by $-5acx$.

Perform the multiplications indicated :

11. $a^2bc(3a^4 - 4a^3b - 5a^2b^2 + 2ab^3 - 16b^4)$.
12. $2xy(5x^2 - 10xy - 36y^2 - 5x + 5y + 120)$.
13. $5m^3(16m^3 - 20m^2n + 13mn^2 - 25n^3)$.
14. $abc(a^2b^2 - 2a^2c^2 - 2b^2c^2 - a^4 - 4b^4 - c^4 - 5abc)$.
15. $-bc(b^4 + c^4 - b^3 - c^3 + b^2c^2 - 4b^2c + 8bc^2 - 2bc)$.
16. $-2x(x^4 - 5x^3y - 16x^2y^2 + 24xy^3 - y^4 - xy - x + 4)$.

87. To multiply a polynomial by a polynomial.

To multiply $p + q + r$ by $a + b$,

$$\S 85, \quad (p + q + r)(a + b) = p(a + b) + q(a + b) + r(a + b)$$

$$\S\S 82, 85, \quad = ap + bp + aq + bq + ar + br$$

$$\S 55, \quad = ap + aq + ar + bp + bq + br.$$

RULE. — *Multiply every term of the multiplicand by each term of the multiplier, and find the algebraic sum of the partial products.*

EXAMPLES

- i. Multiply $x^2 - xy$ by $2x + 3y$.

PROCESS

$$\begin{array}{r}
 x^2 - xy \\
 2x + 3y \\
 \hline
 + 2x \text{ times } (x^2 - xy) = 2x^3 - 2x^2y \\
 + 3y \text{ times } (x^2 - xy) = \quad + 3x^2y - 3xy^2 \\
 \hline
 \therefore (2x + 3y) \text{ times } (x^2 - xy) = 2x^3 + x^2y - 3xy^2
 \end{array}$$

2. Multiply $x^3 - 3x^2y + 3xy^2 - y^3$ by $x^2 - xy$.

PROCESS

$$\begin{array}{r}
 x^3 - 3x^2y + 3xy^2 - y^3 \\
 x^2 - xy \\
 \hline
 x^5 - 3x^4y + 3x^3y^2 - x^2y^3 \\
 - x^4y + 3x^3y^2 - 3x^2y^3 + xy^4 \\
 \hline
 x^5 - 4x^4y + 6x^3y^2 - 4x^2y^3 + xy^4
 \end{array}$$

3. Multiply $5x + x^3 - 2x^2 - x^4$ by $3x + 1 - x^2$.

PROCESS	TEST
$5x - 2x^2 + x^3 - x^4$	$= + 3$
$1 + 3x - x^2$	<u>$= + 3$</u>
$5x - 2x^2 + 1x^3 - 1x^4$	
$15 \quad - 6 \quad + 3 \quad - 3x^5$	
$\quad \quad - 5 \quad + 2 \quad - 1 \quad + x^6$	
<hr/> $5x + 13x^2 - 10x^3 + 4x^4 - 4x^5 + x^6$	$= + 9$

SUGGESTION. — For convenience in writing partial products, both polynomials are arranged so that in passing from left to right the several powers of x are either successively higher or lower. In this process, the polynomials are arranged according to the ascending powers of x .

TEST. — Since the correct product of $5x - 2x^2 + x^3 - x^4$ and $1 + 3x - x^2$ is the same in form whatever value x represents, it is possible, by assigning an arithmetical value to x , to change the process of multiplying one algebraic expression by the other into a process of multiplying one arithmetical number by another as shown in the test.

Let 1 be substituted for x .

Multiplicand = $5x - 2x^2 + x^3 - x^4 = 5 - 2 + 1 - 1 = + 3$	
Multiplier = $1 + 3x - x^2 = 1 + 3 - 1 = + 3$	
Product should be equal to	<u>$+ 9$</u>

$$5x + 13x^2 - 10x^3 + 4x^4 - 4x^5 + x^6 = 5 + 13 - 10 + 4 - 4 + 1 = + 9.$$

In like manner the multiplication of any two literal expressions may be tested arithmetically by assigning any values we please to the letters. It is usually most convenient to substitute $+ 1$ for each letter, since this may be readily done by adding the numerical coefficients.

Multiply, and test each result:

- | | |
|------------------------------|--|
| 4. $2x + 3$ by $x + 2$. | 12. $4y - 6b$ by $2y + b$. |
| 5. $4x + 1$ by $3x + 4$. | 13. $3x - 2y$ by $3x + 2y$. |
| 6. $5n - 1$ by $4n + 5$. | 14. $2b + 5c$ by $5b - 2c$. |
| 7. $h + 2k$ by $3h - k$. | 15. $7x - 2n$ by $4x + 2n$. |
| 8. $3r - 6s$ by $5r - 2s$. | 16. $ab - 15$ by $ab + 10$. |
| 9. $4r + 2s$ by $2r + 9s$. | 17. $ax + by$ by $ax - by$. |
| 10. $3l + 5t$ by $2l + 6t$. | 18. $a^2 - ay + y^2$ by $a + y$. |
| 11. $4a + 3x$ by $4a - 3x$. | 19. $3a^2 - 6ab + 3b^2$ by $2a - 3b$. |

Multiply :

20. $2a^2 - 3b^2 - ab$ by $3a^2 - 4ab - 5b^2$.
21. $5x - 5x^2 + 10$ by $12 - 30x + 2x^2$.
22. $3n^2 + 3m^2 + mn$ by $m^3 - 2mn^2 + m^2n$.
23. $4y^2 - 10 + 2y$ by $2y^2 - 3y + 5$.
24. $4x - 3x^2 + 2x^3$ by $3x - 10x^2 + 10$.
25. $a^5 + a^4 + 4a^3 - a^2 + a$ by $a + 1$.
26. $31x^3 - 27x^2 + 9x - 3$ by $3x + 1$.
27. $4x^3 - 3x^2y + 5xy^2 - 6y^3$ by $5x + 6y$.
28. $a + b + c + d$ by $a - b - c + d$.
29. $a^2 + b^2 + c^2 - ab - ac - bc$ by $a + b + c$.
30. $ax^{2n} + ay^{2n}$ by $ax^{2n} - ay^{2n}$.
31. $ax^{n-1} + y^{n-1}$ by $3ax^{n-1} + 2y^{n-1}$.
32. $x^{2n} + 2x^ny^n + y^{2n}$ by $x^{2n} - 2x^ny^n + y^{2n}$.

An indicated product is said to be *expanded* when the multiplication is performed.

Expand :

33. $(x + y)(x + y)$.
34. $(a + b)(a + b)$.
35. $(c^3 + d^3)(c^3 + d^3)$.
36. $(x^n + y^n)(x^n + y^n)$.
37. $(3a + b)(3a + b)$.
38. $(3a + b)(3a - b)$.
39. $(2a^2 + b)(2a^2 - b)$.
40. $(x^n + y^n)(x^n - y^n)$.
41. $(3ax + 2by)(3ax + 2by)$.
42. $(3ax + 2by)(3ax - 2by)$.
43. $(4m - 5n)(4m + 5n)$.
44. $(a + b + c)(a + b - c)$.
45. $(a + b)(a - b)(a + b)(a - b)$.
46. $(a^2 + x^2)(a^2 - x^2)(a^2 + x^2)(a^2 - x^2)$.
47. $(a - b)(a + b)(a^2 + b^2)(a^4 + b^4)$.
48. $(a^m - b^n)(a^m + b^n)(a^{2m} + b^{2n})$.
49. $(2a + 3b + 5c)(2a + 3b - 5c)$.
50. $(5m - 2n + x)(5m - 2n - x)$.
51. $(x^n - nx^{n-1}y + \frac{1}{2}nx^{n-2}y^2)(x + y)$.
52. $(x^p + 3x^{p-1}z - 6x^{p-2}z^2)(x^2 - y^2)$.

88. When polynomials are arranged according to the ascending or the descending powers of some literal factor, processes may frequently be abridged by using the **Detached Coefficients**.

53. Expand $(2x^4 - 3x^3 + 3x + 1)(3x + 2)$.

FULL PROCESS	DETACHED COEFFICIENTS
$2x^4 - 3x^3 + 3x + 1$	2 -3 +0 +3 +1
$3x + 2$	3 +2
<hr/>	<hr/>
$6x^5 - 9x^4 \qquad + 9x^3 + 3x$	6 -9 +0 +9 +3
$\qquad 4x^4 - 6x^3 \qquad + 6x + 2$	$\qquad 4 \quad -6 \quad +0 \quad +6 \quad +2$
<hr/>	<hr/>
$6x^5 - 5x^4 - 6x^3 + 9x^2 + 9x + 2$	$6x^5 - 5x^4 - 6x^3 + 9x^2 + 9x + 2$

Since the second power of x is wanting in the first factor, the term, if it were supplied, would be $0x^2$, and the detached coefficient of the term would be 0.

The detached coefficients of missing terms should be supplied to prevent confusion in placing the coefficients in the partial products and to prevent errors in determining the result.

54. Multiply $a^4 - 2a^3b + 3a^2b^2 - 5ab^3 + 5b^4$ by $a + 2b$.

PROCESS	TEST
$1 - 2 + 3 - 5 + 5$	$= 2$
$1 + 2$	$= 3$
<hr/>	<hr/>
$1 - 2 + 3 - 5 + 5$	
$\qquad 2 - 4 + 6 - 10 + 10$	
<hr/>	
$1 + 0 - 1 + 1 - 5 + 10$	$= 6$
$= a^5 + 0a^4b - a^3b^2 + a^2b^3 - 5ab^4 + 10b^5$	
$= a^5 - a^3b^2 + a^2b^3 - 5ab^4 + 10b^5$	

Observe that the powers of a decrease uniformly from left to right, and that the powers of b increase uniformly from left to right.

Observe also that the sum of the exponents is the same in every term.

Expand the following, using detached coefficients, and test the results:

55. $(x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4)(x + y)$.

56. $(5x^8 - x^7 - 2x^5 + x^4 + 3x^3 - 1)(x + 2)$.

57. $(a^3 + 4a^2 + 16a - 32)(a^3 + a^2 + a + 1)$.
 58. $(p^2 - 2pq + q^2)(p^2 + 2pq + q^2)$.
 59. $(x - 1)(x - 2)(x - 3)(x + 4)(x + 2)$.
 60. $(15r^4s - 10rs^4 + r^5 + s^5 + 3r^3s^2 + 3r^2s^3)(r - 2s)$.
 61. $(x^{10} - x^9 + x^8 - x^7 + x^6 - x^5 + x^4 - x^3 + x^2 - x + 1)(x + 1)$.
 62. $(x^{10} + 3x^8 + 2x^6 + 5x^4 + 2x^2 + 2)(x^6 + x^4 + x^2 + 1)$.
 63. $(x^4 + x^3 + x^2 + x + 1)(x^4 - x^3 + x^2 - x + 1)(x + 1)(x - 1)$.
 64. $(x^7 - 4x^6y + 5x^5y^2 + 3x^3y^4 - x^2y^5 + xy^6 - y^7)(2x - 3y)$.

Expand:

65. $(2x^3 + 4x^2 + 8x + 16)(3x - 6)$.
 66. $(x^3 + 4x^2 + 5x - 24)(x^2 - 4x + 11)$.
 67. $(x^3 - 4x^2 + 11x - 24)(x^2 + 4x + 5)$.
 68. $(x^2 - x + 1)(x^2 + x + 1)(x^4 - x^2 + 1)$.
 69. $(16a^4 - 8a^3 + 4a^2 - 2a + 1)(2a + 1)$.
 70. $(a^3 - 2a^2 + 3a - 4)(4a^3 + 3a^2 + 2a + 1)$.
 71. $(m^4 + 2m^3 + m^2 - 4m - 11)(m^2 - 2m + 3)$.
 72. $(m^3 + 2m^2n + 2mn^2 + n^3)(m^3 - 2m^2n + 2mn^2 - n^3)$.
 73. $(x^3 - \frac{1}{2}x^2 + \frac{1}{4}x - \frac{3}{16})(x + \frac{1}{2})$.
 74. $(\frac{1}{2}x^3 + \frac{1}{3}x^2 + \frac{2}{3}x + 2)(x - \frac{2}{3})$.
 75. $\left(\frac{x^3}{8} - \frac{3x^2}{4} + \frac{3x}{2} - 1\right)(4x + 4)$.
 76. $\left(a^2 - \frac{2a}{3}\right)(3a^2 + 2a)(3a - 3)$.
 77. $(x^{n-1} - x^{n-2} + x^{n-3} - x^{n-4})(x + 1)$.
 78. $(1 - a^n + a^{2n} - a^{3n})(1 - a^n)$.
 79. $(a^{2n} + 2a^n b^n + b^{2n})(a^n - b^n)$.
 80. $(x^{2n} - x^n y^n + y^{2n})(x^n + y^n)$.

EQUATIONS AND PROBLEMS

89. 1. Given $5(2x - 3) = 7(3x + 5) - 72$, to find the value of x .

SOLUTION

$$5(2x - 3) = 7(3x + 5) - 72.$$

Expanding, $10x - 15 = 21x + 35 - 72.$

Transposing, $10x - 21x = 15 + 35 - 72.$

Uniting terms, $-11x = -22.$

Multiplying by -1 , $11x = 22.$

$$\therefore x = 2.$$

VERIFICATION. — Substituting 2 for x in the given equation,

$$5(4 - 3) = 7(6 + 5) - 72.$$

$$5 = 77 - 72 = 5.$$

Find the value of x , and verify the result, in the following:

2. $4 = 5 - (x - 2) - (x - 3).$
5. $10x - 2(x - 3) = -10.$
3. $2 = 2x - 5 - (x - 3).$
6. $6x - 3(x - 6) = 4(2x - 1) + 2.$
4. $1 = 5(2x - 4) + 5x + 6.$
7. $7(5 - 3x) = 3(3 - 4x) - 1.$
8. $4(2 - 4x) = 4 - 2(x + 5).$
9. $5 + 3x - 4 = 13 + 4(x - 4).$
10. $49 - 2(2x + 3) = 9 + 2(2x - 3).$
11. $3x - 2(4x - 5) = -2(6 + 2x).$
12. $3(x + 1) - 2(2x + 5) = 6(3 - x).$
13. $2(x - 5) + 7 = x + 30 - 9(x - 3).$
14. $5 + 7(x - 5) = 15(x + 2 - 36).$
15. $(x - 2)(x - 2) = (x - 3)(x - 3) + 7.$
16. $(x - 4)(x + 4) = (x - 6)(x + 5) + 25.$
17. $4x^3 - 4(x^3 - x^2 + x - 2) = 4x^2.$
18. $7(2x - 3b) = 2b - 3(2x + b).$
19. $11a = 3(x - 2a) - 5(2x - 2a).$
20. $3(2b - 4x) - (x - b) = -6b.$
24. $3(x - a - 2b) = 3b.$
21. $4x - x^2 = x(2 - x) + 2a.$
25. $5b = 3(2x - b) - 4b.$
22. $2(x + a) = 10c.$
26. $13(x - a) = 5(2x + a).$
23. $5c = 5(x + a - b).$
27. $5(4x - 3a) - 6(3x - 2a) = 3a.$

Solve the following problems and verify the solutions :

28. George and Henry together had 46 cents. If George had 4 cents more than half as many as Henry, how many cents had each ?

FIRST SOLUTION

Let x = the number of cents George had.
 Then, $x - 4$ = the number of cents George had less 4,
 and $2(x - 4)$ = the number of cents Henry had ;
 $\therefore x + 2(x - 4) = 46$.
 Expanding, $x + 2x - 8 = 46$;
 $\therefore x = 18$, the number of cents George had,
 and $2(x - 4) = 28$, the number of cents Henry had.

SECOND SOLUTION

Since George had 4 cents more than half as many as Henry,
 let $2x$ = the number of cents Henry had ;
 then, $x + 4$ = the number of cents George had,
 and $2x + x + 4 = 46$;
 $\therefore x = 14$,
 $2x = 28$, the number of cents Henry had,
 and $x + 4 = 18$, the number of cents George had.

VERIFICATION

The answers obtained should be tested by the conditions of the problem. If they satisfy the conditions of the problem, the solution is presumably correct.

1st condition : — They had together 46 cents.

$$18 + 28 = 46.$$

2d condition : — George had 4 cents more than half as many as Henry.

$$18 = \frac{1}{2} \text{ of } 28 + 4.$$

29. In a certain election at which 8000 votes were polled, B received 500 votes more than half as many as A. How many votes did each receive ?

30. A had \$40 more than B; B had \$10 more than one third as much as A. How much money had each ?

31. Mary is 25 years younger than her mother. If she were one year older, she would be $\frac{1}{4}$ as old as her mother. What is the age of each?

32. If John had 3 more marbles, he would have 3 times as many as Clarence. Both have 41 marbles. How many has each?

33. Two boys together had \$8.20, and one had 50 cents less than half as much as the other. What amount had each?

34. If 5 is subtracted from twice a certain number, and the difference is multiplied by 3, the result is 9 less than 5 times the number. What is the number?

35. A is $\frac{2}{3}$ as old as B; 8 years ago he was $\frac{1}{2}$ as old as B. What is the age of each?

SUGGESTION. — Let $5x = B$'s present age.

36. In 2 years A will be twice as old as he was 2 years ago. How old is he?

37. Two wheelmen start at the same time from A to ride to B. One rides at the rate of 10 miles an hour, and rests 3 hours; the other rides at the rate of 8 miles an hour, and by resting only 1 hour arrives at B as soon as the faster rider. How far is it from A to B, and how many hours are occupied in making the trip?

38. A man had two flocks of sheep with the same number of sheep in each. After selling 100 sheep from one flock, and 20 from the other, the numbers remaining were as 2 to 3. How many sheep had he in each flock at first?

39. Mary bought 17 apples for 61 cents. For a certain number of them she paid 5 cents each, and for the rest she paid 3 cents each. How many of each kind did she buy?

40. George is $\frac{1}{2}$ as old as his father; a years ago he was $\frac{1}{3}$ as old as his father. What is the age of each?

41. A rug is 3 feet longer than it is wide. When it is placed on the floor of a certain room, it leaves a margin of 2 feet on every side. If the area of the floor is 172 square feet greater than that of the rug, what are the dimensions of the floor?

SPECIAL CASES IN MULTIPLICATION

90. The square of the sum of two numbers.

$$(a + b)(a + b) = a^2 + 2ab + b^2.$$

$$(x + y)(x + y) = x^2 + 2xy + y^2.$$

1. When a number is multiplied by itself, what power is obtained? What is the second power, or square of $(a + b)$? of $(x + y)$?

2. How are the terms of the square of the sum of two numbers obtained from the numbers?

3. What signs have the terms?

91. PRINCIPLE. — *The square of the sum of two numbers is equal to the square of the first number, plus twice the product of the first and second, plus the square of the second.*

Since $5a^3 \times 5a^3 = 25a^6$, $3a^4b^5 \times 3a^4b^5 = 9a^8b^{10}$, etc., it is evident that in squaring a number the exponents of literal factors are doubled.

EXAMPLES

Expand by inspection:

- | | |
|--------------------------|-------------------------------|
| 1. $(m + n)(m + n)$. | 13. $(3b + c)^2$. |
| 2. $(p + q)(p + q)$. | 14. $(2a + 3b)^2$. |
| 3. $(r + s)(r + s)$. | 15. $(5x + 2y)^2$. |
| 4. $(a + x)(a + x)$. | 16. $(7z + 3c)^2$. |
| 5. $(x + 4)(x + 4)$. | 17. $(3b + 10x)^2$. |
| 6. $(m + 5)(m + 5)$. | 18. $(a^3 + b^3)^2$. |
| 7. $(a + 6)(a + 6)$. | 19. $(a^5 + b^5)^2$. |
| 8. $(y + 7)(y + 7)$. | 20. $(a^n + b^n)^2$. |
| 9. $(z + 1)(z + 1)$. | 21. $(x^m + y^n)^2$. |
| 10. $(2a + x)(2a + x)$. | 22. $(3a^2 + 5b^3)^2$. |
| 11. $(3m + n)(3m + n)$. | 23. $(1 + 2a^3b)^2$. |
| 12. $(5x + z)(5x + z)$. | 24. $(x^{n-1} + y^{n-1})^2$. |

25. Find the square of 41.

SOLUTION

$$41^2 = (40 + 1)^2 = 40^2 + 2 \times 40 \times 1 + 1^2 = 1681.$$

Square:

$$26. \ 21. \qquad 29. \ 45. \qquad 32. \ 22. \qquad 35. \ 81.$$

$$27. \ 24. \qquad 30. \ 83. \qquad 33. \ 72. \qquad 36. \ 91.$$

$$28. \ 25. \qquad 31. \ 65. \qquad 34. \ 43. \qquad 37. \ 101.$$

38. Find the square of $7\frac{1}{2}$.

SOLUTION

$$(7\frac{1}{2})^2 = (7 + \frac{1}{2})^2 = 7^2 + 2 \times 7 \times \frac{1}{2} + (\frac{1}{2})^2 = 49 + 7 + \frac{1}{4} = 56\frac{1}{4}.$$

Observe that the *middle term* of the square of any number expressed by an integer and the fraction $\frac{1}{2}$ is equal to the integer. Hence, the square of such a number is equal to the square of the integer, + the integer, + the square of the fraction. Observe also that the sum of the first two terms of the square may be found by multiplying the integer by the integer increased by 1.

$$\text{Thus,} \qquad (3\frac{1}{2})^2 = 9 + 3 + \frac{1}{4} = 12\frac{1}{4},$$

$$\text{or} \qquad (3\frac{1}{2})^2 = 3 \times 4 + \frac{1}{4} = 12\frac{1}{4}.$$

Find the square of

$$39. \ 5\frac{1}{2}. \qquad 41. \ 2\frac{1}{2}. \qquad 43. \ 1.5. \qquad 45. \ 6.5.$$

$$40. \ 4\frac{1}{2}. \qquad 42. \ 12\frac{1}{2}. \qquad 44. \ 5.5. \qquad 46. \ 8.5.$$

92. The square of the difference of two numbers.

$$(a - b)(a - b) = a^2 - 2ab + b^2.$$

$$(x - y)(x - y) = x^2 - 2xy + y^2.$$

1. What is the square of $(a - b)$? of $(x - y)$?

2. How is the square of the difference of two numbers obtained from the numbers?

3. How does the square of $(a - b)$ differ from the square of $(a + b)$?

93. PRINCIPLE. — *The square of the difference of two numbers is equal to the square of the first number, minus twice the product of the first and second, plus the square of the second.*

EXAMPLES

Expand by inspection:

- | | | |
|-------------------|------------------|-----------------------------|
| 1. $(x-m)(x-m)$. | 10. $(2a-x)^2$. | 19. $(3x-2)^2$. |
| 2. $(m-n)(m-n)$. | 11. $(3m-n)^2$. | 20. $(2x-5y)^2$. |
| 3. $(x-6)(x-6)$. | 12. $(4x-y)^2$. | 21. $(2x-4y)^2$. |
| 4. $(p-8)(p-8)$. | 13. $(5m-n)^2$. | 22. $(5m-3n)^2$. |
| 5. $(q-7)(q-7)$. | 14. $(m-4n)^2$. | 23. $(3p-5q)^2$. |
| 6. $(a-c)(a-c)$. | 15. $(p-3q)^2$. | 24. $(a^n-b^n)^2$. |
| 7. $(r-t)(r-t)$. | 16. $(a-7b)^2$. | 25. $(x^m-y^n)^2$. |
| 8. $(a-x)(a-x)$. | 17. $(4a-3)^2$. | 26. $(x^{n-1}-y^{n-1})^2$. |
| 9. $(x-1)(x-1)$. | 18. $(5x-4)^2$. | 27. $(mx^m-ny^n)^2$. |
28. Find the square of 19.

SOLUTION

$$19^2 = (20-1)^2 = 20^2 - 2 \times 20 \times 1 + 1^2 = 361.$$

Find the square of

- | | | | |
|---------|---------|---------|----------|
| 29. 49. | 32. 29. | 35. 67. | 38. 998. |
| 30. 69. | 33. 38. | 36. 89. | 39. 999. |
| 31. 79. | 34. 48. | 37. 99. | 40. 595. |

94. The square of any polynomial.

By actual multiplication,

$$(a+b+c)^2 = a^2 + b^2 + c^2 + 2ab + 2ac + 2bc.$$

$$(a+b-c+d)^2 = a^2 + b^2 + c^2 + d^2 + 2ab - 2ac + 2ad - 2bc + 2bd - 2cd.$$

$$(a + \dots + k + \dots + m)^2 = a^2 + \dots + k^2 + \dots + m^2 + 2ak + \dots + 2am + \dots + 2km + \dots$$

1. In the square of each polynomial what terms are squares?
2. How are the other terms formed from the terms of the polynomial?
3. What signs have the squares? How are the signs of the other terms determined?

95. PRINCIPLE. — *The square of a polynomial is equal to the sum of the squares of the terms and twice the product of each term by each term that follows it.*

EXAMPLES

Expand by inspection:

- | | | |
|--------------------------|---------------------------|-----------------------|
| 1. $(x + y + z)^2$. | 4. $(x - y + z)^2$. | 7. $(a - 2b + c)^2$. |
| 2. $(x + y - z)^2$. | 5. $(x + y - 3z)^2$. | 8. $(2a - b - c)^2$. |
| 3. $(x - y - z)^2$. | 6. $(x - y + 3z)^2$. | 9. $(b - 2a + c)^2$. |
| 10. $(ax - by + cz)^2$. | 15. $(2a - 5b + 3c)^2$. | |
| 11. $(ma + nb - rz)^2$. | 16. $(2m - 4n - r)^2$. | |
| 12. $(qb - pc - rd)^2$. | 17. $(12 - 2x + 3y)^2$. | |
| 13. $(ac - bd - de)^2$. | 18. $(a + m + b + n)^2$. | |
| 14. $(3x - 2y + 4z)^2$. | 19. $(a - m + b - n)^2$. | |

96. The product of the sum and difference of two numbers.

$$(a + b)(a - b) = a^2 - b^2.$$

$$(x^n + y^n)(x^n - y^n) = x^{2n} - y^{2n}.$$

1. Since $a + b$ represents the sum and $a - b$ the difference of two numbers, to what is the product of the sum and the difference of two numbers equal?

2. How are the terms of the product obtained from the numbers?

3. What sign connects the squares?

97. PRINCIPLE. — *The product of the sum and difference of two numbers is equal to the difference of their squares.*

EXAMPLES

Expand by inspection:

- | | |
|-----------------------|----------------------------|
| 1. $(x + y)(x - y)$. | 6. $(r + s)(r - s)$. |
| 2. $(a + c)(a - c)$. | 7. $(x + 1)(x - 1)$. |
| 3. $(m + n)(m - n)$. | 8. $(x^2 + 1)(x^2 - 1)$. |
| 4. $(p + q)(p - q)$. | 9. $(x^3 + 1)(x^3 - 1)$. |
| 5. $(p + 5)(p - 5)$. | 10. $(x^4 - 1)(x^4 + 1)$. |

- | | |
|--------------------------------|--|
| 11. $(x^5 - 1)(x^5 + 1)$. | 20. $(2x^3 + 5y^2)(2x^3 - 5y^2)$. |
| 12. $(x^a + y^3)(x^a - y^3)$. | 21. $(3x^6 + 2y^6)(3x^6 - 2y^6)$. |
| 13. $(ab + 5)(ab - 5)$. | 22. $(2a^2 + 2b^2)(2a^2 - 2b^2)$. |
| 14. $(cd + d^2)(cd - d^2)$. | 23. $(-5n - 2b)(-5n + 2b)$. |
| 15. $(2x + 3y)(2x - 3y)$. | 24. $(-x - 2y)(-x + 2y)$. |
| 16. $(3m + 4n)(3m - 4n)$. | 25. $(-5 - 3m)(-5 + 3m)$. |
| 17. $(12 + xy)(12 - xy)$. | 26. $(x^{m-1} + y^{n+1})(x^{m-1} - y^{n+1})$. |
| 18. $(3m^2n - b)(3m^2n + b)$. | 27. $(3x^m + 7y^n)(3x^m - 7y^n)$. |
| 19. $(ab + cd)(ab - cd)$. | 28. $(5a^3b^2 + 2x^r)(5a^3b^2 - 2x^r)$. |

One or both of the numbers may consist of more than one term.

29. Expand $(a + m - n)(a - m + n)$.

SOLUTION

$$a + m - n = a + (m - n).$$

$$a - m + n = a - (m - n).$$

$$\therefore [a + m - n][a - m + n] = [a + (m - n)][a - (m - n)].$$

$$\text{Prin.,} \quad = a^2 - (m - n)^2$$

$$\S 93, \quad = a^2 - (m^2 - 2mn + n^2)$$

$$= a^2 - m^2 + 2mn - n^2.$$

Expand:

- | | |
|--------------------------------|--------------------------------------|
| 30. $(a + x - y)(a - x + y)$. | 36. $(y + c + d)(y + c - d)$. |
| 31. $(x + c - d)(x - c + d)$. | 37. $(a + x + y)(a + x - y)$. |
| 32. $(r + p - q)(r - p + q)$. | 38. $(x^2 + 2x + 1)(x^2 + 2x - 1)$. |
| 33. $(r + p + q)(r - p - q)$. | 39. $(x^2 + 2x - 1)(x^2 - 2x + 1)$. |
| 34. $(x + b + n)(x - b - n)$. | 40. $(x^2 + 3x - 2)(x^2 - 3x + 2)$. |
| 35. $(y + c + d)(y - c - d)$. | 41. $(x^2 + 3x + 2)(x^2 - 3x + 2)$. |

$$42. (m^4 - 2m^2 + 1)(m^4 + 2m^2 + 1).$$

$$43. (2x + 3y - 4z)(2x + 3y + 4z).$$

$$44. (2x^2 - xy + 3y^2)(2x^2 + xy - 3y^2).$$

$$45. (x^2 + xy + y^2)(x^2 - xy + y^2).$$

$$46. [(a+b)+(c+d)][(a+b)-(c+d)].$$

$$47. (a+b+x+y)(a+b-x-y).$$

$$48. (a+b+m-n)(a+b-m+n).$$

$$49. (x-m+y-n)(x-m-y+n).$$

$$50. (p-q+r+s)(p-q-r-s).$$

$$51. (a-m-b-n)(a+m-b+n).$$

$$52. (a+x+b-y)(a-x+b+y).$$

53. Find the product of 32×28 .

SOLUTION

$$\begin{aligned} 32 \times 28 &= (30+2)(30-2) \\ &= 30^2 - 2^2 = 900 - 4 = 896. \end{aligned}$$

Find the product of

$$54. 31 \times 29.$$

$$57. 48 \times 52.$$

$$60. 98 \times 102.$$

$$55. 42 \times 38.$$

$$58. 57 \times 63.$$

$$61. 99 \times 101.$$

$$56. 69 \times 71.$$

$$59. 95 \times 85.$$

$$62. 95 \times 105.$$

63. What is the square of 95?

SOLUTION

$$(a+b)(a-b) = a^2 - b^2. \quad (1)$$

Transposing, etc.,

$$a^2 = (a+b)(a-b) + b^2. \quad (2)$$

Let

$$a = 95 \text{ and } b = 5.$$

Equation (2) becomes

$$\begin{aligned} 95^2 &= (95+5)(95-5) + 5^2 \\ &= 100 \times 90 + 25 = 9025. \end{aligned}$$

64. What is the square of 48?

SOLUTION

Let

$$a = 48 \text{ and } b = 2.$$

Equation (2) becomes

$$\begin{aligned} 48^2 &= (48+2)(48-2) + 2^2 \\ &= 50 \times 46 + 4 = 2304. \end{aligned}$$

Square by a similar process:

$$65. 98.$$

$$67. 93.$$

$$69. 58.$$

$$71. 87.$$

$$73. 68.$$

$$66. 96.$$

$$68. 97.$$

$$70. 49.$$

$$72. 79.$$

$$74. 129.$$

98. The product of two binomials that have a common term.

$$\begin{aligned}(x+2)(x+5) &= x^2 + 2x + 5x + 10 \\ &= x^2 + 7x + 10.\end{aligned}$$

$$\begin{aligned}(x+2)(x-5) &= x^2 + 2x - 5x - 10 \\ &= x^2 - 3x - 10.\end{aligned}$$

$$\begin{aligned}(x-2)(x-5) &= x^2 - 2x - 5x + 10 \\ &= x^2 - 7x + 10.\end{aligned}$$

$$\begin{aligned}(x+a)(x-b) &= x^2 + ax - bx - ab \\ &= x^2 + (a-b)x - ab.\end{aligned}$$

1. How many terms are alike in each pair of factors?
2. How is the first term of each product obtained from the binomial factors?
3. How is the third term of each product obtained from the factors?
4. How is the second term of the product in the first example obtained from the factors? in the second example? in the third example? in the fourth example?
5. How can the formation of the second, or middle term be described so as to apply to all of the examples?

99. PRINCIPLE. — *The product of two binomials that have a common term is the algebraic sum of the square of the common term, the common term multiplied by the algebraic sum of the unlike terms, and the algebraic product of the unlike terms.*

EXAMPLES

Expand by inspection:

- | | |
|-------------------|---------------------|
| 1. $(x+5)(x+6)$. | 7. $(x-5)(x-1)$. |
| 2. $(x+7)(x+8)$. | 8. $(x+5)(x+8)$. |
| 3. $(x-7)(x+8)$. | 9. $(p-4)(p+1)$. |
| 4. $(x+7)(x-8)$. | 10. $(r-3)(r-1)$. |
| 5. $(x-5)(x-4)$. | 11. $(m-6)(m+5)$. |
| 6. $(x-3)(x-2)$. | 12. $(m-2)(m+10)$. |

- | | |
|--|-------------------------------|
| 13. $(n-8)(n-12)$. | 24. $(y-2a)(y+3b)$. |
| 14. $(n-6)(n+15)$. | 25. $(z-4a)(z+3a)$. |
| 15. $(x^2+5)(x^2-3)$. | 26. $(2x+5)(2x+3)$. |
| 16. $(x^3-7)(x^2+6)$. | 27. $(2x-7)(2x+5)$. |
| 17. $(x^4-3)(x^4+9)$. | 28. $(3y-1)(3y+2)$. |
| 18. $(x+c)(x+d)$. | 29. $(4x^2+1)(4x^2-7)$. |
| 19. $(m+a)(m+b)$. | 30. $(ab-6)(ab+4)$. |
| 20. $(r+a)(r-b)$. | 31. $(x^2y^2-a)(x^2y^2+2a)$. |
| 21. $(s-a)(s+n)$. | 32. $(2m-ab)(2m+3ab)$. |
| 22. $(x^a-5)(x^a+4)$. | 33. $(5p-ac^2)(2ac^2+5p)$. |
| 23. $(x^a-a)(x^a-b)$. | 34. $(3xy+y^2)(y^2-xy)$. |
| 35. $(\overline{a+b+5})(\overline{a+b+2})$. | |
| 36. $(\overline{a-b-4})(\overline{a-b+10})$. | |
| 37. $(\overline{x+y-1})(\overline{x+y+2})$. | |
| 38. $(\overline{x-y-2})(\overline{x-y-8})$. | |
| 39. $(\overline{x^2+x-1})(\overline{x^2+x+3})$. | |
| 40. $(\overline{2m+n-3})(\overline{2m+n+4})$. | |

By an extension of the method given above, the product of any two binomials may be written.

41. Expand $(3x+2y)(5x-4y)$.

SOLUTION

$$\begin{aligned}(3x+2y)(5x-4y) &= 15x^2 + 10xy - 12xy - 8y^2 \\ &= 15x^2 - 2xy - 8y^2.\end{aligned}$$

Expand by inspection:

- | | |
|------------------------|--------------------------|
| 42. $(2x+5y)(3x+4y)$. | 45. $(3a^2-1)(2a^2+3)$. |
| 43. $(3x-4y)(2x+5y)$. | 46. $(m^2+n)(2m^2-n)$. |
| 44. $(3a-6b)(2a+3b)$. | 47. $(a+2b)(c-2d)$. |

EXERCISES

100. 1. In a certain family there are three children each of whom is 2 years older than the one next younger. When the youngest is x years old, what are the ages of the others? When the oldest is y years old, what are the ages of the others?

2. What two whole numbers are nearest to the whole number x ?

3. Mary read 10 pages in a book, stopping at the top of page a . On what page did she begin to read?

4. A man made three purchases of a , b , and 2 dollars, respectively, and tendered a 10-dollar bill. Express the number of dollars change due him.

5. A sold B grain, hay, and potatoes for a , b , and c dollars, respectively; but some of the grain becoming damaged, and some of the potatoes having been frozen, he deducted $x + y$ dollars from B's indebtedness. If B offered a 100-dollar bill in payment, what was the amount due him in return?

6. What is the cost of 5 apples at b cents each? What will a apples cost at b cents each?

7. How many cents are there in a dollars? How many dimes are there in b dollars? in ax dollars?

8. If a man earns \$2 per day, how much will he earn in a days? in c days? in $a - c$ days?

9. How much will a man whose wages are a dollars per day earn in b days? in c days? in x days? in a days?

10. If a man earns a dollars per month and his expenses are b dollars per month, how much will he save in a year?

11. How far can a wheelman ride in a hours at the rate of b miles an hour? How far will he have ridden after a hours, if he stops c hours of the time to rest?

12. A has x dollars and B y dollars. If A gives B m dollars, how much will each then have?

13. The number 25 may be written $20 + 5$. Write the number whose first digit is x and second y .

14. A book contained x pages. If they averaged y lines to a page and z letters to a line, how many letters were there in the book?

15. How many square rods are there in a square field one of whose sides is $2b$ rods long? x rods long?

16. What is the number of square rods in a rectangular field whose length is 30 rods and width 20 rods? What will be the area in square rods, if the length is a rods and the width b rods?

17. A fence is built across a rectangular field so as to make the part on one side of the fence a square. If the field is a rods long and b rods wide, what is the area of each part?

18. A man who had s dollars bought b bales of hay at n cents a bale and a bushels of oats at m cents a bushel. How many cents had he left?

19. A speculator bought a car loads of wheat at m dollars a car, and sold b car loads of it at n dollars a car. How much did he gain by the transaction, if he sold the rest of the wheat for c dollars a car?

20. A sold a farm which had cost him n dollars an acre to two men, a acres to one and b acres to the other, at the uniform price of m dollars an acre. How much did he gain?

21. In a library there were $p + q$ volumes that averaged $p + q$ pages per volume, $p + q$ words per page, and 7 letters per word. How many letters were there in all these books?

22. A wheelman who had a journey of x miles to make rode the first a hours at the rate of b miles an hour, when he was obliged to stop c hours for repairs. After that he rode 2 miles an hour faster, so that he made the whole journey in 10 hours. What was the length of the journey?

23. A wheelman rode a hours at the rate of m miles an hour, then decreased his speed 5 miles an hour for 3 hours, and finished his journey in b hours more, increasing his first rate 2 miles an hour. How far did he ride?

If $a = 4$, $b = 2$, and $m = 10$, how many miles did he ride, and in what time did he accomplish the journey?

DIVISION

101. 1. Since $+2$ times $+10$ is $+20$, if $+20$ is divided by $+10$, what is the sign of the quotient?

2. What is the sign of the quotient when a positive number is divided by a positive number?

3. Since $+2$ times -10 is -20 , if -20 is divided by -10 , what is the sign of the quotient? if -40 is divided by -5 ?

4. What is the sign of the quotient when a negative number is divided by a negative number?

5. What is the sign of the quotient when the dividend and divisor have like signs?

6. Since $+2$ times -10 is -20 , if -20 is divided by $+2$, what is the sign of the quotient? if -20 is divided by $+5$?

7. Since -2 times -10 is $+20$, if $+20$ is divided by -2 , what is the sign of the quotient? if $+20$ is divided by -4 ?

8. What is the sign of the quotient when the dividend and divisor have unlike signs?

9. How many times is $2a$ contained in $6a$? in $10a$?
How is the coefficient of the quotient obtained?

10. Since $a^3 \times a^2 = a^5$, if a^5 is divided by a^3 , what is the quotient? What is the quotient, if a^5 is divided by a^2 ?

What is the quotient, if b^7 is divided by b^3 ? by b^4 ?

How is the exponent of a number in the quotient obtained?

11. What is the exponent of a in the quotient of $a^4 \div a^4$? of $a^3 \div a^3$? How many times is a^4 contained in a^4 ? a^3 in a^3 ?

12. What is the value of any expression whose exponent is 0?

102. In multiplication two numbers are given and their product is to be found. The *inverse* process, finding one of two numbers when their product and the other number are given, is called **Division**.

$$10 \div 2 = 5 \text{ and } D \div d = q$$

are inverses of

$$5 \times 2 = 10 \text{ and } q \times d = D.$$

The dividend corresponds to the product, the divisor to the multiplier, and the quotient to the multiplicand. Hence, the *quotient* may be defined as *that number which multiplied by the divisor produces the dividend*.

The quotient of a divided by b , indicated by $(a \div b)$, or $\frac{a}{b}$, is defined for all values of a and b by the relation

$$\frac{a}{b} \times b = a.$$

103. PRINCIPLES. — 1. **Law of Signs.** — *The sign of the quotient is + when the dividend and divisor have like signs, and - when they have unlike signs.*

2. **Law of Coefficients.** — *The coefficient of the quotient is equal to the coefficient of the dividend divided by that of the divisor.*

3. **Law of Exponents.** — *The exponent of a letter in the quotient is equal to its exponent in the dividend diminished by its exponent in the divisor.*

An expression whose exponent is 0 is equal to 1.

The Law of Signs may be established as follows:

$$\text{Since } +a \times +b = +ab, \quad +ab \div +b = +a.$$

$$\text{Since } +a \times -b = -ab, \quad -ab \div -b = +a.$$

$$\text{Since } -a \times +b = -ab, \quad -ab \div +b = -a.$$

$$\text{Since } -a \times -b = +ab, \quad +ab \div -b = -a.$$

The Law of Exponents or the Index Law for Division may be established as follows, m and n being *positive integers* and m being greater than n :

$$\text{By § 24, } a^m = a \times a \times a \dots \text{ to } m \text{ factors,}$$

$$a^n = a \times a \times a \dots \text{ to } n \text{ factors;}$$

$$\begin{aligned} \therefore a^m \div a^n &= (a \times a \times a \dots \text{ to } m \text{ factors}) \div (a \times a \times a \dots \text{ to } n \text{ factors}) \\ &= a \times a \times a \dots \text{ to } (m - n) \text{ factors.} \end{aligned}$$

$$\text{Hence, } a^m \div a^n = a^{m-n}.$$

104. Commutative, Associative, and Distributive Laws for Division.**1. The Commutative Law may be established as follows:**

Since, by the definition of division, § 102, $a = a \div c \times c$,

$$\begin{aligned} a \times b \div c &= a \div c \times c \times b \div c \\ \text{§ 82,} \qquad \qquad \qquad &= a \div c \times b \times c \div c \\ &= a \div c \times b. \end{aligned} \tag{1}$$

Also, § 102,

$$\begin{aligned} a \div b \div c &= a \div c \times c \div b \div c \\ \text{by notation, § 29,} \qquad \qquad &= \overline{(a \div c \times c \div b)} \div c \\ \text{by (1),} \qquad \qquad \qquad &= \overline{(a \div c \div b \times c)} \div c \\ \text{by notation,} \qquad \qquad \qquad &= a \div c \div b \times c \div c \\ &= a \div c \div b. \end{aligned} \tag{2}$$

The Commutative Law for division is expressed by (2).

(2) may be written $\frac{a}{b} \div c = \frac{a}{c} \div b$. (1) may be written $\frac{ab}{c} = \frac{a}{c} \times b$.

It follows from (1) and (2) that *in a succession of multiplications and divisions the multipliers and divisors may be arranged in any order.*

2. The Associative Law may be established as follows:

By the Commutative Law just proved,

$$\begin{aligned} a \times b \div c &= b \div c \times a \\ \text{by notation, § 29,} \qquad \qquad \qquad &= (b \div c) \times a \\ \text{by the Commutative Law,} \qquad \qquad &= a \times (b \div c). \end{aligned} \tag{3}$$

Also

$$\begin{aligned} a \div b \div c &= b \times c \div (b \times c) \times a \div b \div c \\ \text{by the Commutative Law,} \qquad \qquad &= b \div b \times c \div c \times a \div (b \times c) \\ &= a \div (b \times c). \end{aligned} \tag{4}$$

The Associative Law for division is expressed by (4).

(4) may be written $\frac{a}{b} \div c = \frac{a}{bc}$. (3) may be written $\frac{ab}{c} = a \times \frac{b}{c}$.

It follows from (3) and (4) that *in a succession of multiplications and divisions the multipliers and divisors may be grouped in any manner, each element keeping its own sign, \times or \div , if the sign \times precedes the sign of grouping, but changing it, if the sign \div precedes the sign of grouping.*

3. The Distributive Law may be established as follows:

$$\begin{aligned} \text{§ 85,} \qquad \qquad \qquad &(\overline{a \div m} + \overline{b \div m}) \times m = \overline{a \div m} \times m + \overline{b \div m} \times m \\ &= a \div b. \end{aligned}$$

Dividing both members by m ,

$$\overline{a \div m} + \overline{b \div m} = (a \div b) \div m;$$

that is,

$$\frac{a \div b}{m} = \frac{a}{m} \div \frac{b}{m}.$$

105. The **Reciprocal** of a number is 1 divided by the number.

The reciprocal of 5 is $\frac{1}{5}$; of b , $\frac{1}{b}$; of $(a + b)$, $\frac{1}{a + b}$.

106. $a \div b = a \times \frac{1}{b}$

by the Associative Law, $= a \times (\frac{1}{b} \times b)$.

Hence, *dividing by a number is equivalent to multiplying by the reciprocal of the number.*

107. To divide a monomial by a monomial.

EXAMPLES

1. Divide $-18a^5b^3$ by $6ab^2$.

PROCESS **EXPLANATION.**—Since the dividend and divisor have unlike signs, the sign of the quotient is $-$ (Prin. 1).

$$\begin{array}{r} 6ab^2 \overline{) -18a^5b^3} \\ \underline{-3a^4b} \end{array}$$
Then, -18 divided by 6 is -3 (Prin. 2); a^5 divided by a is a^4 (Prin. 3); and b^3 divided by b^2 is b (Prin. 3). Therefore, the quotient is $-3a^4b$.

RULE.—*Divide the numerical coefficient of the dividend by the numerical coefficient of the divisor, and to the result annex the letters, each with an exponent equal to its exponent in the dividend minus its exponent in the divisor.*

Write the sign $+$ before the quotient when the dividend and divisor have like signs, and the sign $-$ when they have unlike signs.

	2.	3.	4.	5.	6.
Divide	$12x$	$-12x^2y^2$	$35xy^3$	$-14x^2y^2$	$-26a^2b^3c^2$
By	$4x$	$4x^2y^2$	$-7xy$	$-2x^2y$	$13ab^2c$

Find the quotient of

- | | |
|-----------------------------------|--------------------------------------|
| 7. $28a^4b^2c \div -7abc$. | 10. $-14x^6y^2z^4 \div 7x^4y^2z^2$. |
| 8. $-16x^3y^2z^3 \div 4xyz$. | 11. $-27m^7n^8p^9 \div 9m^2n^3p^6$. |
| 9. $-36a^4m^2n^3 \div -9m^2n^2$. | 12. $-39q^3r^2p \div -13qp$. |

108. To divide a polynomial by a monomial.

EXAMPLES

1. Divide $5x^4y^4 - 10x^3y^5 + 5x^2y^6$ by $5x^2y^3$.

PROCESS. § 104, 3,
$$\begin{array}{r} 5x^2y^3 \overline{) 5x^4y^4 - 10x^3y^5 + 5x^2y^6} \\ \underline{x^2y - 2xy^2 + y^3} \end{array}$$

RULE. — *Divide each term of the dividend by the divisor, and find the algebraic sum of the partial quotients.*

Find the quotient of

$$2. \frac{4a^2b^3 - 12a^3b^2 + 16a^4b}{4a^2b}$$

$$6. \frac{4m^3n - 8m^2n^2 + 4mn^3}{4mn}$$

$$3. \frac{24a^5b^2 + 32a^5b^3 - 40a^4b^4}{8a^4b^2}$$

$$7. \frac{5x^4y - 10x^2y^2 + 20x^2y^3}{5xy}$$

$$4. \frac{-35x^2y^3z^4 + 45x^4y^3z^3}{5x^2y^3z}$$

$$8. \frac{-a - b - c - d - e}{-1}$$

$$5. \frac{-39x^2y^4z^6 + 65x^2y^5z^7}{-13x^2y^4z^6}$$

$$9. \frac{-a - a^2b - a^3c - a^4d - a^5e}{-a}$$

$$10. \frac{25r^8s^6 - 125r^6s^8 - 75r^4s^{10}}{5r^4s^6}$$

$$11. \frac{27c^5d^2 - 39c^4d^3 - 42c^3d^4}{3c^3d^2}$$

$$12. (34a^2x^6y^2 - 51a^4x^4y^4 - 68a^6x^2y^6) \div 17a^2x^2y^2$$

$$13. (8a^7b^3 - 28a^6b^4 - 16a^5b^5 + 4a^4b^6) \div 4a^4b^3$$

$$14. [a(b-c)^3 - b(b-c)^2 + c(b-c)] \div (b-c)$$

$$15. [(x-y) - 3(x-y)^2 + 4x(x-y)^3] \div (x-y)$$

$$16. (x^a - 2x^{a+1} - 5x^{a+2} - x^{a+3} + 3x^{a+4}) \div x^a$$

$$17. (y^{n+1} - 2y^{n+2} - y^{n+3} - 3y^{n+4} + y^{n+5}) \div y^{n+1}$$

$$18. (x^n - x^{n-1} + x^{n-2} - x^{n-3} + x^{n-4} - x^{n-5}) \div x^2$$

109. To divide a polynomial by a polynomial.

EXAMPLES

1. Divide $3x^2 + 35 + 22x$ by $x + 5$.

	PROCESS	TEST
	$3x^2 + 22x + 35$	
$3x(x+5)$. . .	$\begin{array}{r l} 3x^2 + 22x + 35 & x + 5 \\ \hline 3x^2 + 15x & 3x + 7 \\ \hline 7x + 35 & \end{array}$	$+ 60 \div + 6$ $= + 10$
$7(x+5)$. . .	$7x + 35$	

2. Divide $2a^4 - 5a^3b + 6a^2b^2 - 4ab^3 + b^4$ by $a^2 - ab + b^2$.

PROCESS	TEST
$ \begin{array}{r} 2a^4 - 5a^3b + 6a^2b^2 - 4ab^3 + b^4 \\ \underline{2a^4 - 2a^3b + 2a^2b^2} \\ -3a^3b + 4a^2b^2 - 4ab^3 \\ \underline{-3a^3b + 3a^2b^2 - 3ab^3} \\ a^2b^2 - ab^3 + b^4 \\ \underline{a^2b^2 - ab^3 + b^4} \\ 0 \end{array} $	$ \begin{array}{r} a^2 - ab + b^2 \quad 0 + 1 \\ \hline 2a^2 - 3ab + b^2 \quad = 0 \end{array} $

3. Divide $a^4 + 9a^2 + 81$ by $a^2 - 3a + 9$.

PROCESS	TEST
$ \begin{array}{r} a^4 + 9a^2 + 81 \\ \underline{a^4 - 3a^3 + 9a^2} \\ 3a^3 + 81 \\ \underline{3a^3 - 9a^2 + 27a} \\ 9a^2 - 27a + 81 \\ \underline{9a^2 - 27a + 81} \\ 0 \end{array} $	$ \begin{array}{r} a^2 - 3a + 9 \quad 91 \div 7 \\ \hline a^2 + 3a + 9 \quad = 13 \end{array} $

Divide, and test the results:

4. $x^2 - x - 20$ by $x - 5$.
5. $x^2 + 7x + 12$ by $x + 3$.
6. $m^3 - 3m - 18$ by $m - 6$.
7. $l^4 - 6l^2 - 16$ by $l^2 + 2$.
8. $cd - d^2 + 2c^2$ by $c + d$.
9. $x^2 - 11x + 10$ by $x - 10$.
10. $x^2 + 15x + 54$ by $x + 6$.
11. $r^{2n} + 11r^n + 30$ by $r^n + 6$.
12. $a^2m^4 - 4am^3 + 3m^2$ by $am - 1$.
13. $6a^2 + 13ab + 6b^2$ by $3a + 2b$.
14. $a^4 + 16 + 4a^2$ by $2a + a^2 + 4$.
15. $a^5 + a^4 + a^3 + a^2 + 3a - 1$ by $a + 1$.
16. $20x^2y - 25x^2 - 18y^2 + 27xy^2$ by $6y - 5x$.

17. $ax^3 - a^2x^2 - bx^2 + b^2$ by $ax - b$.

18. $a^5 - 41a - 120$ by $a^2 + 4a + 5$.

19. $x^5 - 61x - 60$ by $x^2 - 2x - 3$.

20. $25x^5 - x^3 - 8x - 2x^2$ by $5x^2 - 4x$.

21. $4y^4 - 9y^2 + 6y - 1$ by $2y^2 + 3y - 1$.

22. $a^4 - 4a^3x + 6a^2x^2 - 4ax^3 + x^4$ by $a^2 - 2ax + x^2$.

23.
$$\begin{array}{r} a^5 - 1 \\ a^5 + a^4 \\ \hline -a^4 - 1 \\ -a^4 - a^3 \\ \hline a^3 - 1 \\ a^3 + a^2 \\ \hline -a^2 - 1 \\ -a^2 - a \\ \hline a - 1 \\ a + 1 \\ \hline -2 \end{array} \quad \left| \begin{array}{l} a + 1 \\ a^4 - a^3 + a^2 - a + 1 + \frac{-2}{a + 1} \end{array} \right.$$

24.
$$\begin{array}{r} x^3 + x - 25 \\ x^3 - 3x^2 \\ \hline 3x^2 + x \\ 3x^2 - 9x \\ \hline 10x - 25 \\ 10x - 30 \\ \hline 5 \end{array} \quad \left| \begin{array}{l} x - 3 \\ x^2 + 3x + 10 + \frac{5}{x - 3} \end{array} \right.$$

25.
$$\begin{array}{r} x^4 - 3x^3 + x^2 + 2x - 1 \\ x^4 - x^3 - 2x^2 \\ \hline -2x^3 + 3x^2 + 2x \\ -2x^3 + 2x^2 + 4x \\ \hline x^2 - 2x - 1 \\ x^2 - x - 2 \\ \hline -x + 1 \end{array} \quad \left| \begin{array}{l} x^2 - x - 2 \\ x^2 - 2x + 1 + \frac{-x + 1}{x^2 - x - 2} \end{array} \right.$$

Divide:

26. $m^5 + n^5$ by $m + n$.

27. $x^5 + 32$ by $x + 2$.

28. $x^6 + y^6$ by $x^2 + y^2$.

29. $a^6 + 5a^5 - a^3 + 2a + 3$ by $a - 1$.

30. $2n^5 - 4n^4 - 3n^3 + 7n^2 - 3n + 2$ by $n - 2$.

31. $x^3 + y^3 + z^3 - 3xyz$ by $x + y + z$.

32. $m^3 + n^3 + x^3 + 3m^2n + 3mn^2$ by $m + n + x$.

33. $a^3 - 6a^2 + 12a - 8 - b^3$ by $a - 2 - b$.

34. $y^5 + 3y^4 + 5y^3 + 3y^2 + 3y + 5$ by $y + 1$.

35. $2x^5 - x^7 + 2x^4 - x^3 + x^2 + 5$ by $x + 1$.

36. $x^7 + 2x^6 - 2x^4 + 2x^2 - 1$ by $x + 1$.

37. $a^3 - 2a^2c + 4ac^2 - ax^2 - 4c^2x + 2cx^2$ by $a - x$.

38. $a^3 - b^3 + c^3 + 3abc$ by $a^2 + b^2 + c^2 + ab - ac + bc$.

39. $x^n + y^n$ by $x + y$ to five terms of the quotient.

40. $x^5 - 6x + 5$ by $x^2 - 2x + 1$, using detached coefficients.

PROCESS

$$\begin{array}{r}
 1 + 0 + 0 + 0 + 0 - 6 + 5 \\
 \underline{1 - 2 + 1} \\
 2 - 1 + 0 \\
 \underline{2 - 4 + 2} \\
 3 - 2 + 0 \\
 \underline{3 - 6 + 3} \\
 4 - 3 - 6 \\
 \underline{4 - 8 + 4} \\
 5 - 10 + 5 \\
 \underline{5 - 10 + 5}
 \end{array}
 \quad
 \begin{array}{r}
 1 - 2 + 1 \\
 \hline
 1 + 2 + 3 + 4 + 5 \\
 x^4 + 2x^3 + 3x^2 + 4x + 5
 \end{array}$$

Divide, using detached coefficients when convenient:

41. $x^3 + 8x + 7$ by $x^2 + 2x + 1$.

42. $a^6 + 38a + 12$ by $a + 2$.

43. $m^5 - 19m - 6$ by $m + 2$.

44. $m^7 - 32m^2 - 4m + 8$ by $m - 2$.

45. $a^6 + 27a^2 - 9a - 10$ by $a^2 - 3a + 5$.

46. $21x^4 - 29x^3 - 8x^2 + 6x + 4$ by $3x - 2$.

47. $2x^4 - 11x^3 + 16x^2 - 12x + 9$ by $2x - 3$.

48. $30x^4 - 62x^3 + 60x^2 - 36x + 8$ by $5x - 2$.

49. $27x^4 - 33x^3 + 46x^2 - 119x + 55$ by $9x - 5$.

50. $x^7 - 2x^5 - x^3 - 10x - 36$ by $x - 2$.

51. $x^4 - 4x^3 + 5x^2 - 4x + 1$ by $x^2 - x + 1$.

52. $x^4 - x^3 - 10x^2 + 7x + 15$ by $x^2 - 2x - 3$.

53. $2x^4 + 7x^3 - 27x^2 - 8x + 16$ by $x^2 + 5x - 4$.

54. $28x^4 + 6x^3 + 6x^2 - 6x - 2$ by $4x^2 + 2x + 2$.

55. $7x^5 - 6x^4 + 2x^3 - x - 2$ by $6x^3 + 5x + 2$.

56. $25x^5 - 20x^4 + 3x^3 + 16x - 6$ by $3x^3 - 8x + 2$.

57. $3x^4 + 7x^3 + 6x^2 + 3x - 1$ by $x^2 + x + 1$.

58. $6x^4 - 23x^3 + 30x^2 - 18x + 4$ by $2x^2 - 5x + 2$.

59. $24x^4 + 32x^3 - 16x^2 - 25x - 4$ by $6x^2 - x - 4$.

60. $x^5 - 2x^4 + \frac{1}{12}x^3 + \frac{2}{3}x^2 + \frac{1}{15}x + \frac{5}{4}$ by $x - \frac{3}{2}$.

61. $x^5 - \frac{5}{4}x^4 + \frac{3}{2}x^3 - \frac{3}{4}x^2 + \frac{5}{6}x - \frac{1}{4}$ by $x - \frac{3}{4}$.

62. $x^6 - \frac{1}{6}x^5 + \frac{5}{6}x^4 - \frac{1}{6}x^3 + \frac{3}{8}x^2 - \frac{1}{6}x + \frac{7}{24}$ by $x^2 - \frac{2}{3}x + \frac{1}{2}$.

63. $\frac{1}{3}x^3 + \frac{1}{2}y^3 + z^3 - \frac{1}{2}xyz$ by $\frac{1}{2}x + \frac{1}{3}y + z$.

64. $\frac{1}{1+x}$ to five terms.

65. $\frac{1}{1-x}$ to five terms.

66. $x^{2n-3} + y^{2n+3}$ by $x^{n-1} + y^{n+1}$.

SPECIAL CASES IN DIVISION

110. By actual division,

$$1. \begin{cases} \frac{x^2 - y^2}{x - y} = x + y. \\ \frac{x^3 - y^3}{x - y} = x^2 + xy + y^2. \\ \frac{x^4 - y^4}{x - y} = x^3 + x^2y + xy^2 + y^3. \\ \frac{x^5 - y^5}{x - y} = x^4 + x^3y + x^2y^2 + xy^3 + y^4. \end{cases}$$

From the above we infer that the difference of the same powers of two numbers is divisible by the difference of the numbers.

$$2. \begin{cases} \frac{x^2 - y^2}{x + y} = x - y. \\ \frac{x^3 - y^3}{x + y} = x^2 - xy + y^2, \text{ Rem., } -2y^3. \\ \frac{x^4 - y^4}{x + y} = x^3 - x^2y + xy^2 - y^3. \\ \frac{x^5 - y^5}{x + y} = x^4 - x^3y + x^2y^2 - xy^3 + y^4, \text{ Rem., } -2y^5. \end{cases}$$

From the above we infer that the difference of the same powers of two numbers is divisible by the sum of the numbers only when the powers are even.

$$3. \begin{cases} \frac{x^2 + y^2}{x - y} = x + y, \text{ Rem., } 2y^2. \\ \frac{x^3 + y^3}{x - y} = x^2 + xy + y^2, \text{ Rem., } 2y^3. \\ \frac{x^4 + y^4}{x - y} = x^3 + x^2y + xy^2 + y^3, \text{ Rem., } 2y^4. \\ \frac{x^5 + y^5}{x - y} = x^4 + x^3y + x^2y^2 + xy^3 + y^4, \text{ Rem., } 2y^5. \end{cases}$$

From the above we infer that the sum of the same powers of two numbers is not divisible by the difference of the numbers.

$$4. \begin{cases} \frac{x^2 + y^2}{x + y} = x - y, \text{ Rem., } 2y^2. \\ \frac{x^3 + y^3}{x + y} = x^2 - xy + y^2. \\ \frac{x^4 + y^4}{x + y} = x^3 - x^2y + xy^2 - y^3, \text{ Rem., } 2y^4. \\ \frac{x^5 + y^5}{x + y} = x^4 - x^3y + x^2y^2 - xy^3 + y^4. \end{cases}$$

Observe that the sum of the same powers of two numbers is divisible by the sum of the numbers only when the powers are odd.

111. Hence, when n is a positive integer.

PRINCIPLES. — 1. $x^n - y^n$ is always divisible by $x - y$.

2. $x^n - y^n$ is divisible by $x + y$ only when n is even.

3. $x^n + y^n$ is never divisible by $x - y$.

4. $x^n + y^n$ is divisible by $x + y$ only when n is odd.

112. From § 110, the following law of signs may be readily inferred:

When $x - y$ is the divisor, the signs in the quotient are plus.

When $x + y$ is the divisor, the signs in the quotient are alternately plus and minus.

113. The following law of exponents may also be inferred:

When $x^n \pm y^n$ is divided by $x \pm y$, the quotient is homogeneous, the exponent of x decreasing and that of y increasing by 1 in each successive term.

114. Proofs of preceding principles.

PRINCIPLE 1

	$x^n - y^n$	$x - y$
	$x^n - x^{n-1}y$	$x^{n-1} + x^{n-2}y + \dots$
1st Rem.,	$x^{n-1}y - y^n$	
	$x^{n-1}y - x^{n-2}y^2$	
2d Rem.,	$x^{n-2}y^2 - y^n$	
.	
nth Rem.,	$x^0y^n - y^n$	
or	$x^0y^n - y^n = y^n - y^n = 0$	

By dividing until several remainders are obtained, it is found that the first term of the first remainder is $x^{n-1}y$; of the second, $x^{n-2}y^2$; of the third, $x^{n-3}y^3$; of the fourth, $x^{n-4}y^4$; and consequently of the n th, $x^{n-n}y^n$. But $x^{n-n} = x^0$, which, § 103, equals 1. Therefore, the first term of the n th remainder reduces to y^n .

Since the second term of the n th remainder is $-y^n$, the entire n th remainder is $y^n - y^n$, or 0; that is, there is no remainder, and the division is exact.

Therefore, $x^n - y^n$ is divisible by $x - y$ when x and y represent any two numbers and n is any positive integer.

PRINCIPLE 2

	$\frac{x^n - y^n}{x^n + x^{n-1}y}$	$\frac{x + y}{x^{n-1} - x^{n-2}y + x^{n-3}y^2 - x^{n-4}y^3 + \dots}$
1st Rem.,	$\begin{array}{r} -x^{n-1}y - y^n \\ -x^{n-1}y - x^{n-2}y^2 \end{array}$	
2d Rem.,	$\begin{array}{r} x^{n-2}y^2 - y^n \\ x^{n-2}y^2 + x^{n-3}y^3 \end{array}$	
3d Rem.,	$\begin{array}{r} -x^{n-3}y^3 - y^n \\ -x^{n-3}y^3 - x^{n-4}y^4 \end{array}$	
4th Rem.,	$x^{n-4}y^4 - y^n$	

SUGGESTION.—Since the second term of each remainder is negative, no remainder can reduce to 0 unless its first term is positive. Show for what values of n such remainders reduce to 0 when n is a positive integer.

Prove Principle 3.

Prove Principle 4.

EXAMPLES

Write by inspection the quotient of

1. $\frac{a^2 - b^2}{a - b}$

6. $\frac{m^8 - n^8}{m - n}$

11. $\frac{x^3 - 9}{x + 3}$

2. $\frac{m^5 + n^5}{m + n}$

7. $\frac{n^3 - 1}{n - 1}$

12. $\frac{a^3 - 8}{a - 2}$

3. $\frac{x^6 - y^6}{x + y}$

8. $\frac{c^5 - d^5}{c - d}$

13. $\frac{x^5 - 32}{x - 2}$

4. $\frac{r^2 - s^2}{r - s}$

9. $\frac{1 + a^5}{1 + a}$

14. $\frac{c^2 + 27}{c + 3}$

5. $\frac{a^9 + b^9}{a + b}$

10. $\frac{x^6 - 1}{x + 1}$

15. $\frac{a^7 + 128}{a + 2}$

16. Find five exact binomial divisors of $a^6 - x^6$.

SOLUTION

$a^6 - x^6$ is divisible by $a - x$ (Prin. 1).

$a^6 - x^6$ is divisible by $a + x$ (Prin. 2).

Since $a^6 - x^6 = (a^2)^3 - (x^2)^3$, $a^6 - x^6$ may be regarded as the difference of two odd powers, and is, therefore, divisible by $a^2 - x^2$ (Prin. 1).

Since $a^6 - x^6 = (a^3)^2 - (x^3)^2$, $a^6 - x^6$ may be regarded as the difference of two squares, and is, therefore, divisible by $a^3 - x^3$ (Prin. 1).

Since $a^6 - x^6 = (a^3)^2 - (x^3)^2$, $a^6 - x^6$ may be regarded as the difference of two squares, and is, therefore, divisible by $a^3 + x^3$ (Prin. 2).

Therefore, the exact binomial divisors of $a^6 - x^6$ are $a - x$, $a + x$, $a^2 - x^2$, $a^3 - x^3$, and $a^3 + x^3$.

17. Find an exact binomial divisor of $a^6 + x^6$.

SOLUTION

Since $a^6 + x^6 = (a^2)^3 + (x^2)^3$, $a^6 + x^6$ may be regarded as the sum of the cubes of a^2 and x^2 , and is, therefore, divisible by $a^2 + x^2$ (Prin. 4).

Find exact binomial divisors:

18. $a^2 - m^2$.

24. $x^7 + a^7$.

30. $a^4 - b^4$, four.

19. $a^3 - m^3$.

25. $a^{10} + b^{10}$.

31. $a^6 - 1$, five.

20. $b^3 + x^3$.

26. $a^{10} + b^5$.

32. $a^8 - b^8$, six.

21. $x^5 - a^5$.

27. $a^{12} + b^{12}$.

33. $a^{10} - b^{10}$, five.

22. $c^5 + n^5$.

28. $a^3 - 27$.

34. $a^{16} - b^{16}$, eight.

23. $a^6 + b^6$.

29. $a^6 - 27$.

35. $a^{12} - b^{12}$, nine.

EQUATIONS AND PROBLEMS

115. 1. Find the value of x in the equation $bx - b^2 = cx - c^2$.

SOLUTION

$$bx - b^2 = cx - c^2.$$

Transposing,

$$bx - cx = b^2 - c^2.$$

Collecting coefficients of x ,

$$(b - c)x = b^2 - c^2.$$

Dividing by $b - c$,

$$x = \frac{b^2 - c^2}{b - c} = b + c.$$

2. Find the value of x in the equation $x - a^3 = 2 - ax$.

SOLUTION

$$x - a^3 = 2 - ax.$$

Transposing,

$$ax + x = a^3 + 2.$$

Collecting coefficients of x , $(a + 1)x = a^3 + 2$.

Dividing by $a + 1$,

$$x = \frac{a^3 + 2}{a + 1} = a^2 - a + 1 + \frac{1}{a + 1}$$

Solve the following equations:

3. $7a - 10 = a^2 - ax + 5x$.
6. $cx - c^3 - d^3 + dx = 0$.
4. $x - 1 - c = cx - c^3 - c^4$.
7. $a^2 - ax - 2ab + bx + b^2 = 0$.
5. $2m^3 - mx + nx - 2n^3 = 0$.
8. $2n^2 + 5n + x = n^3 - nx - 2$.
9. $n^2x - 3m^2n^3 + nx + 3m^2 + x = 0$.
10. $a^2x - a^3 + 2a^2 + 5x - 5a + 10 = 0$.
11. $3ab - a^3 - 2bx = 2b^2 - ax$.
12. $9a^2 + 4mx = -(3ax - 16m^2)$.
13. $cy - c^4 - 2c^3 - 2c^2 = 2c - y + 1$.
14. $ay - 2by + 3cy = a - 2b + 3c$.
15. $z + 6n^4 - 4n^3 = 1 - 3nz + 2n - n^2$.
16. $x - 3b^2 - 192b^2c^3 - 4cx + 16c^2x = 0$.
17. $8b^3 - 18b^2 - 57b - 2bx + 7x + 77 = 0$.

Solve the following problems:

18. A drover, who had 5 times as many sheep as oxen and $\frac{1}{2}$ as many oxen as horses, sold all for \$2300, — the horses at \$35 a head, the oxen at \$25 a head, and the sheep at \$4 a head. What was the number of each?

19. A man paid yearly a certain amount of money for taxes and twice that amount for improvements, and received for rent 3 times as much as he paid out for improvements. If his net gain per year was \$300, what were his taxes per year?

20. A owed B a certain sum of money and C twice as much. D owed A 3 times as much as A owed B, and E owed A 5 times the sum A owed B. A found that if he could settle with them all he would have \$5000. How much did he owe B and C?

21. After taking 3 times a number from 11 times the number and adding to the remainder 7 times the number, the result was 12 less than 117. What was the number?

22. A merchant failed in business, owing A 3 times as much as B, C twice as much as A, and D as much as A and B. If the entire debt to A, B, C, and D was \$28,000, how much did he owe each?

23. At a certain election there were three candidates for the office of mayor. A received half as many votes as B and 4 times as many as C. If the total vote lacked 25 votes of being 2300, how many votes did each receive?

24. Three boys together had 140 marbles. If the second boy had twice as many as the first and half as many as the third, how many had each?

25. In a certain school of 600 students there were twice as many Sophomores and 3 times as many Freshmen as Juniors, and 40 more Seniors than Juniors. How many students were there in each class?

26. Divide 25 into three parts such that the first is one third of the second and 5 greater than the third.

27. A, B, and C divided \$40 so that for every \$2 A received, B and C each received \$3. What was the share of each?

28. Divide \$2200 among A, B, and C, so that B shall have twice as much as A and \$200 less than C.

29. Divide \$351 among three persons so that for every dime the first receives the second shall receive 25 cents and the third a dollar.

30. A man gave equal amounts of money to a school and to a library, and $\frac{1}{4}$ the same amount to a hospital. If to all he gave \$28,000, what sum did he give each?

31. When wheat was worth 85 cents a bushel, oats 35 cents a bushel, and corn 60 cents a bushel, a man bought a quantity of wheat, oats, and corn for \$67. If he bought twice as many bushels of oats as of wheat, and also three times as many bushels of corn as of wheat, how many bushels of each did he buy?

REVIEW

116. Simplify:

$$1. x^3 + 2a\sqrt{xy} - 3mn + 4mn - 4x^3 - 5a\sqrt{xy} + 3x^3 + 4a\sqrt{xy} - 2mn.$$

$$2. 5x^3 + 3x^2y + 4xy^2 - y^3 - \sqrt{x} + 6y^3 + xy^2 - \sqrt{y} - 5x^2y - 7x^3 - 5xy^2 + x^3 + 2\sqrt{x} + x^2y - 6y^3 + \sqrt{y} - \sqrt{x} - 2x^2y + 3xy^2 + x^3.$$

$$3. (\frac{5}{2}a - 3bc + \frac{1}{2}c - 7b) - (\frac{3}{2}a + \frac{1}{2}bc + \frac{1}{2}c + 3b).$$

$$4. (a^2x^2 - 4ay + 4bc + ax) - (b^2x^2 - 4by - ax + bc).$$

$$5. x^5 - (x^5 - 5x^4y + 10x^3y^2 - 10x^2y^3 + 5xy^4 - y^5).$$

$$6. \frac{1}{2}a - \frac{5}{8}x - (\frac{3}{4}a - \frac{1}{2}x) - (3b - \frac{1}{4}x - \frac{2}{3}a) + \frac{1}{8}a.$$

$$7. (a^3 + 3a^2y + 3ay^2)(a^3 - 2ay + y^2).$$

$$8. (x^{2n} + 2x^ny^n + y^{2n})(x^{2n} - 2x^ny^n + y^{2n}).$$

$$9. (\frac{1}{4}x^2 + \frac{1}{3}xy + \frac{1}{6}y^2)(\frac{1}{4}x^2 - \frac{1}{3}xy + \frac{1}{6}y^2).$$

$$10. (.2a^2 - .8a + 1.6)(.1a^2 + .4a + .8).$$

Expand:

$$11. (y - 3)(y + 4).$$

$$21. (x + m)(x + m).$$

$$12. (y + 7)(y - 8).$$

$$22. (x^3 + x^2)(x + 1).$$

$$13. (y - 1)(y + 2).$$

$$23. (x - 1)(1 + x).$$

$$14. (y - 5)(y - 9).$$

$$24. (x + 3y)(x - 2y).$$

$$15. (y + 8)(y - 4).$$

$$25. (a^n + b^m)(a^n - b^m).$$

$$16. (m - x)(m + x).$$

$$26. (a^m + b^n)(a^m - b^n).$$

$$17. (m + a)(m - b).$$

$$27. (a + b + c)(a + b - c).$$

$$18. (x - m)(x + n).$$

$$28. (x + y + z)(x - y + z).$$

$$19. (x^2 + x)(x + 2).$$

$$29. (r + s - t)(s - t - r).$$

$$20. (x^2 + 4)(x^2 - 3).$$

$$30. (m + n - p)(m - n + p).$$

$$31. (a - b)(a + b)(a^2 + b^2).$$

$$32. (1 - x)(1 + x)(1 + x^2)(1 + x^4).$$

$$33. (1 - x)(1 + x)(1 - x)(1 + x).$$

$$34. (m + n)(m + n)(m - n)(m - n).$$

35. $(a^4 + a^3 + a^2 + a + 1)(a - 1).$

36. $(x^3 + x^2y + xy^2 + y^3)(x - y).$

37. $(x^5 - x^4 + x^3 - x^2 + x - 1)(x + 1).$

38. $(a^5 + 2a^4 + 4a^3 + 8a^2 + 16a + 32)(a - 2).$

Square

39. $2x - 3y.$

42. $100 - 5.$

45. $a + b + c + d.$

40. $x^4 - ax^2.$

43. $n^2 - m^2.$

46. $2a - 3b - 4c.$

41. $50 - 1.$

44. $7x - b^2y.$

47. $x^{n-1} - y - a^2.$

Expand:

48. $(5a - 4y)(5a - 3y).$

51. $(2a^2x - 5b^2y)(4a^2x - 3b^2y).$

49. $(6x - 4y)(3x + 5y).$

52. $(6amn + 5p)(6amn - 3p).$

50. $(3x + ay)(3x + by).$

53. $(3a^{n+1} - 2b^{n-1})(2a^{n+1} - 3b^{n-1}).$

54. $(x + y)(x - y)(x^2 + y^2)(x^4 + y^4)(x^8 + y^8).$

55. $(m^8 + 1)(m^4 + 1)(m^2 + 1)(m + 1)(m - 1).$

56. $(16x^4 + 1)(4x^2 + 1)(2x + 1)(2x - 1).$

Divide:

57. $x^5 - 2x^4 + 2x^3 + 12x^2 - x - 8$ by $x + 1.$

58. $x^4 - 4x^3 + 5x^2 - 4x + 1$ by $x^2 - 3x + 1.$

59. $x^5 - 45x^4 + 45x^3 - 18x^2 + 9x - 1$ by $x^3 - 4x^2 + 3x - 1.$

60. $a^7 - 12a^2 - a + 12$ by $a^3 - 2a^2 + 4a - 3.$

61. $b^5 - 10b^2 - 5b + 4$ by $b^3 - 2b^2 + 3b - 1.$

62. $m^{10} - 6m^3 + 5m - 2$ by $m^4 + 2m^3 - 3m - 2.$

63. $a^7 - 160a^4 + 127a^3 - 100a^2 - 20a + 16$ by $a^3 - 6a^2 + 5a - 4.$

64. $b^{10} + 29b^4 - 170b^3 - 61b^2 + 210b - 22$ by $b^4 + 2b^2 - 5b - 11.$

65. $6a^7 + \frac{3}{8}a^2y^5 - \frac{3}{8}ay^6 + \frac{3}{8}y^7$ by $a^3 + \frac{1}{2}a^2y - \frac{1}{4}ay^2 + \frac{1}{8}y^3.$

66. $a^2c - ab^2 + acd - ad^2 - abc + b^3 - bcd + bd^2 - ac^2 + cb^2 - c^2d + cd^2$
by $ac - b^2 + cd - d^2.$

67. Divide 1 by $1 - x$ to six terms.

Simplify:

$$68. 4a^2 + 3b^2 - (2a^2 - 3b^2) - (7b^2 - 2a^2) - (-a^2 - b^2).$$

$$69. a^2 - (b^2 - c^2) - (b^2 + c^2 - a^2) + (c^2 - \overline{b^2 - a^2}) - (a^2 - b^2 - c^2).$$

$$70. x^2 - (2xy - y^2) - (x^2 + xy - y^2) - x^2 - \overline{2xy - y^2} + 5y^2.$$

$$71. m + \{2m - [n + 3p - (4p - 3n) - 5n + 2m] - 7p\}.$$

$$72. x - 3y + \{2z - (5y - \overline{4x - 7z}) - (x - y) - z\} - x.$$

$$73. 1 - \{1 - [x^2 - 3 - (2x^2 - 4) + 3x^2 + 1] - (x^2 - 4)\} - 1.$$

$$74. a - (2b + 5a) - 2b - [3a - (6b - 5a) - a] + 12a.$$

$$75. x - \{m - [x + (3m - 2x) + 5x - (2x + m)] - 2x + m\}.$$

$$76. x - \{5x - [6x - (7x - \overline{8x - 9x}) - 10x] + 11x\} + 9x.$$

$$77. 3 - \{c - [5 - (2c - \overline{7 - 3c} - 11) + 5c] - 6c - 20\}.$$

$$78. x + \{3y + [4x - (2y - 7x) - 3y] - (10y + 4x) + 8y\}.$$

$$79. 1 - \{-[-(1 - x) - 1] - 1\} - \{x - (5 - 3x) - 7 + x\}.$$

Collect the coefficients of x , y , and z :

$$80. ax + ay + az - bx - by - bz.$$

$$81. ax - 2y + cz + by - 12x + 4z.$$

$$82. 3mx - nx + by - y + 3cz - 4z.$$

$$83. py - y - 4z + bz - x + mx - nx - z.$$

$$84. cx - by - 3az + x - y - 4z + z - y.$$

$$85. 16ny - 16mx + ax + by + cx - 2y.$$

$$86. mx + ny + az + 2ax - 2my + 2nz.$$

$$87. x - y - az + 8mx + aby - x^2 + y^2 + z.$$

$$88. a^2x + b^2y - 2ax - 2cz + c^2z + x + y + z.$$

$$89. m^2x - n^2y + m^2y - n^2x - 2mnx - 2mny - n^2z + z.$$

$$90. 4(ax - by + cz) - 2(bx - ay - cz) - 2(x - y + z).$$

FACTORING

117. The numbers that multiplied together produce a given number are called its **Factors**.

The factors of $12a$ are 2, 2, 3, and a ; or 4, 3, and a ; or 2, 6, and a ; or 12 and a ; or 2 and $6a$; or $4a$ and 3, etc.

118. A number that has no integral factors except itself and 1 is called a **Prime Number**.

119. A number that has integral factors besides itself and 1 is called a **Composite Number**.

120. A factor that is a prime number is called a **Prime Factor**.

121. The process of separating a number into its factors is called **Factoring**.

122. To factor a monomial.

The factors of the numerical coefficient are found as in arithmetic, but the factors of the literal part are evident.

Thus, in a^3 , the three factors are as evident as if they were written $a \times a \times a$.

It is seldom necessary to resolve a monomial into its simplest factors, but the following problem often occurs:

Given one factor of a monomial, to find the other.

RULE. — *Divide the monomial by the given factor.*

1. In each of the following, if xy is one factor, find the other: $6x^3y$, $15x^4y^2$, $2x^3y^3$, $a^2x^2b^2y^2$, $-mnxy$, $-xy$.

2. In each of the following, if abc is one factor, find the other: a^2bc , ab^2c , abc^2 , $-a^2b^2c^2$, $-a^2bc$, $-\frac{1}{3}abc$.

3. Find two equal positive factors of x^2 ; of $9a^2x^2$; of $64m^4$.

4. Find two equal negative factors of $25x^2$; of $16a^2$; of $9a^4$.

123. To factor a polynomial whose terms have a common factor.

EXAMPLES

1. What are the factors of $3a^2xy - 6ax^2y + 9axy^2$?

PROCESS

$$3a^2xy - 6ax^2y + 9axy^2 \\ = 3axy(a - 2x + 3y)$$

EXPLANATION. — By examining the terms of the polynomial, it is seen that $3axy$ is a factor of every term. Dividing by this common factor, the other factor is found.

Hence, the factors are $3axy$, the monomial factor, and $(a - 2x + 3y)$, the polynomial factor, since, by the Distributive Law for multiplication,

$$3axy(a - 2x + 3y) = 3a^2xy - 6ax^2y + 9axy^2.$$

Find the factors of each of the following polynomials:

- | | |
|---------------------------------------|--|
| 2. $5x^3 - 5x^2.$ | 14. $ac - bc - cy - abc.$ |
| 3. $8x^2 + 2x^4.$ | 15. $3x^3y^3 - 3x^2y^2 + 12xy.$ |
| 4. $3x^3 - 6x^2y.$ | 16. $16a^2b^3c^4 - 24a^3b^2c^3 + 32a^4b^4c^2.$ |
| 5. $4a^2 - 6ab.$ | 17. $60m^2n^3r^2 - 45m^3n^2r^3 + 90m^4n^3r^2.$ |
| 6. $5m^2 - 3mn.$ | 18. $12a^2by - 18ab^2y^2 + 24a^3b^2y^2.$ |
| 7. $3x^3y^2 - 3x^2y^3.$ | 19. $14a^2mn^2 - 21a^3m^2n^3 - 49a^4mn^3.$ |
| 8. $5m^4n - 10m^3n^2.$ | 20. $12x^2y^2z^2 - 16x^3y^2z^2 - 20x^2y^3z^2.$ |
| 9. $4a^3b - 6a^2b^2.$ | 21. $25c^2dx^3 + 35c^3d^2x^4 - 55c^2d^2x^5.$ |
| 10. $5x^4 - 10x^3 - 5x^2.$ | 22. $51xy^2z^3 - 68x^2y^2z^3 + 85x^4y^2z^4.$ |
| 11. $3a^4 - 2a^3b + a^2b^2.$ | 23. $52a^2b^3c^4 - 65a^3b^2c^3 - 91a^2b^2c^2.$ |
| 12. $x^{12} + x^{11} + x^{10} - x^9.$ | 24. $44a^4x^3y^2 + 66a^3x^4y^3 + 88a^2x^5y^4.$ |
| 13. $3m^5 - 12m^3n^2 + 6mn^4.$ | 25. $84x^4y^2 - 36x^3y^3 + 60x^2y^4 - 48xy^5.$ |

124. To factor a polynomial whose terms may be grouped to show a common polynomial factor.

EXAMPLES

1. Factor $ax + ay + bx + by.$

SOLUTION

$$ax + ay + bx + by = a(x + y) + b(x + y) \\ = (a + b)(x + y).$$

2. Factor $ax - ay - bx + by$.

SOLUTION

$$\begin{aligned}
 & ax - ay - bx + by \\
 &= a(x - y) - b(x - y) \\
 &= (a - b)(x - y).
 \end{aligned}$$

Observe that, when the first two terms are factored, $(x - y)$ is found to be the binomial factor. Since $(x - y)$ is to be a factor of the other two terms, the monomial factor is $-b$, not $+b$.

3. Factor $cx + y - dy + cy - dx + x$.

SOLUTION

$$\begin{aligned}
 & cx + y - dy + cy - dx + x \\
 \text{Arranging terms,} \quad &= cx - dx + x + cy - dy + y \\
 &= (c - d + 1)x + (c - d + 1)y \\
 &= (c - d + 1)(x + y).
 \end{aligned}$$

Factor the following:

- | | |
|-------------------------------------|--------------------------------------|
| 4. $am - an + mx - nx$. | 16. $x^3 + x^2 + x + 1$. |
| 5. $bc - bd + cx - dx$. | 17. $y^3 + y^2 - 3y - 3$. |
| 6. $pq - px - rq + rx$. | 18. $x^5 + x^3 + x^2y + y$. |
| 7. $ay - by - ab + b^2$. | 19. $2 - 2n - n^2 + n^3$. |
| 8. $x^2 - xy - 5x + 5y$. | 20. $x^2 - x - a + ax$. |
| 9. $b^2 - bc + ab - ac$. | 21. $3x^3 - 15x + 10y - 2x^2y$. |
| 10. $x^2 + xy - ax - ay$. | 22. $12a^3 - 8ab - 3a^4 + 2a^2b$. |
| 11. $c^2 - 4c + ac - 4a$. | 23. $3m^2n - 9mn^2 + am - 3an$. |
| 12. $2x - y + 4x^2 - 2xy$. | 24. $15ab^2 - 9b^2c - 35ab + 21bc$. |
| 13. $1 - m + n - mn$. | 25. $16ax + 12ay - 8bx - 6by$. |
| 14. $2p + q + 6p^2 + 3pq$. | 26. $ax^2 - ax - axy + ay + x - 1$. |
| 15. $ar - rs - ab + bs$. | 27. $xy + x - 3y^2 - 3y - 4y - 4$. |
| 28. $ax - a - bx + b - cx + c$. | |
| 29. $mx - nx - x - my + ny + y$. | |
| 30. $a^2 - a - ab + b - 2ac + 2c$. | |

31. $mp^2 - np^2 + mq - nq + m - n$.
32. $ax^2 - bx^2 - ax + bx + a - b$.
33. $2mx^2 - nx^2 + n + 2nx - 4mx - 2m$.
34. $bx^2 - b - xy - y + yx^2 - bx$.
35. $a^2x - a^2y - ay - y + x + ax$.
36. $2 - 3b + 3ab - 2a + 4a^2 - 6a^2b$.
37. $m^2 + mn + mn + n^2 + m + n$.

125. To factor a trinomial that is a perfect square.

1. Since $(a+b)(a+b) = a^2 + 2ab + b^2$, what are the factors of $a^2 + 2ab + b^2$? How are they obtained from $a^2 + 2ab + b^2$?
2. Since $(a-b)(a-b) = a^2 - 2ab + b^2$, what are the factors of $a^2 - 2ab + b^2$? How are they obtained from $a^2 - 2ab + b^2$?
3. What determines the sign that connects the terms of each factor?

126. One of the two equal factors of a number is called its **Square Root**.

127. In a trinomial that is a perfect square one term is equal to twice the product of the square roots of the other two terms.

$25x^2 - 20xy + 4y^2$ is a perfect square, for twice the product of the square root of $25x^2$ and the square root of $4y^2$ is $20xy$.

RULE. — *Connect the square roots of the terms that are squares with the sign of the other term, and indicate that the result is to be taken twice as a factor.*

From any expression that is to be factored, the *monomial* factors should usually first be removed.

Thus, $2a^3 - 4a^2 + 2a = 2a(a^2 - 2a + 1) = 2a(a-1)^2$.

EXAMPLES

Factor the following:

- | | |
|------------------------|-----------------------|
| 1. $x^2 + 2xy + y^2$. | 5. $x^2 + 6x + 9$. |
| 2. $p^2 - 2pq + q^2$. | 6. $m^2 - 8m + 16$. |
| 3. $c^2 + 2cd + d^2$. | 7. $x^2 - 2x + 1$. |
| 4. $m^2 - 2mn + n^2$. | 8. $a^2 - 16a + 64$. |

- | | |
|--------------------------|---|
| 9. $x^2 + 4x + 4.$ | 20. $9 + 42b^3 + 49b^6.$ |
| 10. $4 - 4a + a^2.$ | 21. $9m^3 - 6m^4 + 1.$ |
| 11. $4a - 4a^2 + a^3.$ | 22. $4x^2y^2 - 20xy + 25.$ |
| 12. $3x^2 + 6xy + 3y^2.$ | 23. $4x^2 + 12xyz + 9y^2z^2.$ |
| 13. $2m^2 - 4mn + 2n^2.$ | 24. $9a^2m^2 - 6am + 1.$ |
| 14. $5x^2 + 30x + 45.$ | 25. $2x + 20a^2x + 50a^4x.$ |
| 15. $10x^2 - 20x + 10.$ | 26. $18a^2b + 60ab^2 + 50b^3.$ |
| 16. $16p^2 - 24p + 9.$ | 27. $a^2x^6 - 2ax^3by^3 + b^2y^6.$ |
| 17. $9x^2 - 42x + 49.$ | 28. $25a^{2m} - 60a^mb^n + 36b^{2n}.$ |
| 18. $1 + 4b + 4b^2.$ | 29. $x^{2n} - 2x^ny^n z^n + y^{2n}z^{2n}.$ |
| 19. $1 - 6a^3 + 9a^6.$ | 30. $81a^2b^2c^2 + 18ab^2c^2d + b^2c^2d^2.$ |

When either or both of the squares are polynomials, the expression may be factored in a similar manner.

31. Factor $x^2 + 6x(x-y) + 9(x-y)^2.$

SOLUTION

$$\begin{aligned}
 & x^2 + 6x(x-y) + 9(x-y)^2 \\
 &= [x + 3(x-y)][x + 3(x-y)] \\
 &= (x + 3x - 3y)(x + 3x - 3y) \\
 &= (4x - 3y)(4x - 3y).
 \end{aligned}$$

32. Factor $(a-b)^2 + 2(a-b)(b-c) + (b-c)^2.$

SOLUTION

$$\begin{aligned}
 & (a-b)^2 + 2(a-b)(b-c) + (b-c)^2 \\
 &= [(a-b) + (b-c)][(a-b) + (b-c)] \\
 &= (a-b+b-c)(a-b+b-c) \\
 &= (a-c)(a-c).
 \end{aligned}$$

Factor:

33. $x^2 + 2x(x-y) + (x-y)^2.$
34. $a^2 - 4a(a-1) + 4(a-1)^2.$
35. $c^2 - 6c(a-c) + 9(a-c)^2.$
36. $m^2 + 2m(m-n) + (m-n)^2.$
37. $16 - 24(a-b) + 9(a-b)^2.$
38. $x^2 + 25(y^3-x)^2 + 10x(y^3-x).$
39. $14a(x-y) + (x-y)^2 + 49a^2.$
40. $10m(m-4) + 25m^2 + (m-4)^2.$

41. $(a+b)^2 - 2(a+b)(b+c) + (b+c)^2$.
42. $(a-2x)^2 + 4(a-2x)(2x-b) + 4(2x-b)^2$.
43. $16(a-x)^2 + 32(a-x)(x+b) + 16(x+b)^2$.
44. $(a+3b)^2 - 4(a+3b)(3b-2c) + 4(3b-2c)^2$.
45. $(x^2+x+1)^2 + 2(x+1)(x^2+x+1) + (x+1)^2$.
46. $(a+b+c)^2 + 2(a+b-c)(a+b+c) + (a+b-c)^2$.
47. $(x^3-x^2)^2 + 2(x^3-x^2)(x+1) + (x^2+2x+1)^2$.

128. To factor the difference of two squares.

1. Since $(a+b)(a-b) = a^2 - b^2$, what are the factors of $a^2 - b^2$?
How do these two factors differ?

2. Since $(a^2 + b^2)(a^2 - b^2) = a^4 - b^4$, what are the factors of $a^4 - b^4$? How do these two factors differ?

RULE. — Find the square roots of the two terms, and make their sum one factor and their difference the other.

Sometimes the factors of a number may themselves be factored.

EXAMPLES

1. Factor $b^2 - y^2$.

SOLUTION. $b^2 - y^2 = (b+y)(b-y)$.

2. Factor $x^2 - 1$.

SOLUTION. $x^2 - 1 = (x+1)(x-1)$.

3. Factor $x^4 - 1$.

SOLUTION. $x^4 - 1 = (x^2+1)(x^2-1)$
 $= (x^2+1)(x+1)(x-1)$.

Resolve into their simplest factors:

- | | | |
|------------------|-------------------|-----------------------|
| 4. $x^2 - m^2$. | 9. $25 - c^2$. | 14. $a^{16} - b^8$. |
| 5. $a^2 - y^2$. | 10. $x^4 - 49$. | 15. $4x^2 - 25y^2$. |
| 6. $a^2 - 16$. | 11. $x^4 - 81$. | 16. $9a^2 - 49b^2$. |
| 7. $y^2 - x^2$. | 12. $a^4 - 16$. | 17. $a^2x^2 - 4c^2$. |
| 8. $a^2 - 9$. | 13. $a^4 - b^4$. | 18. $m^4 - 16n^4$. |

- | | | |
|------------------------|-------------------------|----------------------------|
| 19. $25x^2 - 1$. | 25. $400x^2 - 100y^2$. | 31. $5x^3 - 5$. |
| 20. $81m^4 - 1$. | 26. $2a^8 - 2b^8$. | 32. $3a^5 - 3a$. |
| 21. $36a^4 - 25$. | 27. $4m^4 - 4b^4$. | 33. $x^3 - xy^2$. |
| 22. $121b^2 - c^2$. | 28. $3x^4 - 3y^4$. | 34. $5a^5y - 5ay$. |
| 23. $400a^2 - 81y^2$. | 29. $5x^4 - 5y^{10}$. | 35. $x^{2n} - y^{2n}$. |
| 24. $100x^{2n} - 1$. | 30. $8x^{10} - 8y^6$. | 36. $x^{2n+1} - xy^{2n}$. |

When either or both of the squares are polynomials, the expression may be factored in a similar manner.

37. Factor $25a^2 - (3a + 2b)^2$.

SOLUTION

One factor is $5a + (3a + 2b)$, and the other is $5a - (3a + 2b)$.

$$5a + (3a + 2b) = 5a + 3a + 2b = 8a + 2b = 2(4a + b).$$

$$5a - (3a + 2b) = 5a - 3a - 2b = 2a - 2b = 2(a - b).$$

$$\begin{aligned}\therefore 25a^2 - (3a + 2b)^2 &= (5a + 3a + 2b)(5a - 3a - 2b) \\ &= (8a + 2b)(2a - 2b) \\ &= 2(4a + b) \cdot 2(a - b) \\ &= 4(4a + b)(a - b).\end{aligned}$$

Factor

- | | |
|--------------------------|-----------------------------|
| 38. $a^2 - (a + b)^2$. | 42. $9b^2 - (a - x)^2$. |
| 39. $b^2 - (2a + b)^2$. | 43. $9a^2 - (2a - 5)^2$. |
| 40. $a^2 - (b + c)^2$. | 44. $x^4 - (3x^2 - 2y)^2$. |
| 41. $4c^2 - (b + c)^2$. | 45. $49a^2 - (5a - 4b)^2$. |
46. Factor $(3a - 2b)^2 - (2a - 5b)^2$.

SOLUTION

$$\begin{aligned}&(3a - 2b)^2 - (2a - 5b)^2 \\ &= [(3a - 2b) + (2a - 5b)][(3a - 2b) - (2a - 5b)] \\ &= (3a - 2b + 2a - 5b)(3a - 2b - 2a + 5b) \\ &= (5a - 7b)(a + 3b).\end{aligned}$$

Factor:

- | | |
|---------------------------------|---------------------------------|
| 47. $(2a + 3b)^2 - (a + b)^2$. | 49. $(2x + 5)^2 - (5 - 3x)^2$. |
| 48. $(5a - 3b)^2 - (a - b)^2$. | 50. $(a - 2b)^2 - (a - 5)^2$. |

51. $(2x - 3y)^2 - (3y + z)^2$. 54. $(9x + 6y)^2 - (4x - 3y)^2$.
 52. $(5b - 4c)^2 - (3a - 2c)^2$. 55. $(x^3 + x^2)^2 - (2x + 2)^2$.
 53. $(4x - 3y)^2 - (2x - 3a)^2$. 56. $(a + b + c)^2 - (a - b - c)^2$.
 57. Factor $a^2 + 4 - c^2 - 4a$.

SOLUTION

$$\begin{aligned} & a^2 + 4 - c^2 - 4a \\ \text{Arranging terms,} \quad & = a^2 - 4a + 4 - c^2 \\ & = (a - 2)^2 - c^2 \\ & = (a - 2 + c)(a - 2 - c). \end{aligned}$$

58. Factor $a^2 + b^2 - c^2 - 4 - 2ab + 4c$.

SOLUTION

$$\begin{aligned} & a^2 + b^2 - c^2 - 4 - 2ab + 4c \\ \text{Arranging terms,} \quad & = a^2 - 2ab + b^2 - c^2 + 4c - 4 \\ & = (a^2 - 2ab + b^2) - (c^2 - 4c + 4) \\ & = (a - b)^2 - (c - 2)^2 \\ & = (a - b + c - 2)(a - b - c + 2). \end{aligned}$$

Factor:

59. $a^2 - 2ax + x^2 - n^2$. 67. $c^2 - a^2 - b^2 - 2ab$.
 60. $b^2 + 2by + y^2 - n^2$. 68. $b^2 - x^2 - y^2 + 2xy$.
 61. $1 - 4q + 4q^2 - a^2$. 69. $4c^2 - x^2 - y^2 - 2xy$.
 62. $r^2 - 2rx + x^2 - 16t^2$. 70. $9c^2 - x^2 - y^2 + 2xy$.
 63. $9a^2b - 6ab^2 + b^3 - 4bc^2$. 71. $x^3 - a^2x - 4b^2x - 4abx$.
 64. $4a^2c + 12abc + 9b^2c - 4c^3$. 72. $bc^2 - 9a^2b - b^3 - 6ab^2$.
 65. $3x^2y - 12xy^2 + 12y^3 - 3x^2y$. 73. $ab^2 - 4a^3 - 12a^2c - 9ac^2$.
 66. $4an^4 - 16a^2n^2 + 16a^3 - 4an^6$. 74. $27c^2 - 12a^2 + 36ab - 27b^2$.
 75. $a^2 - 2ab + b^2 - c^2 + 2cd - d^2$.
 76. $x^2 - 2xy + y^2 - m^2 + 10m - 25$.
 77. $4x^2 + 9 - 12x + 10mn - m^2 - 25n^2$.
 78. $x^3 - a^3 + y^3 - b^3 + 2xy - 2ab$.

129. To factor a quadratic trinomial.

$$(x + 3)(x + 5) = x^2 + 8x + 15.$$

$$(x - 3)(x - 5) = x^2 - 8x + 15.$$

$$(x + 1)(x + 15) = x^2 + 16x + 15.$$

$$(x - 1)(x + 15) = x^2 + 14x - 15.$$

$$(x - 3)(x + 5) = x^2 + 2x - 15.$$

$$(x + 3)(x - 5) = x^2 - 2x - 15.$$

1. How may the first term of each factor be found from the product?

2. When the last term of the product has the sign +, how do the signs of the last terms of the factors compare? How, when the last term of the product has the sign -?

3. How does the coefficient of the middle term of the product compare with the *algebraic* sum of the last terms of the factors?

130. A trinomial of the form $x^2 + bx + c$, in which x^2 is the square of any number, c the product of two numbers, and b their algebraic sum, b and c being either positive or negative, is called a **Quadratic Trinomial**.

RULE. — *Arrange the trinomial according to the descending powers of one of the letters.*

For the first term of each factor take the square root of the first term of the trinomial; and for the second terms, such numbers that their product is the third term of the trinomial, and their algebraic sum multiplied by the first term of the factor will be equal to the second term.

EXAMPLES

1. Resolve $x^2 - 13x - 48$ into two binomial factors.

SOLUTION. — The first term of each factor is evidently x .

Since the product of the second terms of the two binomial factors is -48 , the second terms must have opposite signs; and since their algebraic sum, -13 , is negative, the negative term must be numerically larger than the positive term.

The two factors of -48 whose sum is negative may be 1 and -48 , 2 and -24 , 3 and -16 , 4 and -12 , or 6 and -8 . Since the algebraic sum of 3 and -16 is -13 , 3 and -16 are the factors of -48 sought.

$$\therefore x^2 - 13x - 48 = (x + 3)(x - 16).$$

2. Factor $m^2 + m - 72$.

SOLUTION. — Since +9 and -8 are the only two factors of -72 whose algebraic sum is +1, the coefficient of the middle term, the required factors are $(m + 9)$ and $(m - 8)$.

$$\therefore m^2 + m - 72 = (m + 9)(m - 8).$$

Separate the following into their simplest factors:

- | | |
|-------------------------|---------------------------------------|
| 3. $x^2 + 7x + 12$. | 18. $x^2 + 5ax + 6a^2$. |
| 4. $y^2 - 7y + 12$. | 19. $x^2 - 6ax + 5a^2$. |
| 5. $p^2 - 8p + 12$. | 20. $y^2 - 4by - 12b^2$. |
| 6. $r^2 + 8r + 12$. | 21. $y^2 - 3ny - 28n^2$. |
| 7. $m^2 + 5m - 14$. | 22. $x^2 - anz - 2a^2n^2$. |
| 8. $a^2 - 2a - 15$. | 23. $x^4 + 19cx^2 + 90c^2$. |
| 9. $b^2 + b - 12$. | 24. $x^6 + 12ax^3 + 20a^2$. |
| 10. $r^2 + r - 30$. | 25. $x^{10} - 11b^2x^5 + 24b^4$. |
| 11. $c^2 - c - 72$. | 26. $5nx^2 - 55nx + 150n$. |
| 12. $c^2 - 5c - 14$. | 27. $3a^2bx^2 - 3a^2bx - 6a^2b$. |
| 13. $x^2 - x - 110$. | 28. $12m^2x^2 - 60m^2bx + 72m^2b^2$. |
| 14. $a^2 + 9a - 52$. | 29. $4ax + 2ax^2 - 48a$. |
| 15. $a^2 + 8a - 128$. | 30. $11a^2x - 55ax + 66x$. |
| 16. $x^2 - 25x + 100$. | 31. $20bx + 10b^2 - 630x^2$. |
| 17. $x^2 + 12x - 85$. | 32. $x^2 + (b - a)x - ab$. |

131. To factor trinomials of the form $ax^2 + bx + c$.

EXAMPLES

1. Separate $3x^2 + 11x + 6$ into two binomial factors.

SOLUTION. — Since $3x^2$ is the product of the first terms of the factors, and 6 is the product of their last terms, and since the only factors, each containing x , that $3x^2$ can have are $3x$ and x , one factor is $3x$ plus a factor of 6, and the other is x plus the other factor of 6. By trial, it is found that 2 and 3 are the factors of 6, and that if 2 is added to $3x$ and 3 to x , the middle term of the product $(3x + 2)(x + 3)$ will be $11x$.

$$\therefore 3x^2 + 11x + 6 = (3x + 2)(x + 3).$$

2. Factor $9x^2 + 30x + 16$.

SOLUTION

$$\begin{aligned}
 &9x^2 + 30x + 16 \\
 &= (3x)^2 + 10(3x) + 16 \\
 \text{Put } m \text{ for } 3x, &= m^2 + 10m + 16 \\
 \S 130, &= (m+2)(m+8) \\
 \text{Put } 3x \text{ for } m, &= (3x+2)(3x+8).
 \end{aligned}$$

SUGGESTION. — When the coefficient of x^2 is a square, and when the square root of the coefficient of x^2 is exactly contained in the coefficient of x , the trinomial may be factored by the method illustrated above.

3. Factor $4x^2 - 5x - 6$.

SOLUTION

$$\begin{aligned}
 4x^2 - 5x - 6 &= (4x^2 - 5x - 6) \times \frac{4}{4} = \frac{16x^2 - 20x - 24}{4} \\
 &= \frac{(4x)^2 - 5(4x) - 24}{4} = \frac{(4x-8)(4x+3)}{4} \\
 &= \frac{4(x-2)(4x+3)}{4} = (x-2)(4x+3).
 \end{aligned}$$

EXPLANATION. — Although the first term is a square, its square root is not exactly contained in the second term.

But if such a trinomial is multiplied by the coefficient of x^2 , the resulting trinomial will be one whose second term exactly contains the square root of its first term.

Multiplying the given trinomial by 4, factoring as in example 2, and dividing the result by 4, the factors of the given trinomial are $(x-2)$ and $(4x+3)$.

4. Factor $24x^2 + 14x - 5$.

SOLUTION

$$\begin{aligned}
 24x^2 + 14x - 5 &= (24x^2 + 14x - 5) \times \frac{6}{6} = \frac{144x^2 + 84x - 30}{6} \\
 &= \frac{(12x)^2 + 7(12x) - 30}{6} = \frac{(12x+10)(12x-3)}{6} \\
 &= \frac{2(6x+5) \cdot 3(4x-1)}{2 \times 3} = (6x+5)(4x-1).
 \end{aligned}$$

SUGGESTION. — When the first term is not a square, it may always be made a square whose square root will be exactly contained in the second term by multiplying the trinomial by the coefficient of x^2 .

After factoring, the result should be divided by the coefficient of x^2 .
Frequently the multiplier can be taken smaller than the coefficient of x^2 ,
as in the above example.

Separate into their simplest factors :

- | | |
|--------------------------|----------------------------|
| 5. $2x^2 + x - 15$. | 17. $9x^4 - 10x^2 - 16$. |
| 6. $9x^2 - 42x + 40$. | 18. $27b^4 - 3b^2 - 14$. |
| 7. $5x^2 + 13x + 6$. | 19. $10x^5 - 2x^3 - 44$. |
| 8. $3x^2 - 17x + 10$. | 20. $2x^2 + 5xy + 2y^2$. |
| 9. $25x^2 + 15x + 2$. | 21. $2x^2 + 3xy - 2y^2$. |
| 10. $16x^2 + 20x - 66$. | 22. $3x^2 - 10xy + 3y^2$. |
| 11. $36x^2 - 48x - 20$. | 23. $6x^2 - 11x - 35$. |
| 12. $9x^2 + 43x - 10$. | 24. $6x^2 - 13x + 6$. |
| 13. $25x^2 + 25x - 24$. | 25. $15x^2 - 14x - 8$. |
| 14. $49x^2 - 42x - 55$. | 26. $15x^3 + 17x - 4$. |
| 15. $16x^2 + 50x - 21$. | 27. $21a^2 - a - 10$. |
| 16. $4x^2 - 4x - 35$. | 28. $18x^2 - 3x - 36$. |

132. To factor the sum of the same odd powers of two numbers.

EXAMPLES

1. Factor $a^7 + b^7$.

SOLUTION

$$\S\S 111-113, a^7 + b^7 = (a + b)(a^6 - a^5b + a^4b^2 - a^3b^3 + a^2b^4 - ab^5 + b^6).$$

2. Factor $m^5 + 32x^5$.

SOLUTION

$$\begin{aligned} m^5 + 32x^5 &= m^5 + (2x)^5 \\ \S\S 111-113, &= (m + 2x)(m^4 - 2m^3x + 4m^2x^2 - 8mx^3 + 16x^4). \end{aligned}$$

3. Factor $x^5 + y^5$.

SOLUTION

$$\begin{aligned} x^5 + y^5 &= (x^2)^2 + (y^2)^2 \\ \S\S 111-113, &= (x^2 + y^2)(x^4 - x^2y^2 + y^4). \end{aligned}$$

Factor the following:

- | | | |
|------------------|------------------|----------------------|
| 4. $m^3 + n^3$. | 7. $x^5 + 1$. | 10. $a^5 + 32$. |
| 5. $a^3 + x^3$. | 8. $r^7 + s^7$. | 11. $p^3 + 27$. |
| 6. $a^5 + b^5$. | 9. $x^9 + y^9$. | 12. $x^{14} + 128$. |

133. To factor the difference of the same odd powers of two numbers.

EXAMPLES

1. Factor $a^7 - b^7$.

SOLUTION

$$\S\S 111-113, a^7 - b^7 = (a - b)(a^6 + a^5b + a^4b^2 + a^3b^3 + a^2b^4 + ab^5 + b^6).$$

2. Factor $m^5 - 32x^5$.

SOLUTION

$$\begin{aligned}\S\S 111-113, m^5 - 32x^5 &= m^5 - (2x)^5 \\ &= (m - 2x)(m^4 + 2m^3x + 4m^2x^2 + 8mx^3 + 16x^4).\end{aligned}$$

3. Factor $a^{10} - b^5$.

SOLUTION

$$\S\S 111-113, a^{10} - b^5 = (a^2)^5 - b^5 = (a^2 - b)(a^8 + a^6b + a^4b^2 + a^2b^3 + b^4).$$

Factor the following:

- | | | |
|------------------|-----------------|----------------------|
| 4. $a^5 - b^5$. | 7. $y^3 - 8$. | 10. $x^5 - y^{10}$. |
| 5. $x^3 - b^3$. | 8. $a^5 - 32$. | 11. $x^6 - y^3$. |
| 6. $y^3 - a^3$. | 9. $a^7 - 1$. | 12. $x^5 - 243$. |

134. To factor the difference of the same even powers of two numbers.

EXAMPLES

1. Factor $a^6 - b^6$.

FIRST SOLUTION

$$\begin{aligned}\S\S 111-113, a^6 - b^6 &= (a - b)(a^5 + a^4b + a^3b^2 + a^2b^3 + ab^4 + b^5) \\ &= (a - b)(a^5 + a^2b^3 + a^4b + ab^4 + a^3b^2 + b^5) \\ &= (a - b)[a^2(a^3 + b^3) + ab(a^3 + b^3) + b^2(a^3 + b^3)] \\ &= (a - b)(a^2 + ab + b^2)(a^3 + b^3) \\ \S 132, &= (a - b)(a^2 + ab + b^2)(a + b)(a^2 - ab + b^2).\end{aligned}$$

SECOND SOLUTION

$$\S 128, \quad a^6 - b^6 = (a^3)^2 - (b^3)^2 = (a^3 + b^3)(a^3 - b^3)$$

$$\S\S 132, 133, \quad = (a + b)(a^2 - ab + b^2)(a - b)(a^2 + ab + b^2).$$

In each of the above solutions $a^6 - b^6$ is first separated into two factors. Thus, $a^6 - b^6$ is divisible by $a - b$; also by $a^3 - b^3$, since $a^6 - b^6 = (a^3)^2 - (b^3)^2$.

When the even powers are regarded as squares, as in the second solution, the process may be regarded as *factoring the difference of two squares*.

Separate the following into their simplest factors:

2. $x^6 - y^6$.

5. $x^4 - 16$.

8. $1 - b^8$.

3. $x^6 - 1$.

6. $x^4 - 81$.

9. $64 - y^6$.

4. $a^8 - b^8$.

7. $a^4 - 625$.

10. $1 - x^8$.

Factor the following by previous principles:

11. $p^2 - q^2$.

17. $r^6 - s^6$.

23. $1 + x^6$.

12. $p^3 + q^3$.

18. $r^6 + s^6$.

24. $x + x^6$.

13. $p^3 - q^3$.

19. $m^7 - n^7$.

25. $32n - n^6$.

14. $r^4 - s^4$.

20. $m^7 + n^7$.

26. $m^3 - x^6$.

15. $r^5 - s^5$.

21. $m^8 - n^8$.

27. $b^4 - a^2b^4$.

16. $r^5 + s^5$.

22. $a^8 - 1$.

28. $125 + a^3$.

135. To factor by the Factor Law.

1. If $5x = 0$, what must be the value of x ?

2. If $5(x - 3) = 0$, which factor reduces the first member to zero; that is, what *value of* x reduces it to zero?

3. Since the expression $5(x - 3)$ reduces to zero when $x - 3$ reduces to zero, and $x - 3$ is reduced to zero by substituting 3 for x , what value substituted for x will reduce to zero *every* expression of which $x - 3$ is a factor?

4. How, then, may it be discovered whether $x - 3$ is a factor of a given expression containing x ?

5. What value of x will reduce the expression $x^3 - 8$ to zero? What factor, then, has $x^3 - 8$?

6. What value of x will reduce the expression $x^5 - 8x + 7$ to zero? What factor, then, has $x^5 - 8x + 7$?

7. When does a rational integral expression containing x have the factor $x - 1$? $x - 2$? $x - 3$? $x - a$?

136. Factor Law. — *If a rational integral expression containing x reduces to zero when a is substituted for x , it is exactly divisible by $x - a$.*

DEMONSTRATION. — Let D represent any rational integral expression containing x , and let D reduce to zero when a is substituted for x .

It is to be proved that D is exactly divisible by $x - a$.

Suppose that the dividend D is divided by $x - a$ until the remainder does not contain x . Denote the remainder by R and the quotient by Q .

$$\text{Then,} \quad D = Q(x - a) + R. \quad (1)$$

But, since, by supposition, D reduces to zero when $x = a$; that is, when $x - a = 0$, equation (1) becomes

$$0 = 0 + R;$$

whence,

$$R = 0.$$

That is, the remainder is zero, and the division is exact.

NOTE. a is the known number substituted for x , and it may be either positive or negative.

EXAMPLES

1. Factor $x^3 - 8x^2 + 17x - 10$ by the factor law.

SOLUTION

$$\begin{aligned} & x^3 - 8x^2 + 17x - 10 \\ &= (x - 1)(x^2 - 7x + 10) \\ &= (x - 1)(x - 2)(x - 5). \end{aligned}$$

Since, when 1 is substituted for x , $x^3 - 8x^2 + 17x - 10$ reduces to 0, $x - 1$ is a factor of the expression. Dividing by $x - 1$, the other factor is $x^2 - 7x + 10$, whose factors, § 130, are $(x - 2)$ and $(x - 5)$.

2. Factor $x^4 + x^3 - 16x^2 - 4x + 48$ by the factor law.

SOLUTION

$$\begin{aligned} & x^4 + x^3 - 16x^2 - 4x + 48 \\ &= (x - 2)(x^3 + 3x^2 - 10x - 24) \\ &= (x - 2)(x + 2)(x^2 + x - 12) \\ &= (x - 2)(x + 2)(x - 3)(x + 4). \end{aligned}$$

Since, when 2 is substituted for x , the expression reduces to 0, $x - 2$ is a factor of the expression. Dividing by $x - 2$, the other factor is $x^3 + 3x^2 - 10x - 24$. Since, when -2 is substituted for x , $x^3 + 3x^2 - 10x - 24$ reduces to 0, $x + 2$ is a factor of $x^3 + 3x^2 - 10x - 24$.

Dividing by $x + 2$, the other factor is $x^2 + x - 12$, a quadratic trinomial whose factors, § 130, are $(x - 3)$ and $(x + 4)$. Hence, the factors of the given expression are $(x - 2)$, $(x + 2)$, $(x - 3)$, and $(x + 4)$.

Factor the following polynomials by the factor law :

3. $x^2 - 31x + 30$.
4. $4x^2 - 7x + 3$.
5. $26x^2 - 10x - 16$.
6. $48x^2 - 31x - 17$.
7. $36x^2 - 61x + 25$.
8. $x^3 - 9x^2 + 23x - 15$.
9. $x^3 - 13x^2 + 47x - 35$.
10. $x^3 - 14x^2 + 35x - 22$.
11. $x^3 - 4x^2 - 7x + 10$.
12. $x^3 - 6x^2 - 9x + 14$.
13. $x^3 - 12x^2 + 41x - 30$.
14. $x^3 - 11x^2 + 31x - 21$.
15. $x^3 - 10x^2 + 29x - 20$.
16. $x^3 - 16x^2 + 71x - 56$.
17. $x^3 - 57x + 56$.
18. $x^3 - 21x + 20$.
19. $x^3 - 31x - 30$.
20. $x^3 - 13x + 12$.
21. $x^3 - 7x + 6$.
22. $x^3 - 19x + 30$.
23. $x^3 - 67x - 126$.
24. $x^3 - 39x - 70$.
25. $a^3 + 4a^2 - 11a - 30$.
26. $a^3 + 9a^2 + 26a + 24$.
27. $m^3 - 6m^2 - m - 30$.
28. $b^3 - 5b^2 - 29b + 105$.
29. $a^3 + 10a^2 - 17a - 66$.
30. $m^3 + 7m^2 + 2m - 40$.
31. $b^3 + 16b^2 + 73b + 90$.
32. $n^3 + 12n^2 + 41n + 42$.
33. $x^4 - 15x^2 + 10x + 24$.
34. $x^4 - 25x^2 + 60x - 36$.
35. $x^4 + 13x^2 - 54x + 40$.
36. $x^4 + 22x^2 + 27x - 50$.
37. $x^4 - 9x^2y^2 - 4xy^3 + 12y^4$.
38. $x^4 - 9x^2y^2 + 12xy^3 - 4y^4$.
39. $x^4 - x^3 - 7x^2 + x + 6$.
40. $x^4 - 9x^3 + 21x^2 + x - 30$.
41. $x^4 + 8x^3 + 14x^2 - 8x - 15$.
42. $x^5 - 4x^4 + 19x^3 - 28x^2 + 12x$.
43. $x^5 - 18x^3 + 30x^2 - 19x + 30$.
44. $x^5 - 10x^4 + 40x^3 - 80x^2 + 80x - 32$.

SPECIAL APPLICATIONS AND DEVICES

EXAMPLES

137. 1. Factor

$$a^2 + b^2 + c^2 + d^2 + 2ab - 2ac + 2ad - 2bc + 2bd - 2cd.$$

SOLUTION. — Since the polynomial consists of the squares of four numbers together with twice the product of each of them by each succeeding number, the polynomial is the square of the sum of four numbers, § 95, and may be separated into two equal factors containing a , b , c , and d with proper signs.

Since $-2ac$, $-2bc$, and $-2cd$, the products that contain c , are negative, while $2ab$, $2ad$, and $2bd$, the products that do not contain c , are positive, it is evident that the sign of c is the opposite of that of a , b , and d .

Therefore, the factors are either

$$(a + b - c + d)(a + b - c + d),$$

or

$$(-a - b + c - d)(-a - b + c - d).$$

Factor the following:

$$2. 9x^2 + 4y^2 + 25z^2 - 12xy + 30xz - 20yz.$$

$$3. 25m^2 + 36n^2 + p^2 - 60mn - 10mp + 12np.$$

$$4. a^2 + 16x^4 + 36y^2 - 8ax^2 + 12ay - 48x^2y.$$

$$5. x^2 + 4a^2 + b^2 + y^2 + 4ax - 2bx + 2xy - 4ab + 4ay - 2by.$$

$$6. m^2 + 4n^2 + a^2 + 9 - 4mn - 2am + 6m + 4an - 12n - 6a.$$

138. The principle by which the difference of two squares is factored has many special applications.

1. Factor $a^4 + a^2b^2 + b^4$.

SOLUTION. — Since $a^4 + a^2b^2 + b^4$ lacks $+a^2b^2$ of being a perfect square, and since the value of the polynomial will not be changed by adding a^2b^2 and also subtracting a^2b^2 , the polynomial may be written

$$a^4 + 2a^2b^2 + b^4 - a^2b^2,$$

which is the difference of two squares.

$$\begin{aligned} a^4 + a^2b^2 + b^4 &= a^4 + 2a^2b^2 + b^4 - a^2b^2 \\ &= (a^2 + b^2)^2 - (ab)^2 \\ &= (a^2 + ab + b^2)(a^2 - ab + b^2). \end{aligned}$$

2. Factor $4a^4 - 13a^2 + 9$.

$$\begin{aligned} \text{SOLUTION.} \quad 4a^4 - 13a^2 + 9 &= 4a^4 - 12a^2 + 9 - a^2 \\ &= (2a^2 - 3)^2 - a^2 \\ &= (2a^2 + a - 3)(2a^2 - a - 3). \end{aligned}$$

3. Factor $a^4 + 4$.

$$\begin{aligned}\text{SOLUTION.} \quad a^4 + 4 &= a^4 + 4a^2 + 4 - 4a^2 \\ &= (a^2 + 2)^2 - (2a)^2 \\ &= (a^2 + 2a + 2)(a^2 - 2a + 2).\end{aligned}$$

Factor the following:

- | | | |
|---------------------------------|----------------------------------|------------------------|
| 4. $x^4 + x^2y^2 + y^4$. | 11. $x^4 + x^2 + 1$. | |
| 5. $a^8 + a^4b^4 + b^8$. | 12. $n^8 + n^4 + 1$. | |
| 6. $p^4 + p^2q^2 + q^4$. | 13. $16x^4 + 4x^2y^2 + y^4$. | |
| 7. $9x^4 + 20x^2y^2 + 16y^4$. | 14. $a^4b^4 - 21a^2b^2 + 36$. | |
| 8. $4a^4 + 11a^2b^2 + 9b^4$. | 15. $c^4 + c^2a^2x^2 + d^4x^4$. | |
| 9. $16a^4 - 17a^2x^2 + x^4$. | 16. $25a^4 - 14a^2b^4 + b^8$. | |
| 10. $25x^4 - 29x^2y^2 + 4y^4$. | 17. $9a^4 + 26a^2b^2 + 25b^4$. | |
| 18. $b^4 + 64$. | 21. $a^4 + 324$. | 24. $x^4 + 64y^4$. |
| 19. $a^4 + 4b^4$. | 22. $a^8 - 16$. | 25. $4a^4 + 81$. |
| 20. $m^8 + 4$. | 23. $m^5 + 4mn^4$. | 26. $x^5y^2 + 4xy^2$. |

139. Many polynomials may be written as quadratic trinomials in which x^2 and x are replaced by polynomials.

1. Factor $9x^2 + 4y^2 + 12z^2 + 21xz + 14yz + 12xy$.

$$\begin{aligned}\text{SOLUTION.} \quad &9x^2 + 4y^2 + 12z^2 + 21xz + 14yz + 12xy \\ &= (9x^2 + 12xy + 4y^2) + (21xz + 14yz) + 12z^2 \\ &= (3x + 2y)^2 + 7z(3x + 2y) + 4z \cdot 3z \\ \S 130, \quad &= (3x + 2y + 4z)(3x + 2y + 3z).\end{aligned}$$

Factor the following:

- $a^2 + 2ab + b^2 + 8ac + 8bc + 15c^2$.
- $x^2 - 6xy + 9y^2 + 6xz - 18yz + 5z^2$.
- $m^2 + n^2 - 2mn + 7mp - 7np - 30p^2$.
- $16n^2 + 55 - 64n - 16m + m^2 + 8mn$.
- $9m^4 + k^2 - 30 + 39m^2 + 13k + 6m^2k$.
- $25a^2 + y^2 + 10x^2 + 10ay - 35ax - 7xy$.
- $4x^2 + y^2 - 6z^2 - 4xy + 2xz - yz$.
- $a^2 + b^2 + c^2 + 2ab + 2ac + 2bc + 5a + 5b + 5c + 6$.

REVIEW OF FACTORING

140. Factor the following:

- | | | |
|-----------------------------------|----------------------------------|----------------------|
| 1. $m^2 - n^2$. | 11. $p^4 + 4$. | 21. $4x^4 - 4x$. |
| 2. $x^2 - 1$. | 12. $1 + x^{12}$. | 22. $7y^4 - 175$. |
| 3. $y^4 - 1$. | 13. $y - a^4y$. | 23. $8 - 27a^3x^2$. |
| 4. $1 - x^2$. | 14. $x^2y - y^3$. | 24. $32x - 2x^3$. |
| 5. $x^{10} - 1$. | 15. $a^{13} - ab^{12}$. | 25. $6b^4 + 24$. |
| 6. $x^6 - 1$. | 16. $a^4 - 256$. | 26. $a^5 + 27a^2$. |
| 7. $1 - a^3$. | 17. $3a^3 - 3a$. | 27. $b^2 - 196$. |
| 8. $1 - b^4$. | 18. $64 - 2y^2$. | 28. $450 - 2a^3$. |
| 9. $a - a^7$. | 19. $7n^7 - 7n$. | 29. $4m^3 + .004$. |
| 10. $b^7 + b$. | 20. $a^7 - 9a$. | 30. $125 - 8x^6$. |
| 31. $x^2 - 25x + 100$. | 45. $y^2 - xy - 42x^2$. | |
| 32. $x^2 - xy - 132y^2$. | 46. $x^2 - ax - 72a^2$. | |
| 33. $ax^2 - 3ax - 4a$. | 47. $n^2 - an - 90a^2$. | |
| 34. $x^3 + 5x^2 - 6x$. | 48. $a^2b^2 + ab - 56$. | |
| 35. $3x^2 + 30x + 27$. | 49. $10a^2c + 33ac - 7c$. | |
| 36. $128a^2 - 250a^5$. | 50. $60ny^2 - 61ny - 56n$. | |
| 37. $5x^{10} + 10x^5 - 15$. | 51. $25x^2 + 60xy + 36y^2$. | |
| 38. $6x^2 - 19x + 15$. | 52. $6ax^2 + 5axy - 6ay^2$. | |
| 39. $x^{2n} + 2x^ny^p + y^{2p}$. | 53. $169x^4 - 26ax^2 + a^2x^2$. | |
| 40. $7x^2 - 77xy - 84y^2$. | 54. $a^4c^4 + a^2b^2c^2 + b^4$. | |
| 41. $y^2 - 25yx + 136x^2$. | 55. $16x^4 + 4x^2y^2 + y^4$. | |
| 42. $9x^2 - 24xy + 16y^2$. | 56. $b^4c - 13b^2c + 42c$. | |
| 43. $289x^2 - 34xy + y^2$. | 57. $2a^2 - 6ab - 140b^2$. | |
| 44. $3bx^2 + bxy - 10by^2$. | 58. $m^2n - 21mn^2 + 80n^3$. | |

59. $17x^2 + 25x - 18.$
60. $5x^2 - 26xy + 5y^2.$
61. $y^2 + 16ay - 36a^2.$
62. $8a^2 - 21ab - 9b^2.$
63. $60a^2 + 8ax - 3x^2.$
64. $30x^3 - 37x - 77.$
65. $2x^3 + 28x^2 + 66x.$
66. $a^2 + b^2 - c^2 - 2ab.$
67. $ax^2 + 10ax - 39a.$
68. $n^4 + n^2a^2b^4 + a^4b^8.$
69. $a^2z^4 + a^2x^2 + a^2.$
70. $a^2 - 16a - 17.$
71. $a^2x^2 - 4ax + 3.$
72. $b^3 + b^4y^2 + y^4.$
73. $x^7 - 2x^6 + x.$
74. $x^3 + x^2y - 41xy^2 - 105y^3.$
75. $x^2 - cx + 2dx - 2cd.$
76. $x^2y + 4x^2y - 31xy - 70y.$
77. $x^2 - 3ax + 4bx - 12ab.$
78. $ax^3 - 9ax^2 + 26ax - 24a.$
79. $12ax - 8bx - 9ay + 6by.$
80. $25x^2 - 9y^2 - 24yz - 16z^2.$
81. $x^2 - z^2 + y^2 - a^2 - 2xy + 2az.$
82. $2b^2m - 3ab^2 + 2bmx - 3abx.$
83. $a^2 + b^2 + c^2 - 2ab - 2ac + 2bc.$
84. $x^2y + 14x^2y + 43xy + 30y.$
85. $x^2y - 15x^2y + 38xy - 24y.$
86. $abx^3 + 3abx^2 - abx - 3ab.$
87. $3bmx + 2bm - 3anx - 2an.$
88. $20ax^3 - 28ax^2 + 5a^2x - 7a^2.$
89. $x^3 + 9y^2 + 25z^2 - 6xy - 10xz + 30yz.$
90. $9x^2 + y^2 + 16z^2 - 6xy - 8yz + 24zx.$
91. $x^2y^2z^2 + a^2b^2 + 1 + 2abxyz + 2xyz + 2ab.$
92. $a^2b^2 + b^2c^2 + c^2d^2 - 2ab^2c + 2abcd - 2bc^2d.$
93. $x^3 + n^4x^4 + n^8 + 2n^2x^5 + 2n^4x^4 + 2n^6x^2.$
94. $a^2b^2x^2 - a^2b^2 - b^2x^2 + b^2 - a^2x^2 + a^2 + x^2 - 1.$
95. $(a + b)^6 - 1.$
96. $a^3 - 2a^2 + 1.$
97. $b^3 - 4b^2 + 8.$
98. $x^3 - 10x^2 + 125.$
99. $8x^4 - 6x^2 - 35.$
100. $3x^5 + 96x.$
101. $(a - 2)^3 + (a - 1)^3.$
102. $12x^3 + 3x^2 - 8x - 2.$
103. $2x^3 + 10x + ax + 5a.$
104. $x^3 + 5x^2 - 29x - 105.$
105. $m^2n^2 + a^2b^2 + b^2n^2 + 2bmn^2 + 2ab^2n + 2abmn.$

106. $a^5 + a^4b + a^3b^2 + a^2b^3 + ab^4 + b^5$.
 107. $a^2b^3 - 4abx - 4x + 2ab + 4x^2$.
 108. $(a+b)^2(x-y) - (a+b)(x^2-y^2)$.
 109. $1 - x^2 + abx^2 + bx^3 - bx - ab$.
 110. $x^2 - x^3 + x^2y - xy + x^2y - xy^2$.
 111. $x^{2n-2} + b^2y^2 + 2x^{n-1}by$.
 112. $x^3 + 15x^2 + 75x + 125$.
 113. $4(ab+cd)^2 - (a^2+b^2-c^2-d^2)^2$.
114. $x^{3n} - a^{3n}$.
 115. $(a^2+b^2-c^2)^2 - 4a^2b^2$.
 116. $a^4b^2 + a^2b - 12$.
 117. $x^3 - xy - x^2y + y^2$.
 118. $x^4 - 4x^2y^2 + 2x^3 - 16y^3$.
 119. $a^4 - b^4 - (a+b)(a-b)$.
 120. $x^3 - 6x^2 + 12x - 8$.
 121. $1000x^3 - 27y^3$.
 122. $(a+x)^4 - x^4$.
 123. $1 + (x+1)^3$.
 124. $ab - bx^n + x^ny^m - ay^m$.
125. $x^9 + 4x$.
 126. $x^5 - x^2 - x^4 + x^3$.
 127. $(a+b)^4 - (b-c)^4$.
 128. $3ab(a+b) + a^3 + b^3$.
 129. $(x+y)^3 + (x-y)^3$.
 130. $a^3 - (a+b)^3$.
 131. $x^4 - 119x^2y^2 + y^4$.
 132. $m^3 + m^2 - mn - mn^2$.
 133. $(x^2 - y^2)^2 - (x^2 - xy)^2$.
 134. $x^6 - y^6 - 3x^2y^2(x^2 - y^2)$.
 135. $(x^2+6x+9)^2 - (x^2+5x+6)^2$.
136. $x^3 + (a+b-c)x^2 + (ab-ac-bc)x - abc$.
 137. Factor $32 - x^5$ by the factor law.
 138. Factor $16 + 5x - 11x^2$ by the factor law.
 139. If n is odd, factor $x^n - a^n$ by the factor law.
 140. If n is odd, factor $x^n + a^n$ by the factor law.
 141. Factor $x^3 - 6bx^2 + 12b^2x - 8b^3$ by the factor law.
 142. Discover by the factor law for what values of n , between 1 and 20, $x^n + a^n$ has no binomial factors.

EQUATIONS SOLVED BY FACTORING

141. 1. Find the value of x in $x^2 + 1 = 10$.

FIRST PROCESS

$$\begin{aligned}x^2 + 1 &= 10 \\x^2 &= 9 \\x \cdot x &= 3 \cdot 3 \quad \therefore x = 3 \\ \text{or } x \cdot x &= -3 \cdot -3 \quad \therefore x = -3 \\ \therefore x &= \pm 3\end{aligned}$$

EXPLANATION. — Transposing the known term 1 to the second member, the first member contains the second power, only, of the unknown number. Separating each member into two equal factors,

$$x \cdot x = 3 \cdot 3 \quad \text{or } x \cdot x = -3 \cdot -3.$$

Since, if $x = 3$, $x \cdot x = 3 \cdot 3$, and if $x = -3$, $x \cdot x = -3 \cdot -3$, the value of x that makes $x^2 = 9$, or that makes $x^2 + 1 = 10$, is either $+3$ or -3 ; that is, $x = \pm 3$.

Find the values of x in the following equations:

2. $x^2 + 3 = 28$.

7. $x^2 - 24 = 120$.

3. $x^2 + 1 = 50$.

8. $x^2 + 11 = 180$.

4. $x^2 - 5 = 59$.

9. $x^2 - 11 = 110$.

5. $x^2 - 7 = 29$.

10. $x^2 - b^2 = a^2 - 2ab$.

6. $x^2 + 3 = 84$.

11. $x^2 - 4n^2 = m^2 - 4mn$.

12. Find the value of x in $x^2 + 1 = 10$.

SECOND PROCESS

$$\begin{aligned}x^2 + 1 &= 10 \\x^2 - 9 &= 0 \\(x - 3)(x + 3) &= 0 \\ \therefore x - 3 &= 0, \text{ whence } x = 3 \\ \text{or } x + 3 &= 0, \text{ whence } x = -3 \\ \therefore x &= \pm 3\end{aligned}$$

EXPLANATION. — The first process is given in example 1.

In the second process, all terms are brought to the first member, which is factored as the difference of the squares of two numbers.

Since the product of the two factors is 0, one of them is equal to 0. This gives $x - 3 = 0$ or $x + 3 = 0$; whence $x = 3$ or $x = -3$; that is, $x = \pm 3$.

Solve the following equations:

13. $x^2 + 35 = 39$.

16. $x^2 - 31^2 = 0$.

14. $x^2 - 50 = 50$.

17. $x^2 - 4b^2 = 0$.

15. $x^2 + 90 = 91$.

18. $x^2 - 9n^2 = 0$.

19. $x^2 - 21 = 4.$

24. $32 - x^2 = 28.$

20. $x^2 - 56 = 8.$

25. $65 - x^2 = 16.$

21. $x^2 - 3a^2 = 6a^2.$

26. $4x^2 - 8b^2 = 8b^2.$

22. $x^2 + 5b^4 = 6b^4.$

27. $x^2 + 25 = 25 + m^2.$

23. $x^2 - 40 = 24.$

28. $x^2 - 30 = 2(2b^2 - 15).$

29. Solve $x^2 + 2am = a^2 + m^2.$

SOLUTION

$$x^2 + 2am = a^2 + m^2.$$

$$x^2 = a^2 - 2am + m^2.$$

$$x \cdot x = (a - m)(a - m),$$

or

$$x \cdot x = -(a - m) \cdot -(a - m).$$

$$\therefore x = \pm(a - m).$$

Solve the following equations:

30. $x^2 - c^2 = d^2 - 2cd.$

36. $x^2 - c^2 = 36 - 12c.$

31. $x^2 - b^2 = 4bc + 4c^2.$

37. $x^2 - 4b^2 = 36 - 24b.$

32. $x^2 - n^2 = 6n + 9.$

38. $x^2 - a^2 = 9 - 6a.$

33. $x^2 + 10a = a^2 + 25.$

39. $x^2 - b^4 = 4 - 4b^2.$

34. $x^2 - a^2 = 2a + 1.$

40. $x^2 - a^2b^2 = 2ab + 1.$

35. $x^2 - m^2 = 8m + 16.$

41. $x^2 - r^4 = b^4 - 2r^2b^2.$

42. Find the value of x in $x^2 + 4x = 45.$

FIRST PROCESS

$$x^2 + 4x = 45$$

$$x^2 + 4x - 45 = 0$$

$$(x - 5)(x + 9) = 0$$

$$\therefore x - 5 = 0$$

or

$$x + 9 = 0$$

$$\therefore x = 5 \text{ or } -9$$

SECOND PROCESS

$$x^2 + 4x = 45$$

$$x^2 + 4x + 4 = 49$$

$$(x + 2)(x + 2) = 7 \cdot 7 \text{ or } -7 \cdot -7$$

$$\therefore x + 2 = 7 \text{ or } -7$$

$$x = 7 - 2 \text{ or } -7 - 2$$

$$\therefore x = 5 \text{ or } -9$$

EXPLANATION. — The explanation given for example 12 will serve for the first process.

In the second process, it is seen that, by adding 4 to each member of the equation, the first member will become the square of the binomial $(x + 2)$. Solving for $(x + 2)$ as for x in previous examples, $x + 2 = \pm 7$; whence $x = \pm 7 - 2 = 5$ or -9 .

SUGGESTION. — In the following examples, when the coefficient of the first power of the unknown number is *even*, either of the above processes may be used; but when it is *odd*, the first process is simpler.

Solve the following equations:

43. $x^2 - 6x = 40$.

62. $x^2 + 4x + 3 = 0$.

44. $x^2 - 8x = 48$.

63. $x^2 + 6x + 8 = 0$.

45. $x^2 - 5x = -4$.

64. $x^2 - 9x + 20 = 0$.

46. $x^2 - 7x = 18$.

65. $x^2 + 11x + 30 = 0$.

47. $x^2 + 10x = 56$.

66. $x^2 + x - 132 = 0$.

48. $x^2 + 12x = 28$.

67. $32 = 4x + x^2$.

49. $x^2 - 3x = 40$.

68. $3x = 88 - x^2$.

50. $x^2 - 9x = 36$.

69. $160 = x^2 - 6x$.

51. $x^2 + 11x = 26$.

70. $4y = y^2 - 192$.

52. $x^2 - 12x = 45$.

71. $600 = y^2 - 10y$.

53. $y^2 - 15y = 54$.

72. $x^2 + 16x - 36 = 0$.

54. $y^2 - 21y = 46$.

73. $x^2 + 15x - 34 = 0$.

55. $x^2 - 10x = 96$.

74. $y^2 - 8y - 84 = 0$.

56. $y^2 - 20y = 96$.

75. $y^2 - 2ay + a^2 = 0$.

57. $y^2 + 12y = 85$.

76. $x^2 + 2bx + b^2 = 0$.

58. $y^2 + 42 = 13y$.

77. $x^2 + 4ax + 4a^2 = 0$.

59. $y^2 + 63 = 16y$.

78. $z^2 + 22z + 121 = 0$.

60. $v^2 - 60 = 11v$.

79. $x^2 - (a + b)x + ab = 0$

61. $y^2 + 140 = 72y$.

80. $x^2 + (c + d)x + cd = 0$.

81. $x^2 + (a + 2)x + 2a = 0.$ 83. $x^2 - (a - d)x - ad = 0.$

82. $y^2 - (c - n)y - nc = 0.$ 84. $x^2 - (b + 7)x + 7b = 0.$

85. $(2x + 3)(2x - 5) - (3x - 1)(x - 2) = 1.$

86. $(2x - 6)(3x - 2) - (5x - 9)(x - 2) = 4.$

87. Solve $6x^2 + 5x - 21 = 0.$

SOLUTION

$$6x^2 + 5x - 21 = 0.$$

Factoring, § 131, $(2x - 3)(3x + 7) = 0.$

$$\therefore 2x - 3 = 0,$$

$$3x + 7 = 0.$$

or

$$\therefore x = \frac{3}{2} \text{ or } -\frac{7}{3}.$$

Solve the following equations:

88. $3x^2 + 2x - 1 = 0.$ 93. $7x^2 + x - 1 = 0.$

89. $5x^2 + 4x - 1 = 0.$ 94. $2v^2 - 9v - 35 = 0.$

90. $3y^2 + y - 10 = 0.$ 95. $6y^2 - 22y + 20 = 0.$

91. $3y^2 - 4y - 4 = 0.$ 96. $3x^2 + 13x - 30 = 0.$

92. $4y^2 + 9y - 9 = 0.$ 97. $4x^2 + 13x - 12 = 0.$

98. Solve the equation $x^3 - 2x^2 - 5x + 6 = 0.$

SOLUTION

$$x^3 - 2x^2 - 5x + 6 = 0.$$

Factoring by the factor law, $(x - 1)(x - 3)(x + 2) = 0.$

$$x - 1 = 0, \text{ or } x - 3 = 0, \text{ or } x + 2 = 0.$$

$$\therefore x = 1, \text{ or } 3, \text{ or } -2.$$

Solve the following equations:

99. $x^3 - 15x^2 + 71x - 105 = 0.$ 101. $x^3 - 12x + 16 = 0.$

100. $x^3 + 10x^2 + 11x - 70 = 0.$ 102. $x^3 - 19x - 30 = 0.$

103. $x^4 + x^3 - 21x^2 - x + 20 = 0.$

104. $x^4 - 7x^3 + x^2 + 63x - 90 = 0.$

105. $x^4 + 8x^3 - x^2 - 68x + 60 = 0.$

106. $x^5 - 11x^4 + 45x^3 - 85x^2 + 74x - 24 = 0.$

HIGHEST COMMON DIVISOR

142. 1. Name all the numbers that will exactly divide both a^4 and a^3 . Which of these is of the highest degree?

2. What is the highest divisor common to b^2 and b^5 ? to x^2 and x ?

3. Since the highest divisor common to a^4 and a^3 is a^3 , to b^2 and b^5 is b^2 , and to x^2 and x is x , what is the highest divisor common to $a^4b^2x^2$ and a^3b^5x ?

4. What is the highest common divisor of $36a^2b$ and $90ab^2$? What prime factors, or divisors, are common to $36a^2b$ and $90ab^2$? How may the highest common divisor of $36a^2b$ and $90ab^2$ be found from their factors?

143. A number that exactly divides each of two or more algebraic expressions is called a **Common Divisor** of them.

The common divisors of $12a^2$ and $4a^3$ are 2, a , 4, a^2 , $2a$, $4a$, $2a^2$, and $4a^2$.

An exact divisor of an expression is a *factor* of it.

Two expressions whose only common divisor, or factor, is 1 are said to be *prime to each other*.

$3x$ and $2a$ are prime to each other; also $x + y$ and $x - y$.

144. That common divisor, or factor, of two or more algebraic expressions which is of the highest degree is called their **Highest Common Divisor**, or **Highest Common Factor**.

The highest common divisor, or factor, of $12a^2$ and $4a^3$ is $4a^2$.

The abbreviation H. C. D. is used for *Highest Common Divisor*.

The *highest* common divisor in algebra corresponds to the *greatest* common divisor in arithmetic. But there would be an inaccuracy in applying the term *greatest* common divisor to literal numbers, since letters may represent any numbers, as, for instance, fractions.

Thus, if $a = \frac{1}{2}$, then $a^2 = \frac{1}{4}$, $a^3 = \frac{1}{8}$, and the higher the degree of the literal number, the less will be its arithmetical value. Consequently, it is inaccurate to speak of a^3 , the *highest* common divisor of a^3 , $2a^4b$, and $3a^5$, as their *greatest* common divisor, because a may represent a proper fraction.

145. PRINCIPLE. — *The highest common divisor, or factor, of two or more algebraic expressions is the product of all their common prime factors.*

146. To find the highest common divisor of expressions that may be factored readily by inspection.

EXAMPLES

1. What is the highest common divisor of $12a^4b^2c$ and $8a^2b^3c^3$?

FIRST PROCESS

$$\begin{aligned} 12a^4b^2c &= 3 \times 2 \times 2 \times aaaa \times bb \times c \\ 8a^2b^3c^3 &= 2 \times 2 \times 2 \times aa \times bbb \times ccc \end{aligned}$$

$$\text{H. C. D.} = 2 \times 2 \times aa \times bb \times c = 4a^2b^2c$$

SECOND PROCESS

$$\begin{aligned} 12a^4b^2c &= 4a^2b^2c \times 3a^2 \\ 8a^2b^3c^3 &= 4a^2b^2c \times 2bc^3 \end{aligned}$$

$$\text{H. C. D.} = 4a^2b^2c$$

EXPLANATION. — Since the highest common divisor of the expressions is the product of all their common prime factors (Prin.), and since the only prime factors common to the given expressions are 2, 2, a , a , b , b , and c , their product, $4a^2b^2c$, is the highest common divisor.

SUGGESTION. — Frequently the work may be abridged by grouping common factors, as in the second process. Since $3a^2$ and $2bc^3$ are prime to each other, $4a^2b^2c$ must contain all the common factors, and be the highest common divisor.

2. What is the H. C. D. of $3x^3 - 3xy^2$ and $x^3 - 2x^2y + xy^2$?

PROCESS

$$\begin{aligned} 3x^3 - 3xy^2 &= 3x(x+y)(x-y) \\ x^3 - 2x^2y + xy^2 &= x(x-y)(x-y) \end{aligned}$$

$$\therefore \text{H. C. D.} = x(x-y)$$

EXPLANATION. — For convenience in selecting the common divisors, the expressions are resolved into their simplest factors.

Since the only common prime factors are x and $(x-y)$, the highest common divisor sought is their product, $x(x-y)$ (Prin.).

RULE. — *Separate the expressions into their prime factors.*

The product of all the common prime factors, each factor being taken the least number of times it occurs in any of the given expressions, is the highest common factor.

The factors that enter into the H. C. D. can often be selected without actually separating the expressions into their *prime* factors.

3. What is the H. C. D. of $5a^2c^2 - 5b^2c^2$ and $a^2x^3 - b^2x^3 + a^2y^3 - b^2y^3$?

SOLUTION

$$\begin{aligned} 5a^2c^2 - 5b^2c^2 &= 5c^2(a^2 - b^2) \\ a^2x^3 - b^2x^3 + a^2y^3 - b^2y^3 &= (x^3 + y^3)(a^2 - b^2) \\ \hline \therefore \text{H. C. D.} &= a^2 - b^2 \end{aligned}$$

Find the highest common divisor of

4. $10x^3y^2$, $10x^2y^3$, and $15xy^4z$.
5. $70a^6b^3$, $21a^4b^4$, and $35a^4b^5$.
6. $8m^7n^3$, $28m^6n^4$, and $56m^5n^2$.
7. $4b^3cd$, $6b^2c^2$, and $24abc^3$.
8. $10(x - y)^4z^3$ and $15(z - y)(x - y)^3$.
9. $4(a + b)^3(a - b)$ and $b(a + b)^2(a - b)^2$.
10. $3(a^2 - b^2)^3$ and $a(a - b)(a^2 - b^2)$.
11. $x^2 - 2x - 15$ and $x^2 - x - 20$.
12. $x^4 - y^4$, $x^2 - y^2$, and $x + y$.
13. $a^2 + 7a + 12$ and $a^2 + 5a + 6$.
14. $x^3 + y^3$ and $x^2 + 2xy + y^2$.
15. $a^2 - x^2$ and $a^2 - 2ax + x^2$.
16. $a^2 - b^2$ and $a^2 + 2ab + b^2$.
17. $x^4 + x^2y^2 + y^4$ and $x^2 + xy + y^2$.
18. $x^3 + y^3$, $x^5 + y^5$, and $x^2y + xy^2$.
19. $a^4 + a^2b^4 + b^8$ and $3a^2 - 3ab^2 + 3b^4$.
20. $a^3 - x^2$, $a^2 + 2ax + x^2$, and $a^3 + x^3$.
21. $ax - y + xy - a$ and $ax^2 + x^2y - a - y$.
22. $a^2b - b - a^2c + c$ and $ab - ac - b + c$.

- 23. $1 - 4x^2$, $1 + 2x$, and $4a - 16ax^2$.
- 24. $(a - b)(b - c)$ and $(c - a)(a^2 - b^2)$.
- 25. $24x^2y^3 + 8x^2y^3$ and $8x^2y^3 - 8x^2y^3$.
- 26. $6x^2 + x - 2$ and $2x^2 - 11x + 5$.
- 27. $16x^2 - 25$ and $20x^2 - 9x - 20$.
- 28. $x^2 - 6x + 5$ and $x^3 - 5x^2 + 7x - 3$.
- 29. $x^2 - 4$ and $x^3 - 10x^2 + 31x - 30$.
- 30. $x^2 - 9$ and $x^3 - 12x^2 + 41x - 42$.
- 31. $x^3 - 4x + 3$ and $x^3 + x^2 - 37x + 35$.

147. To find the highest common divisor of expressions that cannot be factored readily by inspection.

1. What are the exact divisors of ab ? Will they be factors of 2 times ab ? of a times ab ? of m times ab ?

2. If a number is an exact divisor of an expression, what will be its relation to any number of times the expression?

3. What common divisor have ax and ay ? any number of times ax and ay , as $m \cdot ax$ and $n \cdot ay$?

4. If two numbers have a common divisor, what divisor has their sum? their difference? the sum or difference of any number of times the numbers?

5. What is the highest common divisor of $2am(x+y)$ and $3bm(x+y)$? How will it be affected, if the second number is multiplied by 7 or z ? by 2 or a ? How will it be affected, if the first number is multiplied by 5? by b ?

6. By what numbers may one of two expressions be multiplied without affecting their highest common divisor?

7. How will the highest common divisor of $2am(x+y)$ and $3bm(x+y)$ be affected, if the first number is divided by 2? by a ? by m ? by $(x+y)$? How, if the second number is divided by b ? by m ?

8. By what numbers may one of two expressions be divided without affecting their highest common divisor?

148. PRINCIPLES.—1. *A divisor of an expression is a divisor of any number of times the expression.* Hence, by § 85,

2. *A common divisor of two expressions is a divisor of their sum, of their difference, and of the sum or the difference of any number of times the expressions; also,*

3. *The highest common divisor of two expressions is not affected by multiplying or dividing either of them by numbers that are not factors of the other.*

EXAMPLES

1. Find the H. C. D. of $x^3 + 5x + 6$ and $4x^3 + 21x^2 + 30x + 8$.

PROCESS

$$\begin{array}{r}
 x^3 + 5x + 6 \quad 4x^3 + 21x^2 + 30x + 8(4x + 1 \\
 \underline{4x^3 + 20x^2 + 24x} \\
 x^2 + 6x + 8 \\
 \underline{x^2 + 5x + 6} \\
 x + 2 \quad x^2 + 5x + 6(x + 3 \\
 \underline{x^2 + 2x} \\
 3x + 6 \\
 \underline{3x + 6} \\
 0
 \end{array}$$

\therefore H. C. D. = $x + 2$.

EXPLANATION.—Since the highest common divisor cannot be higher than $x^2 + 5x + 6$, it will be $x^2 + 5x + 6$, if $x^2 + 5x + 6$ is exactly contained in $4x^3 + 21x^2 + 30x + 8$. By trial, it is found that it is not exactly contained in $4x^3 + 21x^2 + 30x + 8$, since there is a remainder of $x + 2$. Therefore, $x^2 + 5x + 6$ is not the highest common divisor.

Since $x^2 + 5x + 6$ contains the highest common divisor, $(4x + 1)$ times $x^2 + 5x + 6$ will also contain the highest common divisor (Prin. 1); and since both $4x^3 + 21x^2 + 30x + 8$ and $(4x + 1)(x^2 + 5x + 6)$ contain the highest common divisor, their difference, $x + 2$, must contain the highest common divisor (Prin. 2). Hence, the highest common divisor cannot be higher than $x + 2$.

$x + 2$ will be the highest common divisor, if it is exactly contained in $x^2 + 5x + 6$, since, if it is contained in $x^2 + 5x + 6$, it will be contained in any number of times $x^2 + 5x + 6$, as $(4x + 1)(x^2 + 5x + 6)$ (Prin. 1); and in the sum of $(4x + 1)(x^2 + 5x + 6)$ and $x + 2$, or $4x^3 + 21x^2 + 30x + 8$ (Prin. 2). By trial, $x + 2$ is found to be exactly contained in $x^2 + 5x + 6$. Therefore, $x + 2$ is the highest common divisor of the given expressions.

2. Find the H. C. D. of $6x^3 + 33x - 63$ and $2x^3 + 11x^2 - x - 30$.

PROCESS

$$\begin{array}{r}
 3) 6x^3 + 33x - 63 \quad | \quad 2x^3 + 11x^2 - x - 30(x \\
 \underline{2x^3 + 11x - 21} \quad | \quad 2x^3 + 11x^2 - 21x \\
 10) 20x - 30 \\
 \quad 2x - 3) 2x^3 + 11x - 21(x + 7 \\
 \underline{2x^3 - 3x} \\
 14x - 21 \\
 \underline{14x - 21}
 \end{array}$$

\therefore H. C. D. = $2x - 3$.

SUGGESTION. — Since only common factors are sought, factors that are not common to the given expressions, as 3 and 10, may be *rejected* from any expression before it is used as a divisor (Prin. 3).

3. Find the H. C. D. of

$$2x^3 + 5x^2 - 22x + 15 \text{ and } 5x^3 + 18x^2 - 33x + 10.$$

PROCESS

$$\begin{array}{r}
 2x^3 + 5x^2 - 22x + 15) 5x^3 + 18x^2 - 33x + 10 \\
 \underline{2} \\
 10x^3 + 36x^2 - 66x + 20(5 \\
 \underline{10x^3 + 25x^2 - 110x + 75} \\
 11) 11x^2 + 44x - 55 \\
 \underline{x^2 + 4x - 5} \\
 2x^3 + 5x^2 - 22x + 15(2x - 3 \\
 \underline{2x^3 + 8x^2 - 10x} \\
 - 3x^2 - 12x + 15 \\
 \underline{- 3x^2 - 12x + 15}
 \end{array}$$

\therefore H. C. D. = $x^2 + 4x - 5$.

SUGGESTION. — When the first term of the divisor is not contained in the first term of the dividend an integral number of times, fractional quotients may be avoided by multiplying the polynomial taken for the dividend by some number not a factor of the divisor (Prin. 3). In the above example the simplest factor that may thus be introduced is 2, if $5x^3 + 18x^2 - 33x + 10$ is taken for the dividend; or 5, if $2x^3 + 5x^2 - 22x + 15$ is taken for the dividend.

4. Find the H. C. D. of

$$30 amx^2 - 21 amx - 99 am \text{ and } 42 abx^3 + 33 abx^2 - 45 abx.$$

PROCESS

$$30 amx^2 - 21 amx - 99 am) 42 abx^3 + 33 abx^2 - 45 abx$$

Reject m (Prin. 3). Reject bx (Prin. 3).

$$30ax^2 - 21ax - 99a) 42ax^2 + 33ax - 45a$$

Reserve the common factor $3a$ as a factor of the H. C. D.

$$10x^2 - 7x - 33 \mid 14x^2 + 11x - 15$$

$$\begin{array}{r} 7 \\ \hline 70x^2 - 49x - 231 \end{array} \quad \begin{array}{r} 5 \\ \hline 70x^2 + 55x - 75(1) \\ 70x^2 - 49x - 231 \end{array}$$

$$52) 104x + 156$$

$$2x + 3$$

$$0x^2 + 7x$$

$$\frac{2x+5}{10x^2-5x-33} = \frac{1}{x} = \frac{33}{33x} = \frac{11}{11x}$$

$$\therefore \text{H.C.D.} \equiv 5a(2x+3), \quad \underline{10x-7x-35}$$

SUGGESTION.—Since each of the polynomials contains a factor not found in the other, these two factors may be rejected (Prin. 3). Consequently, m is rejected from the first polynomial, and bx from the second.

To simplify the process the common factor $3a$ is removed and *reserved* as a factor of the H. C. D.

5. Find the H. C. D. of $9x^2 - 35x + 24$ and $29x - 8x^2 - 15$.

PROCESS

$$-8x^2 + 29x - 15$$

$$9x^2 - 35x + 24$$

$$x^2 - 6x + 9)(9x^2 - 35x + 24)(9$$

$$9x^2 - 54x + 81$$

10\10 57

$x = 57$

$$\therefore \text{H.C.D.} = x - 3.$$

SUGGESTION.—Since the sum or the difference of two expressions contains their highest common divisor (Prin. 2), it is evident that at the outset a simpler expression that will contain the highest common divisor may be obtained by adding the given expressions, giving $x^2 - 6x + 9$.

Find the H. C. D. of

6. $x^2 + 2x - 24$ and $2x^2 + 7x - 30$.

7. $2x^2 - x - 21$ and $4x^2 + 4x - 63$.

8. $3x^2 + 10x - 8$ and $6x^2 - 7x + 2$.

9. $2x^3 - 6x^2 + 7x - 6$ and $2x^3 + 4x^2 - 3x + 9$.

10. $x^3 + 9x^2 + 26x + 24$ and $2x^3 + 14x^2 + 20x$.

11. Find the H. C. D. of

$3ax^3 - 4ax^2 - 13ax + 14a$ and $3abx^3 + 5abx^2 - 10abx - 42ab$.

FIRST PROCESS

$3ax^3 - 4ax^2 - 13ax + 14a) 3abx^3 + 5abx^2 - 10abx - 42ab$

Reserve the common factor a as a factor of the H. C. D.

$$\begin{array}{r}
 3x^3 - 4x^2 - 13x + 14) 3bx^3 + 5bx^2 - 10bx - 42b(b \\
 \underline{3bx^3 - 4bx^2 - 13bx + 14b} \\
 b) 9bx^2 + 3bx - 56b \\
 \underline{9x^2 + 3x - 56} 3x^3 - 4x^2 - 13x + 14 \\
 3 \\
 \underline{9x^3 - 12x^2 - 39x + 42} (x \\
 9x^3 + 3x^2 - 56x \\
 \underline{-15x^2 + 17x + 42} \\
 -15x^2 + 17x + 42) 9x^2 + 3x - 56 \\
 5 \\
 \underline{45x^2 + 15x - 280} (-3 \\
 45x^2 - 51x - 126 \\
 \underline{22) 66x - 154} \\
 3x - 7) -15x^2 + 17x + 42 (-5x - 6 \\
 \underline{-15x^2 + 35x} \\
 -18x + 42 \\
 \underline{-18x + 42}
 \end{array}$$

$\therefore \text{H. C. D.} = a(3x - 7).$

Since the arrangement of the dividend, divisor, and quotient may be either: Divisor) Dividend (Quotient; or Quotient) Dividend (Divisor; by using these two arrangements *alternately*, the above process may be more compactly written as follows:

SECOND PROCESS

$$3ax^3 - 4ax^2 - 13ax + 14a) 3abx^3 + 5abx^2 - 10abx - 42ab$$

Reserve the common factor a as a factor of the H.C.D.

$$\begin{array}{r|l}
 3x^3 - 4x^2 - 13x + 14 \\
 3 \\
 \hline
 9x^3 - 12x^2 - 39x + 42 \\
 x \quad 9x^3 + 3x^2 - 56x \\
 \hline
 - 15x^2 + 17x + 42 \\
 -5x \quad -15x^2 + 35x \\
 \hline
 -18x + 42 \\
 -6 \quad -18x + 42 \\
 \hline
 \end{array}
 \begin{array}{r|l}
 3bx^3 + 5bx^2 - 10bx - 42b \\
 3bx^3 - 4bx^2 - 13bx + 14b \\
 \hline
 b) 9bx^2 + 3bx - 56b \\
 9x^2 + 3x - 56 \\
 5 \\
 \hline
 45x^2 + 15x - 280 \\
 45x^2 - 51x - 126 \\
 \hline
 22) 66x - 154 \\
 3x - 7 \\
 \hline
 \end{array}
 \begin{array}{l}
 b \\
 -3
 \end{array}$$

SUGGESTION. — When the quotient consists of more than one term, for convenience each term is placed opposite the corresponding part of the dividend or product.

RULE. — Divide one expression by the other, and if there is a remainder, divide the divisor by it; then divide the preceding divisor by the last remainder, and so on, until there is no remainder. The last divisor will be the highest common divisor.

If any remainder does not contain the letter of arrangement, the expressions have no common divisor in that letter.

If more than two expressions are given, find the highest common divisor of any two, then of this divisor and another, and so on. The last divisor will be the highest common divisor.

1. If either expression contains a monomial factor not found in the other, it should be rejected before beginning the process.
2. A common factor of the expressions should be removed before beginning the division, but it must appear as a factor of the highest common divisor.
3. When necessary, to avoid fractional quotients, any dividend or divisor may be multiplied or divided by any number not a factor of the other.
4. The highest common divisor has an ambiguous sign. For, if a positive divisor is contained in a dividend, the same negative divisor also will be contained in that dividend, but the signs of the quotient will be changed. It is not customary to write both divisors.

149. The principle, that the exact divisor reached by the process given in the rule is the highest common divisor, may be proved as follows :

Let A and B represent any two polynomials freed of monomial factors, the degree of B being not higher than that of A .

Divide A by B , and let the quotient be m and the remainder D ; divide B by D , and let the quotient be n and the remainder E ; divide D by E , and let the quotient be r and the remainder zero; that is, let E be an exact divisor of D .

It is to be proved that E is the highest common divisor of A and B .

$$\begin{array}{r}
 \text{PROCESS} \\
 B)A(m \\
 \underline{mB} \\
 D)B(n \\
 \underline{nD} \\
 E)D(r \\
 \underline{rE} \\
 0
 \end{array}$$

Since the minuend is equal to the subtrahend plus the remainder,

$$A = mB + D, \text{ and } A - mB = D;$$

$$B = nD + E, \text{ and } B - nD = E; \text{ and } D = rE.$$

Since the division has terminated, E is a common divisor of D and nD (Prin. 1); also of D and $nD + E$, or B (Prin. 2); also of B and mB (Prin. 1); also of B and $mB + D$, or A (Prin. 2). That is, E is a common divisor of B and A .

Every common divisor of A and B is a divisor of mB (Prin. 1); and of $A - mB$, or D (Prin. 2). Therefore, every common divisor of A and B is a divisor of nD (Prin. 1); and of $B - nD$, or E (Prin. 2).

But, since no divisor of E can be of higher degree than E itself, E is the highest common divisor of A and B .

150. The principle, that the highest common divisor of several expressions may be obtained by finding the highest common divisor of two of them, then of this result and a third expression, and so on, may be proved as follows :

Let P be the highest common divisor of A and B , and Q the highest common divisor of P and a third expression C .

Then, since P contains all the common factors of A and B , and Q contains of these particular factors only such as are factors of C also, Q is the highest common divisor of A , B , and C .

This method may be extended to embrace any number of expressions.

Find the H. C. D. of

12. $2x^3 - 7x^2 + 2x + 3$ and $2x^3 + 7x^2 - 5x - 4$.
13. $9x^3 + 18x^2 - x - 10$ and $3x^3 + 13x^2 + 2x - 8$.
14. $1 - 2x - 5x^2 + 6x^3$ and $1 + 5x + 2x^2 - 8x^3$.
15. $1 - 4x + x^2 + 6x^3$ and $1 + 3x - 6x^2 - 8x^3$.
16. $1 - x - 14x^2 + 24x^3$ and $36x^3 - 24x^2 + x + 1$.
17. $m^3 - 4m^2 - 20m + 48$ and $m^3 - m^2 - 14m + 24$.
18. $3a^3 + 20a^2 - a - 2$ and $3a^3 + 17a^2 + 21a - 9$.
19. $8ax^2 + 22ax + 15a$ and $6bx^2 + 11bx + 3b$.
20. $20b^2c - 2bc - 4c$ and $8a^2b^3c - 4a^2bc + a^2c$.
21. $21ax - 17ax^2 - 5ax^3 + ax^4$ and $7ax + 34ax^2 - 5ax^3$.
22. $x^3 - 7x + 6$, $x^4 - 2x^3 - 9x^2 + 18x$, $x^3 + x^2 - 4x - 4$.
23. $x^3 - 5x + 4$, $x^4 - 2x^3 + 1$, $x^6 + 4x^3 - 3x - 2$.
24. $1 + 4x^2 + 5x^3$, $2 + 5x + 3x^4$, $x^6 - 4x^4 + 5x^2 - 2$.
25. $3 + x - 8x^2 + 4x^3$, $3 - 8x - 8x^2 + 8x^3$, $16x^4 - 48x^3 + 81$.
26. $x^3 - 6x^2 - 5x - 14$, $x^3 - 10x^2 + 20x + 7$, $x^4 - 310x - 231$.
27. Find the H. C. D. of $x^6 + x^4 + x^3 - x - 2$ and $2x^6 + x^5 - x^3 - x^2 - 1$.

PROCESS BY DETACHED COEFFICIENTS

$$\begin{array}{l}
 1 \left| \begin{array}{r} 1 + 0 + 1 + 1 + 0 - 1 - 2 \\ 1 - 2 - 3 - 1 + 2 + 3 \\ \hline 2 + 4 + 2 - 2 - 4 - 2 \\ 2 - 4 - 6 - 2 + 4 + 6 \\ \hline 8 + 8 + 0 - 8 - 8 \\ 1 + 1 + 0 - 1 - 1 \end{array} \right| \begin{array}{r} 2 + 1 + 0 - 1 - 1 + 0 - 1 \\ 2 + 0 + 2 + 2 + 0 - 2 - 4 \\ \hline 1 - 2 - 3 - 1 + 2 + 3 \\ \hline 1 + 1 + 0 - 1 - 1 \\ - 3 - 3 + 0 + 3 + 3 \\ \hline - 3 - 3 + 0 + 3 + 3 \end{array} \left| \begin{array}{l} 2 \\ 1 \\ 1 \\ -3 \end{array} \right.
 \end{array}$$

\therefore H. C. D. = $x^4 + x^3 - x - 1$.

Find the H. C. D. of

28. $x^5 - x^4 - 2x^3 - x^2 + x + 2$ and $x^5 + 3x^4 + 3x^3 + x^2 - x - 1$.
29. $x^5 + x^4 - x^2 - 7x - 4$ and $2x^5 + 3x^4 + 3x^3 + 3x^2 - 7x - 4$.

30. $x^5 - 2x^4 - 2x^3 - 11x^2 - x - 15$ and $2x^5 - 7x^4 + 4x^3 - 15x^2 + x - 10$.
31. $a^5 - 3a^4 - 3a^3 - 3a^2 - 19a - 15$ and $a^5 + 3a^4 - 3a^3 + 9a^2 - a - 15$.
32. $5a^5 + a^4 - 11a^3 + 9a^2 - 8a + 4$ and $2a^5 - a^4 - 5a^3 + 8a^2 - 4a$.
33. $x^5 - 5x + 4$ and $x^4 - x^3 - 3x^2 - 5x - 12$.
34. $a^3 + 3a^2 - 2a - 6$ and $a^5 + 4a^4 + 4a^3 + 4a^2 - a - 12$.
35. $1 - 4a^3 + 3a^4$ and $1 + a - a^2 - 5a^3 + 4a^4$.
36. $2 - a + 3a^2 + 5a^3 - a^4$ and $4 - 4a + a^2 - 9a^4$.
37. $y^5 + 13y^2 + 20y - 14$ and $7 - 3y - 20y^2 + 2y^3 - y^5$.
38. $6x^3 - 11x^2 - 35x$, $30x^3 - 115x + 35$, $2x^3 - 5x^2 - 5x - 7$.



LOWEST COMMON MULTIPLE



151. 1. What number exactly contains 2, 5, a , and b , or is a multiple of 2, 5, a , and b ?

2. What different prime factors must enter into every number that will contain $4a^3b$, a^2b^2 , and $10ab^3$, or must be found in every common multiple of $4a^3b$, a^2b^2 , and $10ab^3$?

3. What is the lowest power of a that common multiples of $4a^3b$, a^2b^2 , and $10ab^3$ can contain? What is the lowest power of b ? of 2? of 5?

What, then, is the lowest common multiple of $4a^3b$, a^2b^2 , and $10ab^3$?

To what is the *lowest common multiple* of two or more expressions equal?

152. An expression that exactly contains each of two or more given expressions is called a **Common Multiple** of them.

$6abx$ is a common multiple of a , $3b$, $2x$, and $6abx$. These numbers may have other common multiples, as $12abx$, $6a^2b^2x$, $18a^3bx^2$, etc.

153. The expression of *lowest degree* that will exactly contain each of two or more given expressions is called their **Lowest Common Multiple**.

$6abx$ is the lowest common multiple of a , $3b$, $2x$, and $6abx$.

The abbreviation L. C. M. is used for *Lowest Common Multiple*.

The *lowest* common multiple in algebra corresponds to the *least* common multiple in arithmetic. But, since letters may represent any numbers, as, for instance, numbers not prime to each other or fractions, the term *least* is not applicable to algebraic common multiples.

Thus, the algebraic lowest common multiple of a^2b^2 , ab^3 , and bx is a^2b^3x . If $a = 4$, $b = 3$, and $x = 2$, a^2b^3x , the *lowest* common multiple of the given expressions, is equal to 864. If, however, the values of a , b , and x are substituted for those letters, the given expressions become 144, 108, and 6; and their *least* common multiple is 432.

It is thus seen that the *lowest* common multiple of two or more expressions is not necessarily their *least* common multiple.

154. PRINCIPLE. — *The lowest common multiple of two or more algebraic expressions is the product of all their different prime factors, using each factor the greatest number of times it occurs in any of the expressions.*

155. To find the lowest common multiple of expressions that may be factored readily by inspection.

EXAMPLES

1. What is the L. C. M. of $12x^2yz^4$, $6a^2xy^2$, and $8axyz^2$?

PROCESS

$$12x^2yz^4 = 2 \cdot 2 \cdot 3 \cdot x^2 \cdot y \cdot z^4$$

$$6a^2xy^2 = 2 \cdot 3 \cdot a^2 \cdot x \cdot y^2$$

$$8axyz^2 = 2 \cdot 2 \cdot 2 \cdot a \cdot x \cdot y \cdot z^2$$

$$\text{L. C. M.} = 2 \cdot 2 \cdot 2 \cdot 3 \cdot a^2 \cdot x^2 \cdot y^2 \cdot z^4 = 24a^2x^2y^2z^4$$

EXPLANATION. — The lowest common multiple of the numerical coefficients is found as in arithmetic. It is 24.

The literal factors of the lowest common multiple are each letter with the highest exponent it has in any of the given expressions (Prin.). They are, therefore, a^2 , x^2 , y^2 , and z^4 .

The product of the numerical and literal factors, $24a^2x^2y^2z^4$, is the lowest common multiple of the given expressions.

2. What is the L. C. M. of $x^3 - 2xy + y^2$, $x^2 - y^2$, and $x^3 + y^3$?

PROCESS

$$\begin{array}{rcl}
 x^3 - 2xy + y^2 & = & (x - y)(x - y) \\
 x^2 - y^2 & = & (x - y)(x + y) \\
 x^3 + y^3 & = & (x + y)(x^2 - xy + y^2) \\
 \hline
 \text{L. C. M.} & = & (x - y)^2(x + y)(x^2 - xy + y^2) \\
 & = & (x - y)^2(x^3 + y^3)
 \end{array}$$

RULE.—Factor the expressions as far as may be necessary to discover their different prime factors.

Find the product of all their different prime factors, using each factor the greatest number of times it occurs in any of the given expressions.

The factors of the L. C. M. may often be selected without separating the expressions into their prime factors.

Find the L. C. M. of

3. a^3x^2y , a^2xy^3 , and ax^2y .
4. $10a^2b^2c^2$, $5ab^2c$, and $25b^3c^3a^3$.
5. $16a^2b^3c$, $24c^3de$, and $36a^4b^2a^2e^2$.
6. $18a^2br^2$, $12p^2q^2r$, and $54ab^2p^3q$.
7. $x^m y^2$, $x^{m-1} y^3$, $x^{m-3} y^4$, and $x^{m+1} y$.
8. $x^2 - y^2$ and $x^3 + 2xy + y^2$.
9. $x^2 - y^2$ and $x^2 - 2xy + y^2$.
10. $x^2 - y^2$, $x^2 + 2xy + y^2$, and $x^2 - 2xy + y^2$.
11. $a^2 - n^2$ and $3a^3 + 6a^2n + 3an^2$.
12. $x^4 - 1$ and $a^2x^2 + a^2 - b^2x^2 - b^2$.
13. $a^3 + 1$, $ab - b$, $a^2 + a$, and $a^3 - 1$.
14. $2x + y$, $2xy - y^2$, and $4x^2 - y^2$.
15. $1 + x$, $x - x^2$, $1 + x^2$, and $x^2(1 - x)$.
16. $3 + a$, $9 - a^2$, $3 - a$, and $5a + 15$.
17. $a - b$, $b - c$, $b + a$, and $a^2 - b^2$.

18. $2x + 2$, $5x - 5$, $3x - 3$, and $x^2 - 1$.
19. $3x - 9y$, $3x^2 + 27y^2$, and $2x + 6y$.
20. $16b^2 - 1$, $12b^2 + 3b$, $20b - 5$, and $2b$.
21. $1 - 2x^2 + x^4$, $(1 - x)^2$, and $1 + 2x + x^2$.
22. $1 - a$, $1 + a$, $1 + a^2$, $1 + a^4$, and $1 + a^8$.
23. $xy - y^2$, $x^2 + xy$, $xy + y^2$, and $x^2 + y^2$.
24. $x^3 - y^3$, $x^2 + xy + y^2$, and $x^2 - xy$.
25. $b^2 - 5b + 6$, $b^2 - 7b + 10$, and $b^2 - 10b + 16$.
26. $x^2 + 7x - 8$, $x^2 - 1$, $x + x^2$, and $3ax^2 - 6ax + 3a$.
27. $a^2 - x^2$, $a - 2x$, $a^2 + 2ax$, and $a^3 - 3a^2x + 2ax^2$.
28. $m^3 - x^3$, $m^2 + mx$, $m^2 + mx + x^2$, and $(m + x)x^2$.
29. $x^2 - 3x + 2$, $x^2 + 4x + 4$, $x^2 + 3x + 2$, and $x^2 - 1$.
30. $x^2 - y^2$, $x^4 + x^2y^2 + y^4$, $x^3 + y^3$, and $x^2 + xy + y^2$.
31. $x^3 + x^2y + xy^2 + y^3$ and $x^3 - x^2y + xy^2 - y^3$.
32. $a^2 + 4a + 4$, $a^2 - 4$, and $a^4 - 16$.
33. $a^2 - (b + c)^2$, $b^2 - (c + a)^2$, and $c^2 - (a + b)^2$.
34. $1 - a + a^2$, $1 + a + a^2$, and $1 + a^3 + a^4$.
35. $a^4 + 4$ and $a^4 - 2a^3 + 4a - 4$.
36. $a^6 - b^3$ and $a^6 + a^4b^2 + b^4$.
37. $x^6 + y^6$ and $a^2x^2 - b^2y^2 + a^2y^2 - b^2x^2$.
38. $a^4 - 2a^2b + a^2b^2 - 9b^4$ and $a^4 + 5a^2b^2 + 9b^4$.
39. $a^4 - a^2 + 1$, $a^6 + 1$, $a^4 + a^2 + 1$, and $a^2 - 1$.
40. Find the lowest common multiple of $x^3 + 6x^2 + 5x - 12$ and $x^3 - 8x^2 + 19x - 12$.

SOLUTION

$$\begin{aligned} x^3 + 6x^2 + 5x - 12 &= (x - 1)(x^2 + 7x + 12) \\ &= (x - 1)(x + 3)(x + 4). \end{aligned}$$

$$\begin{aligned} x^3 - 8x^2 + 19x - 12 &= (x - 1)(x^2 - 7x + 12) \\ &= (x - 1)(x - 3)(x - 4). \end{aligned}$$

$$\begin{aligned} \therefore \text{L. C. M.} &= (x - 1)(x + 3)(x - 3)(x + 4)(x - 4) \\ &= (x - 1)(x^2 - 9)(x^2 - 16). \end{aligned}$$

SUGGESTION.—In solving the following the Factor Law will be found useful.

41. $x^3 - 6x^2 + 11x - 6$ and $x^3 - 9x^2 + 26x - 24$.

42. $x^3 - 5x^2 - 4x + 20$ and $x^3 + 2x^2 - 25x - 50$.

43. $x^3 + 3x^2 - 4$ and $x^3 + x^2 - x - 1$.

44. $x^3 - 4x^2 + 5x - 2$ and $x^3 - 8x^2 + 21x - 18$.

45. $x^3 + 5x^2 + 7x + 3$ and $x^3 - 7x^2 - 5x + 75$.

46. $x^3 + 2x^2 - 4x - 8$, $x^3 - x^2 - 8x + 12$, $x^3 + 4x^2 - 3x - 18$.

47. $x^3 - 9x^2 + 23x - 15$, $x^3 + x^2 - 17x + 15$, $x^3 + 7x^2 + 7x - 15$.

48. $x^3 + 7x^2 + 14x + 8$, $x^3 + 3x^2 - 6x - 8$, $x^3 + x^2 - 10x + 8$.

156. To find the lowest common multiple of expressions that cannot be factored readily by inspection.

$$x^2 - 3x + 2 = (x - 1)(x - 2). \quad (1)$$

$$x^2 - 5x + 6 = (x - 2)(x - 3). \quad (2)$$

$$x^2 - 7x + 12 = (x - 3)(x - 4). \quad (3)$$

$$\text{L. C. M.} = (x - 1)(x - 2)(x - 3)(x - 4).$$

1. Find the lowest common multiple of expressions (1) and (2) from their factors; from the product of their factors. By what factor of the two expressions must the product be divided to obtain the lowest common multiple?

2. How, then, may the lowest common multiple of *two* expressions be found?

3. Since $(x - 1)(x - 2)(x - 3)$ is the lowest common multiple of the first two expressions, what factor of the third expression must the lowest common multiple of all the expressions contain?

157. PRINCIPLES.—1. *The lowest common multiple of two expressions is equal to their product divided by their highest common divisor; or, it is equal to either of them multiplied by the quotient of the other divided by the highest common divisor.*

2. *The lowest common multiple of several expressions may be obtained by finding the lowest common multiple of two of them; then of this result and a third expression; and so on.*

Proof of Principle 1.

Let F be the highest common divisor, or factor, of A and B , and L their lowest common multiple. Let F be contained a times in A and b times in B , or let $A = aF$ and let $B = bF$.

It is to be proved that $L = \frac{A \times B}{F}$, or $A \times \frac{B}{F}$, or $B \times \frac{A}{F}$.

Since F contains *all* the common factors of A and B , a and b have no common factors; consequently, since $A = aF$ and $B = bF$,

$$L = abF.$$

Multiplying by F , $FL = abFF$;

but $A \times B = aF \times bF = abFF$.

Therefore, Ax. 1, $FL = A \times B$,

and $L = \frac{A \times B}{F}$, or $\frac{A}{F} \times B$, or $A \times \frac{B}{F}$

Proof of Principle 2.

Let L be the lowest common multiple of A and B , and M the lowest common multiple of L and a third expression C .

It is to be proved that M is the lowest common multiple of A , B , and C .

Since L is the expression of lowest degree that is exactly divisible by both A and B , and M is the expression of lowest degree that is exactly divisible by both L and C , M is the expression of lowest degree that is exactly divisible by A , B , and C .

EXAMPLES

1. Find the L. C. M. of $x^3 + 6x^2 + 11x + 6$ and $x^3 - 4x^2 + x + 6$.

PROCESS

$$\begin{aligned} \text{Prin. 1, L. C. M.} &= \frac{(x^3 + 6x^2 + 11x + 6)(x^3 - 4x^2 + x + 6)}{\text{H. C. D.}} \\ &= \frac{(x+1)(x^2 + 5x + 6)(x+1)(x^2 - 5x + 6)}{x+1} \\ &= (x+1)(x^2 + 5x + 6)(x^2 - 5x + 6) \end{aligned}$$

$$\text{or} \quad (x+1)(x+2)(x+3)(x-2)(x-3)$$

$$\text{or} \quad (x+1)(x^2 - 4)(x^2 - 9)$$

Find the L. C. M. of

$$2. \quad 4a^3 + 7a^2 + 10a - 3 \text{ and } 4a^3 + 9a^2 + 14a + 3.$$

$$3. \quad 2a^3 - 11a^2 + 18a - 14 \text{ and } 2a^3 + 3a^2 - 10a + 14.$$

$$4. \quad 5x^3 - 11x^2 + 3x + 12 \text{ and } 5x^3 - 19x^2 + 27x - 12.$$

5. $4x^3 - 14x^2 + 22x - 8$ and $2x^4 - 3x^3 - x^2 + 12x$.
6. $6a^3 + 3a^2 - 15a - 75$ and $2a^3 + 11a^2 + 25a + 25$.
7. $4a^3 - 27a^2 - 2a + 15$ and $2a^4 - 9a^3 - 28a^2 - 15a$.
8. $3c^3 - 11c^2 - 32c - 16$ and $3c^3 - 19c^2 + 8c + 16$.
9. $4x^4 - 7x^3 + 7x^2 - 11x + 6$ and $2x^4 + x^3 - x^2 - x - 6$.
10. $x^4 - x^3 - 3x + 9$ and $3ax^4 - 3ax^3 - 18ax^2 + 45ax - 27a$.
11. $20x^3 + 40x^2 + 25x + 125$ and $6x^3 + 7x^2 + 10x + 25$.
12. $12m^3 - 18m^2 + 26m - 10$ and $15m^3 - 9m^2 + 19m + 10$.
13. $6a^3x - 5a^2x - 18ax - 8x$ and $6a^3b - 13a^2b - 6ab + 8b$.
14. $4x^3 + 4x^2y - 5xy^2 + 25y^3$ and $4x^3 - 16x^2y + 25xy^2 - 25y^3$.
15. $10a^3 + 29a^2 - 36a + 9$ and $8a^3 + 34a^2 + 9a - 9$.
16. $4x^4 - 17x^2y^2 + 4y^4$ and $2x^4 - x^2y - 3x^2y^2 - 5xy^3 - 2y^4$.
17. $5x^4 + 8x^3 - 27x^2 + 14x - 10$ and $3x^4 + 4x^3 - 17x^2 + 14x - 10$.
18. $2x^4 - 9x^3 + 18x^2 - 18x + 9$ and $3x^4 - 11x^3 + 17x^2 - 12x + 6$.
19. $3a^4 + 13a^3 - 19a^2 + 12a - 4$ and $4a^4 + 22a^3 - 2a^2 + 2a + 4$.
20. $6x^2 + 5x - 6$, $8x^2 + 10x - 3$, $10x^2 + 9x - 9$.
21. $x^4 - 2x^3 + x^2 - 1$, $x^4 - x^2 + 2x - 1$, $x^4 - 3x^2 + 1$.
22. $x^4 - 7x^2 + 9$, $x^4 + 2x^3 + x^2 - 9$, $x^4 - x^2 - 6x - 9$.
23. $x^4 - 4x^3 + 4x^2 - 16$, $x^4 - 12x^2 + 16$, $x^4 - 4x^3 + 16x - 16$.
24. $x^4 - 4x^3 + 4x^2 - 25$, $x^4 - 4x^3 + 20x - 25$, $x^4 - 14x^2 + 25$.
25. $4x^4 + 5x^2 - x - 1$, $6x^4 + x^3 + 8x^2 - 1$, $36x^4 - 13x^2 + 1$.
26. $10x^4 + 7x^3 - 33x^2 + 26x - 10$ and $2x^4 + 7x^3 + 5x^2 - 4x - 10$.
27. $16x^4 + 16x^3 - 48x^2 - 36x + 27$ and $24x^4 + 20x^3 - 74x^2 - 45x + 45$.
28. $10x^4 + 7x^3 + 2x^2 - x - 2$ and $6x^3 + 5x^2 + 4x + 1$.
29. $5x^4 + 3x^3 + 6x^2 + x + 3$ and $15x^3 + 14x^2 + x + 12$.
30. $2x^3 - x^2 - 3x + 2$, $4x^3 + 6x^2 - 2x - 4$, $4x^3 - 5x + 2$.
31. $x^3 - 1$, $2x^3 + 2x^2 - 5x + 1$, $x^3 - 3x + 2$.

FRACTIONS

158. A fraction is *expressed* by two numbers, one called the *numerator*, written above a line, and the other the *denominator*, written below the line. Thus, $\frac{a}{b}$ is a fraction.

If a and b represent positive integers, as 3 and 4, the fraction $\frac{a}{b}$ is equal to $\frac{3}{4}$; that is, it represents 3 of the 4 equal parts of anything. This is the arithmetical notion of a fraction.

But, since a and b may represent any numbers, positive or negative, integral or fractional, rational or irrational, $\frac{a}{b}$ may represent an expression like $\frac{4}{5\frac{1}{2}}$. Since a thing cannot be divided into $5\frac{1}{2}$ equal parts, algebraic fractions are not accurately described by the definition commonly given in arithmetic. But, since an expression like $\frac{20}{4}$, regarded as 20 fourths, is equivalent to 5, or $20 \div 4$, it is evident that the numerator of a fraction may be regarded as a dividend, and the denominator as its divisor; and this interpretation of a fraction is broad enough to include the fraction $\frac{a}{b}$ when a and b represent any numbers whatever. Hence,

The expression of an unexecuted division, in which the dividend is the numerator and the divisor the denominator, is an Algebraic Fraction.

The fraction $\frac{a}{b}$ is read, 'a divided by b.'

159. The numerator and denominator of a fraction are called its **Terms**.

160. An expression, some of whose terms are integral and some fractional, is called a **Mixed Number**, or a **Mixed Expression**.

$a - \frac{a-b}{c}$, $\frac{x^2}{a^2} - 2 + \frac{a^2}{x^2}$, and $a - b + \frac{1}{ab}$ are mixed expressions.

REDUCTION OF FRACTIONS

161. The process of changing the form of an expression without changing its value is called **Reduction**.

162. To reduce fractions to higher or lower terms.

163. A fraction is in its **Lowest Terms** when its terms have no common divisor.

164. 1. How many eighths are there in $\frac{1}{2}$? in $\frac{x}{2}$? in $\frac{3x}{4}$? in $\frac{5ab}{4}$? in $\frac{4x}{16}$? in $\frac{9a}{24}$? in $\frac{8xy}{32}$?

2. How many tenths are there in $\frac{a}{2}$? in $\frac{6a}{5}$? in $\frac{6a}{20}$?

3. If a dividend is multiplied by any number, as 2, and the divisor is multiplied by the same number, how is the quotient affected?

4. If a dividend is divided by any number, as 2, and the divisor is divided by the same number, how is the quotient affected?

5. Since a fraction may be regarded as an indicated division, what may be done to the terms of a fraction without changing the *value* of the fraction?

165. PRINCIPLE. — *Multiplying or dividing both terms of a fraction by the same number does not change the value of the fraction.*

The proof of the principle is as follows:

Let a and b be any two numbers, a the dividend, b the divisor, and $\frac{a}{b}$ the quotient. Also, let m be any number.

It is to be proved that $\frac{a}{b} = \frac{ma}{mb}$.

Since the quotient multiplied by the divisor equals the dividend,

$$\frac{a}{b} \times b = a. \quad (1)$$

$$\text{Multiplying (1) by } m, \text{ Ax. 4, } \frac{a}{b} \times mb = ma. \quad (2)$$

$$\text{Dividing (2) by } mb, \text{ Ax. 5, } \frac{a}{b} = \frac{ma}{mb} \quad (3)$$

Hence, the terms of any fraction, as $\frac{a}{b}$, may be multiplied by any number, or the terms of any fraction, as $\frac{ma}{mb}$, may be divided by any number, without changing the value of the fraction.

EXAMPLES

1. Reduce $\frac{a}{a+b}$ to a fraction whose denominator is $a^2 - b^2$.

PROCESS

$$(a^2 - b^2) \div (a + b) = a - b$$

$$\therefore \frac{a}{a+b} = \frac{a(a-b)}{(a+b)(a-b)} = \frac{a^2 - ab}{a^2 - b^2}$$

EXPLANATION. — Since the required denominator is $(a - b)$ times the given denominator, both terms of the fraction must be multiplied by $(a - b)$ (Prin.).

2. Reduce $\frac{21 a^2 x^2 y}{30 a^3 x z}$ to its lowest terms.

PROCESS

$$\frac{21 a^2 x^2 y}{30 a^3 x z} = \frac{7 xy}{10 az}$$

EXPLANATION. — Since a fraction is in its lowest terms when its terms have no common divisor, the given fraction may be reduced to its lowest terms by removing in succession all common divisors of its numerator and denominator (Prin.), as, 3, a , a , and x ; or by dividing the terms by their highest common divisor, $3 a^2 x$.

3. Change $\frac{a}{2b}$ to a fraction whose denominator is $4b^2$.

4. Change $\frac{5a}{6}$ to a fraction whose denominator is 42.

5. Change $\frac{3x}{11b}$ to a fraction whose denominator is $55b$.

6. Change $\frac{3a}{14x}$ to a fraction whose denominator is $84xy$.

7. Change $\frac{4a^2}{5y}$ to a fraction whose denominator is $20y^3$.

8. Change $\frac{x-3}{x-1}$ to a fraction whose denominator is $(x-1)^2$.
9. Change $\frac{2x-5}{2x+5}$ to a fraction whose denominator is $(2x+5)^2$.
10. Change $\frac{a}{3-a}$ to a fraction whose numerator is $3a+a^2$.
11. Reduce $\frac{a-b}{a+b}$ to a fraction whose denominator is a^2-b^2 .
12. Reduce $\frac{x-y}{2x+y}$ to a fraction whose numerator is x^2-y^2 .
13. Reduce $\frac{d-c}{b-a}$ to a fraction whose denominator is $a-b$.
14. Reduce $\frac{-1}{x-2}$ to a fraction whose denominator is $4-x^2$.

Reduce the following to their lowest terms :

- | | |
|---|--|
| 15. $\frac{a^2xy^2}{a^3xy}$ | 22. $\frac{-25x^2y^5z^3}{-100x^4y^3}$ |
| 16. $\frac{m^3n^3}{am^2n^4}$ | 23. $\frac{x^{m+2}y^2}{x^my^4}$ |
| 17. $\frac{a^2b^2x^2}{b^3xy^2}$ | 24. $\frac{x^{m-2}a^2}{x^m}$ |
| 18. $\frac{16m^2nx^2z^2}{40am^3yz^3}$ | 25. $\frac{x^{m+1}y}{xy^{m+1}}$ |
| 19. $\frac{210bc^2d}{750ab^2c}$ | 26. $\frac{x^{m-n+1}}{ax}$ |
| 20. $\frac{35a^2bcd^3}{42ab^2cd^4}$ | 27. $\frac{a^2b^7}{3a^{2a}b}$ |
| 21. $\frac{77a^7x^2b^3y}{121a^3b^5c^2}$ | 28. $\frac{a^{m+2}y^{2a}}{2a^ry^{4r}}$ |

29. $\frac{ax^{m-n+1}}{bx^{m-n}}$.
30. $\frac{na^{n-2}b^2}{n(n-2)ab}$.
31. $\frac{a^2 - b^2}{a^2 + 2ab + b^2}$.
32. $\frac{a^2 - 2ab + b^2}{a^2 - b^2}$.
33. $\frac{4a^2 - 9x^2}{8a^3 + 27x^3}$.
34. $\frac{3a^2 + 3ab}{a^4 + ab^3}$.
35. $\frac{3x^2y - 6xy}{x^4y - 8xy}$.
36. $\frac{3a^2b - 3b^3}{2a^3b - 2b^4}$.
37. $\frac{4a^3 - ab^2}{8a^4 + ab^3}$.
38. $\frac{2x^2y^2 - 8y^4}{4x^2y - 32y^4}$.
39. $\frac{a^4bc - b^5c}{3a^6b + 3b^7}$.
40. $\frac{10nx + 10ny}{25nx^2 - 25ny^2}$.
41. $\frac{x^{n+2} - x^n}{x^{n+3} - x^n}$.
42. $\frac{a^{n+4} - a^ny^4}{a^{n+3} + a^{n+1}y^2}$.
43. $\frac{x^4y + x^2y^3 + y^5}{x^6 - y^6}$.
44. $\frac{x^4y - x^2y^3 + y^5}{x^6 + y^6}$.
45. $\frac{a^2 - 11a + 24}{a^2 - a - 6}$.
46. $\frac{x^3 - 6x^2 + 5x}{x^3 + 2x^2 - 35x}$.
47. $\frac{7x - 2x^2 - 3}{2x^2 + 7x - 4}$.
48. $\frac{a(a + 2b)^4}{b(a^2 - 4b^2)^2}$.
49. $\frac{a^3 + 2a^2b + ab^2}{a^5 - 2a^3b^2 + ab^4}$.
50. $\frac{x^2 - 2x^4 + x^6}{x^2 - x^6}$.
51. $\frac{x^3 + 5x^2 - 6x}{2x^2 - 2}$.
52. $\frac{x^3 - 7x + 6}{x^4 - 10x^2 + 9}$.
53. $\frac{x^3 - 21x + 20}{x^4 - 26x^2 + 25}$.
54. $\frac{x^3 + 3x^2 + 3x + 1}{x^3 + x^2 - 4x - 4}$.
55. $\frac{a^3 - 3a^2b + 3ab^2 - b^3}{3a^2b - 3ab^2}$.
56. $\frac{3a^2 + 4ax - 4x^2}{9a^2 - 12ax + 4x^2}$.
57. $\frac{2ax - ay - 4bx + 2by}{4ax - 2ay - 2bx + by}$.
58. $\frac{9x^3 - 13a^2x - 4a^3}{3bx + 3xy - 4ab - 4ay}$.
59. $\frac{m - m^2 - n + mn}{m - mn + n^2 - n}$.
60. $\frac{am - an - m + n}{am - an + m - n}$.

61. Reduce $\frac{3x^3 - 16x^2 + 25x - 12}{3x^3 - 8x^2 - 7x + 12}$ to its lowest terms.

SOLUTION

The process of finding the H. C. D. of the terms of the fraction can be shortened in some instances by finding the sum or the difference of the terms, since the result will either be the H.C.D. or some multiple of it, § 148.

$$\frac{3x^3 - 16x^2 + 25x - 12}{3x^3 - 8x^2 - 7x + 12}$$

Subtracting the numerator from the denominator,

$$\begin{aligned} & 8x^2 - 32x + 24 \\ &= 8(x^2 - 4x + 3). \end{aligned}$$

By trial, $x^2 - 4x + 3$ is found to be the H. C. D.

Dividing the terms of the fraction by $x^2 - 4x + 3$, the fraction in its

lowest terms is $\frac{3x - 4}{3x + 4}$.

Reduce the following to their lowest terms:

62. $\frac{x^3 + 5x^2 - 9x - 45}{x^3 + 3x^2 - 25x - 75}$.

63. $\frac{x^3 + 2x^2 - 23x - 60}{x^3 - 11x^2 - 10x + 200}$.

64. $\frac{4x^3 + 7x^2 + 10x - 3}{4x^3 + 9x^2 + 14x + 3}$.

65. $\frac{x^3 + 5x^2 + 8x + 6}{x^3 + 3x^2 + 4x + 2}$.

66. $\frac{3x^3 - 7x^2 + 4}{5x^3 - 17x^2 + 16x - 4}$.

67. $\frac{5x^3 - 14x^2 + 22x + 5}{5x^3 - 18x^2 + 34x - 15}$.

68. $\frac{x^3 - 6x^2y + 2xy^2 + 3y^3}{x^3 + 6x^2y - 2xy^2 - 5y^3}$.

69. $\frac{a^2 + b^2 + 2c^2 + 2ab + 3ac + 3bc}{a^2 + b^2 + c^2 + 2ab + 2ac + 2bc}$.

$$70. \frac{a^2 + b^2 + c^2 + 2ab - 2ac - 2bc}{a^2 + b^2 - c^2 + 2ab}.$$

$$71. \frac{a^2 + b^2 + c^2 - 2ab - 2ac + 2bc}{a^2 + b^2 + 5c^2 - 2ab - 6ac + 6bc}.$$

$$72. \frac{4a^2 + 9b^2 + 16c^2 + 12ab + 16ac + 24bc}{4a^2 - 9b^2 + 16c^2 + 16ac}.$$

$$73. \frac{ab(x^2 + y^2) + xy(a^2 + b^2)}{ab(x^2 - y^2) + xy(a^2 - b^2)}.$$

166. Signs in fractions.

167. The sign written before the dividing line of a fraction is called the **Sign of the Fraction**.

It belongs to the fraction as a *whole*, and not to either the numerator or the denominator.

In $-\frac{x}{3z}$ the sign of the fraction is $-$, while the signs of x and $3z$ are $+$.

168. An expression like $\frac{-a}{-b}$ indicates a process in division, in which the quotient is to be found by dividing a by b and prefixing the sign according to the law of signs in division; that is,

$$\begin{array}{ll} \frac{-a}{-b} = +\frac{a}{b}, & \frac{+a}{+b} = +\frac{a}{b}, \\ \frac{-a}{+b} = -\frac{a}{b}, & \frac{+a}{-b} = -\frac{a}{b}. \end{array}$$

By comparing the above fractions and their values the following principles may be deduced:

169. PRINCIPLES. — 1. *The signs of both the numerator and the denominator of a fraction may be changed without changing the sign of the fraction.*

2. *The sign of either the numerator or the denominator of a fraction may be changed, provided the sign of the fraction is changed.*

When either term of a fraction is a polynomial, its sign is changed by changing the sign of each of its terms. Thus, the sign of $a - b$ is changed by writing it $-a + b$, or $b - a$.

EXAMPLES

Reduce to fractions having positive numbers in both terms:

1. $\frac{-3}{-4}$ 3. $\frac{-a-x}{2x}$ 5. $-\frac{-a-b}{c+d}$ 7. $-\frac{2-m}{2+n}$
2. $\frac{2}{-5}$ 4. $\frac{-4c}{-b-y}$ 6. $-\frac{-2}{-a-y}$ 8. $\frac{-4(a+b)}{5(-x-y)}$

170. Since, from the laws of signs, changing the signs of an even number of factors does not change the sign of the product, it follows that:

PRINCIPLE 1. — *The signs of an even number of factors of the numerator or of the denominator may be changed without changing the sign of the fraction.*

Since, from the laws of signs, changing the signs of an odd number of factors changes the sign of the product, it follows from Prin. 2, § 169, that:

PRINCIPLE 2. — *The signs of an odd number of factors of the numerator or of the denominator may be changed, provided the sign of the fraction is changed.*

EXAMPLES

1. Show that $\frac{-b}{b-a} = \frac{b}{a-b}$.
2. Show that $\frac{-a}{b-a+c} = \frac{a}{a-b-c}$.
3. Show that $\frac{2}{a(b-a)} = -\frac{2}{a(a-b)}$.
4. Show that $\frac{1}{(a-b)(c-b)} = -\frac{1}{(a-b)(b-c)}$.
5. Show that $\frac{m-n}{(a-c)(b-a)} = \frac{m-n}{(c-a)(a-b)}$.
6. Show that $\frac{1}{(b-a)(c-b)(a-c)} = -\frac{1}{(a-b)(b-c)(c-a)}$.
7. Show that $\frac{n-m}{(y-x)(z-y)(x-z)} = \frac{m-n}{(x-y)(y-z)(z-x)}$.

171. To reduce a fraction to an integral or a mixed expression.

- How many units are there in $\frac{25}{4}$? in $\frac{32}{5}$? in $\frac{17}{8}$?
- How many units are there in $\frac{4a+8b}{4}$? in $\frac{10a-5b}{5}$?

EXAMPLES

- Reduce $\frac{ax+b}{x}$ to a mixed number.

PROCESS EXPLANATION. — Since, § 158, a fraction may be regarded as an expression of unexecuted division, by performing the division indicated the fraction is changed into the form of a mixed number.

$$\frac{ax+b}{x} = a + \frac{b}{x}$$

When the degree of the numerator is lower than the degree of the denominator, the fraction cannot be reduced to an integral or mixed expression.

Reduce the following to integral or mixed expressions:

- $\frac{a^3 - 3a^2 + a - 1}{a^2}$
- $\frac{4x^2 - 8x^3 + 2x - 1}{2x}$
- $\frac{ab - bc - cd + d^2}{b}$
- $\frac{a^2x^3 - ax^3 - x - 1}{ax}$
- $\frac{x^2 - x - 15}{x - 4}$
- $\frac{x^3 - 2xy - y^3}{x - y}$
- $\frac{x^3 - 6xy + 4y^3}{2xy}$
- $\frac{a^2 - 2a - 26}{a + 5}$
- $\frac{a^3 + 2ab + b^3}{a + b}$
- $\frac{x^3 + y^3}{x^2 - xy + y^2}$
- $\frac{x^3 + 5x^2 + 3x - 6}{x + 2}$
- $\frac{a^3 + 9a^2 + 24a + 22}{a + 3}$
- $\frac{x^3 - 6x^2 + 14x - 9}{x - 2}$
- $\frac{x^4 - 3x^2 + 5x - 1}{x - 3}$
- $\frac{a^4 + 3a^2b^2 + b^4}{a^2 + b^2}$
- $\frac{4x^2 + 22x + 21}{2x + 4}$
- $\frac{x^3 - 3x^2 + 4x - 3}{x - 4}$
- $\frac{a^2 + 2ab + b^2 + c^2}{a + b + c}$
- $\frac{a^3 - 6a^2b + 12ab^2 - 10b^3}{a - 2b}$
- $\frac{x^4 + 4x^3y + 6x^2y^2 + 4xy^3}{x + y}$

172. To reduce an integral or a mixed expression to a fraction.

1. How many fifths are there in 6? in 10? in a ? in $3b$?

2. How many fifths are there in $6\frac{4}{5}$? in $a + \frac{4a}{5}$? in $2a + \frac{a}{5}$?

EXAMPLES

1. Reduce $a + \frac{b}{c}$ to a fractional form.

PROCESS

$$a = \frac{ac}{c}$$

$$a + \frac{b}{c} = \frac{ac}{c} + \frac{b}{c} = \frac{ac + b}{c}$$

EXPLANATION. — Since $1 = c \div c$, $a = ac \div c$.
Since, § 158,

$\frac{b}{c}$ means $b \div c$, $a + \frac{b}{c} = ac \div c + b \div c$.

§ 104, 3,

$$= (ac + b) \div c.$$

§ 158,

$$= \frac{ac + b}{c}.$$

RULE. — *Multiply the integral part by the denominator of the fraction; to this product add the numerator when the sign of the fraction is plus, subtract it when the sign of the fraction is minus, and write the result over the denominator.*

If the *sign of the fraction* is $-$, the signs of *all the terms in the numerator* must be changed when it is subtracted.

Reduce the following to fractional forms:

2. $a + \frac{b}{2}.$

8. $a - \frac{a^2 - ab}{b}.$

3. $x - \frac{y}{2}.$

9. $a - \frac{a - b - c}{2}.$

4. $5c + \frac{a^2 - c^2}{c}.$

10. $b - \frac{b - a - c}{2}.$

5. $3b - \frac{a + 2b^2}{b}.$

11. $a - \frac{c - a - b}{2}.$

6. $4x + \frac{1 - x}{3}.$

12. $\frac{a + b + c}{2} - c.$

7. $4b + \frac{2bc - 1}{c}.$

13. $6 + \frac{a^2 + ab}{ab - b^2}.$

$$14. a + x - \frac{x^2}{a - x}.$$

$$17. x^2 + xy + y^2 + \frac{y^3}{x - y}.$$

$$15. x + 5 - \frac{x^2 + 4}{x - 4}.$$

$$18. 3a - 6b - \frac{16b^2 - 5c^2}{a + 2b}.$$

$$16. a^2 - ab + b^2 - \frac{b^3}{a + b}.$$

$$19. x - x^2 - x^3 - \frac{x^4}{1 + x}.$$

173. To reduce dissimilar fractions to similar fractions.

1. Into what fractions having the same denominator may $\frac{2}{3}$, $\frac{3}{4}$, and $\frac{5}{6}$ be changed?

2. Express $\frac{1}{3x}$, $\frac{2}{5x}$, and $\frac{4}{15x}$ by fractions whose common denominator is the lowest common multiple of the given denominators.

174. Fractions that have the same denominator are called **Similar Fractions**.

175. Fractions that have different denominators are called **Dissimilar Fractions**.

176. PRINCIPLE. — *The lowest common denominator of two or more fractions is the lowest common multiple of their denominators.*

The abbreviation L.C.D. is used instead of *Lowest Common Denominator*.

EXAMPLES

1. Reduce $\frac{a}{3bc}$ and $\frac{c}{6ab}$ to similar fractions having their lowest common denominator.

$$\begin{aligned} \text{PROCESS} \\ \frac{a}{3bc} &= \frac{a \times 2a}{3bc \times 2a} = \frac{2a^2}{6abc} \\ \frac{c}{6ab} &= \frac{c \times c}{6ab \times c} = \frac{c^2}{6abc} \end{aligned}$$

EXPLANATION. — Since the L. C. D. of the given fractions is the lowest common multiple of their denominators (Prin.), the lowest common multiple of their denominators must be found. This is $6abc$.

To reduce the fractions to equivalent fractions having the common denominator $6abc$, § 165, the terms of each fraction must be multiplied by the quotient of $6abc$ divided by the denominator of the fraction.

RULE. — Find the lowest common multiple of the denominators of the fractions for the lowest common denominator.

Divide this denominator by the denominator of the first fraction, and multiply the terms of the fraction by the quotient.

Proceed in a similar manner with the other fractions.

All mixed expressions should first be reduced to the fractional form, and all fractions to their lowest terms.

Reduce to similar fractions having their L. C. D.:

$$2. \frac{x}{2} \text{ and } \frac{3y}{5}.$$

$$8. \frac{m-n}{a}, 2, \frac{a}{m+n}.$$

$$3. \frac{ab}{4} \text{ and } \frac{x}{y}.$$

$$9. \frac{4bc}{3a^2b}, \frac{3ac}{4b^2c}, \frac{7ab}{6ac^2}.$$

$$4. \frac{a}{3} \text{ and } \frac{4x}{6y}.$$

$$10. \frac{3ab}{8a^2c}, \frac{7a^2}{4b^2c}, \frac{5}{a^3bc^2}.$$

$$5. \frac{2a}{5b} \text{ and } \frac{3x}{4a}.$$

$$11. \frac{3}{x^2y^2}, \frac{-6}{x^2y^3}, \frac{3}{x^4y^2}.$$

$$6. \frac{a^2b}{c^2d} \text{ and } \frac{ab^2}{cd^2}.$$

$$12. \frac{x+y}{2}, \frac{2x-y}{4}, \frac{x^2-y^2}{6}.$$

$$7. \frac{1}{2xy} \text{ and } \frac{3}{4ay}.$$

$$13. \frac{a-2b}{x}, \frac{2b-a}{y}, a - \frac{c}{3x}.$$

$$14. \frac{x^2}{x^2-1}, \frac{x}{x+1}, \frac{x}{x-1}.$$

$$15. \frac{a^3}{a^4-16}, \frac{a}{a^2+4}, \frac{2a}{a^2-4}.$$

$$16. \frac{4a}{a-b}, \frac{3b}{b+a}, \frac{1}{a^2-b^2}.$$

$$17. \frac{a}{1-ax}, \frac{x}{1+ax}, \frac{-ax}{ax+1}.$$

$$18. \frac{1}{x^2+7x+10}, \frac{1}{x^2+x-2}, \frac{1}{x^2+4x-5}.$$

$$19. \frac{a+5}{a^2-4x+3}, \frac{a-2}{a^2-8x+15}, \frac{a+1}{a^2-6a+5}.$$

ADDITION AND SUBTRACTION OF FRACTIONS

177. 1. Find the value of $\frac{4}{x} + \frac{3}{x}$; $\frac{7}{a} - \frac{2}{a}$; $\frac{a}{y} + \frac{b}{y}$; $\frac{1}{x} + \frac{2}{y}$.

2. What kind of fractions can be added or subtracted without changing their form?

3. What must be done to dissimilar fractions before they can be added or subtracted? How are dissimilar fractions made similar?

178. PRINCIPLE. — *Only similar fractions may be united by addition or subtraction into one term.*

EXAMPLES

1. Find the algebraic sum of $\frac{a}{b} + \frac{c}{b} - \frac{d}{b}$.

PROCESS

$$\S 104, 3, \quad \frac{a}{b} + \frac{c}{b} - \frac{d}{b} = \frac{a + c - d}{b}$$

2. What is the sum of $\frac{3x}{4}$, $\frac{7x}{10}$, and $\frac{5y}{12}$?

PROCESS

$$\begin{aligned} & \frac{3x}{4} + \frac{7x}{10} + \frac{5y}{12} \\ &= \frac{45x}{60} + \frac{42x}{60} + \frac{25y}{60} \\ &= \frac{87x + 25y}{60} \end{aligned}$$

EXPLANATION. — Since the fractions are dissimilar, they must be made similar before they can be united into one term (Prin.).

The lowest common denominator is 60. $\frac{3x}{4} = \frac{45x}{60}$; $\frac{7x}{10} = \frac{42x}{60}$; $\frac{5y}{12} = \frac{25y}{60}$.

Therefore, the sum is $\frac{45x}{60} + \frac{42x}{60} + \frac{25y}{60} = \frac{45x + 42x + 25y}{60} = \frac{87x + 25y}{60}$.

3. Find the algebraic sum of $\frac{5x-1}{8} - \frac{3x-2}{7} + \frac{x-5}{4}$.

PROCESS

$$\begin{aligned}\frac{5x-1}{8} - \frac{3x-2}{7} + \frac{x-5}{4} &= \frac{35x-7}{56} - \frac{(24x-16)}{56} + \frac{14x-70}{56} \\ &= \frac{35x-7-(24x-16)+14x-70}{56} \\ &= \frac{35x-7-24x+16+14x-70}{56} \\ &= \frac{25x-61}{56}\end{aligned}$$

SUGGESTION. — When a fraction is preceded by the sign $-$, it is expedient for the beginner to inclose the numerator in parenthesis, if it is a polynomial, as is shown above.

RULE. — Reduce the fractions to similar fractions having their lowest common denominator.

Change the signs of all the terms of the numerators of fractions preceded by the sign $-$, then find the algebraic sum of the numerators, and write it over the common denominator.

1. Reduce the resulting fraction to its lowest terms, if necessary.
2. The integral and fractional parts of mixed expressions may be united separately.
3. An integer may be expressed as a fraction whose denominator is 1.

Find the algebraic sum of

4. $\frac{a-b}{ab} + \frac{b-c}{bc}$.

9. $\frac{b-c}{bc} - \frac{a-c}{ac}$.

5. $\frac{a+b}{a-b} + \frac{a-b}{a+b}$.

10. $\frac{a+b}{a-b} - \frac{a-b}{a+b}$.

6. $\frac{ax}{a-x} + a$.

11. $\frac{5a^2+b^2}{a^2-b^2} - 2$.

7. $a+b + \frac{a^2+b^2}{a-b}$.

12. $x+y - \frac{x^2+y^2}{x-y}$.

8. $\frac{x+1}{x^2+x+1} + \frac{1}{x-1}$.

13. $\frac{x}{x-2} - \frac{x-2}{x+2}$.

Simplify:

14. $\frac{2x+1}{3} + \frac{x-2}{4} - \frac{x-3}{6} + \frac{5-x}{2}.$
15. $\frac{x-2}{6} - \frac{x-4}{9} + \frac{2-3x}{4} - \frac{2x+1}{12}.$
16. $\frac{x-1}{3} - \frac{x-2}{18} - \frac{4x-3}{27} + \frac{1-x}{6}.$
17. $\frac{2-6x}{5} + \frac{4x-1}{2} - \frac{5x-3}{6} - \frac{1-x}{3}.$
18. $\frac{x+3}{4} - \frac{x-2}{5} + \frac{x-4}{10} - \frac{x+3}{6}.$
19. $\frac{x-4}{3} - \frac{x-6}{8} + 2 - \frac{x+8}{6}.$
20. $\frac{2-3x}{3} - \frac{3-2x}{4} + x - \frac{1-4x}{5}.$
21. $\frac{1-2a}{5} + \frac{2a-1}{4} - \frac{2a-a^2+1}{8}.$
22. $\frac{3+x-x^2}{4} - \frac{1-x+x^2}{6} - \frac{1-2x-2x^2}{3}.$
23. $\frac{3a}{5} + \frac{b}{2} - 3 + \frac{1}{b}.$
24. $\frac{a-b}{a+b} - \frac{a+b}{a-b} + \frac{6ab}{a^2-b^2}.$
25. $\frac{a+x}{a-x} + \frac{a-x}{a+x} + \frac{2ax}{a^2-x^2}.$
26. $x+1 + \frac{x^3-3}{x-1}.$
27. $1 - \frac{ax-bx+ab}{x^2}.$
28. $\frac{a+1}{a^2+a+1} + \frac{a-1}{a^2-a+1}.$
29. $3x + \frac{5}{ax} - \left(2x + \frac{3}{ax}\right).$
30. $\frac{a-b}{2(a+b)} + \frac{a^2+b^2}{a^2-b^2} - \frac{a}{a-b}.$
31. $\frac{a+1}{a^2-9} - \frac{6}{a+5} + \frac{10}{a+3}.$
32. $2a-3b - \frac{4a^2+9b^2}{2a+3b}.$
33. $3a-2x - \frac{8a^2-4x}{3a+2x}.$
34. $m - \frac{m^2+n^2}{m-n} + n.$
35. $\frac{1}{2(x-1)} - \frac{1}{2(x+1)} + \frac{1}{x^2}.$
36. $\frac{1}{x} + 1 + \frac{2x}{1+x} - 2.$

$$37. \frac{a}{a-2} - \frac{a-2}{a+2} + \frac{3}{4-a^2}.$$

SUGGESTION. — By Prin. 1, § 100, $\frac{3}{4-a^2} = \frac{-3}{a^2-4}$.

$$38. \frac{a+1}{a-1} + \frac{a-1}{a+1} + \frac{2a^2}{1-a^2}.$$

$$39. \frac{5x+2}{x^2-4} + \frac{2}{x-2} - \frac{3}{2-x}.$$

$$40. \frac{x(a+x)}{a-x} - \frac{3ax-x^2}{x-a} + 4a.$$

$$41. \frac{a+b}{a-b} - \frac{a^2+b^2}{b^2-a^2} + \frac{b-a}{a+b}.$$

$$42. \frac{a}{x(x-a)} + \frac{2x}{a(a-x)} + \frac{1}{x-a}.$$

$$43. \frac{1}{a^3+8} - \frac{1}{8-a^3} + \frac{1}{4-a^2}.$$

$$44. \frac{x-1}{x^2-5x+6} + \frac{x}{x-2} - \frac{3}{x+1}.$$

$$45. \frac{2}{x} + \frac{3}{1-2x} - \frac{2x-3}{4x^2-1} + \frac{1}{1+2x}.$$

$$46. \frac{x^2+a^2}{x^3-a^3} + \frac{x+a}{x^2+ax+a^2} - \frac{2}{x-a}.$$

$$47. \frac{3m}{(m-2x)^2} + \frac{m+2x}{(m+x)(m-2x)} - \frac{5}{m+x}.$$

$$48. \frac{3}{y^2-my-12m^2} - \frac{2}{y^2-5my+4m^2}.$$

$$49. \frac{ab}{a^2-ab+b^2} - \frac{ab}{a^2+ab+b^2} - 1.$$

$$50. \frac{1}{x^2-3x+2} - \frac{1}{x^2+2x-3} + \frac{1}{x^2+x-6}.$$

$$51. \frac{2}{x^2+5x+6} - \frac{1}{x^2+6x+8} - \frac{1}{x^2+7x+12}.$$

$$52. \frac{5(x-3)}{x^2-x-2} - \frac{2(x+2)}{x^2+4x+3} - \frac{x-1}{6-x-x^2}.$$

$$53. \frac{a-b-c}{a+b+c} - \frac{a-b+c}{a+b-c} + \frac{4ac}{(a+b)^2 - c^2}.$$

$$54. \text{Simplify } \frac{a^2+2a+1}{a^2-2a+1} - 2 + \frac{a^2-2a+1}{a^2+2a+1}.$$

SOLUTION

$$\begin{aligned} \frac{a^2+2a+1}{a^2-2a+1} - 2 + \frac{a^2-2a+1}{a^2+2a+1} &= \left(1 + \frac{4a}{a^2-2a+1}\right) - 2 + \left(1 - \frac{4a}{a^2+2a+1}\right) \\ &= \frac{4a}{a^2-2a+1} - \frac{4a}{a^2+2a+1} = \frac{16a^2}{(a-1)^2(a+1)^2}. \end{aligned}$$

SUGGESTION. — Frequently, by reducing one or more of the given fractions to mixed numbers, the integers cancel each other and the numerators are thus simplified.

Simplify:

$$55. \frac{a^2+2ab+b^2}{a^2+b^2} - 1 + \frac{2ab}{a^2-b^2}.$$

$$56. \frac{a^2+3ab+2b^2}{a^2+3ab-4b^2} - \frac{a^2-13b^2}{a^2-16b^2}.$$

$$57. \frac{x^2+x+1}{x^2-x+1} - \frac{x^2-x+1}{x^2-2x+1}.$$

$$58. \frac{x^3+x^2+x+1}{x^3-x^2+x-1} - 1 - \frac{3}{x-1}.$$

$$59. \frac{x+1}{x-1} + \frac{x-1}{x+1} - \frac{x+2}{x-2} - \frac{x-2}{x+2}.$$

$$60. \frac{x+3}{x-3} - \frac{x-3}{x+3} + \frac{x+4}{x-4} - \frac{x-4}{x+4}.$$

$$61. \frac{a}{a-b} - \frac{a}{a+b} - \frac{2ab}{a^2+b^2} - \frac{4ab^3}{a^4+b^4}.$$

SOLUTION

Combining first two fractions, $\frac{a}{a-b} - \frac{a}{a+b} = \frac{2ab}{a^2-b^2}. \quad (1)$

Combining (1) with the third fraction, $\frac{2ab}{a^2-b^2} - \frac{2ab}{a^2+b^2} = \frac{4ab^3}{a^4-b^4}. \quad (2)$

Combining (2) with the fourth fraction, $\frac{4ab^3}{a^4-b^4} - \frac{4ab^3}{a^4+b^4} = \frac{8ab^7}{a^8-b^8}. \quad (3)$

Hence, $\frac{a}{a-b} - \frac{a}{a+b} - \frac{2ab}{a^2+b^2} - \frac{4ab^3}{a^4+b^4} = \frac{8ab^7}{a^8-b^8}.$

$$62. \frac{a+b}{a-b} - \frac{a-b}{a+b} - \frac{4ab}{a^2+b^2} + \frac{8ab^3}{a^4+b^4}.$$

$$63. \frac{1}{a-b} - \frac{1}{a+b} - \frac{2b}{a^2+b^2} + \frac{2b^3}{a^4+b^4}.$$

$$64. \frac{1}{x-1} + \frac{1}{x+1} - \frac{x}{x^2+1} - \frac{x^3+3x}{x^4+1}.$$

$$65. \frac{a+x}{a-x} + \frac{a^2+x^2}{a^2-x^2} - \frac{a-x}{a+x} - \frac{a^2-x^2}{a^2+x^2} - \frac{4a^3x+4ax^3}{a^4-x^4}.$$

$$66. \text{Simplify } \frac{a}{(b-c)(c-a)} - \frac{b}{(a-c)(a-b)} + \frac{c}{(b-a)(c-b)}.$$

SOLUTION

$$\begin{aligned} & \frac{a}{(b-c)(c-a)} - \frac{b}{(a-c)(a-b)} + \frac{c}{(b-a)(c-b)} \\ \S 170, & = \frac{a}{(b-c)(c-a)} + \frac{b}{(c-a)(a-b)} + \frac{c}{(a-b)(b-c)} \\ & = \frac{a(a-b) + b(b-c) + c(c-a)}{(b-c)(c-a)(a-b)}. \end{aligned}$$

Simplify:

$$67. \frac{1}{(b-c)(a-c)} + \frac{1}{(c-a)(a-b)} + \frac{1}{(b-a)(b-c)}.$$

$$68. \frac{a+1}{(a-b)(a-c)} + \frac{b+1}{(b-c)(b-a)} + \frac{c+1}{(a-c)(b-c)}.$$

$$69. \frac{c^2ab}{(c-a)(b-c)} - \frac{b^2ca}{(b-a)(b-c)} - \frac{a^2bc}{(a-b)(a-c)}.$$

$$70. \frac{b-c}{(b-a)(a-c)} - \frac{c-a}{(b-c)(a-b)} - \frac{a+b}{(a-c)(b-c)}.$$

$$71. \frac{c+a}{(a-b)(b-c)} - \frac{b+c}{(c-a)(b-a)} + \frac{a+b}{(c-b)(a-c)}.$$

$$72. \frac{c+a}{(a+b)(b-c)} - \frac{c-b}{(c-a)(a+b)} - \frac{a-b}{(b-c)(a-c)}.$$

$$73. \frac{c+a-b}{(a-b)(b-c)} + \frac{b+c-a}{(c-a)(a-b)} - \frac{a+b+c}{(b-c)(a-c)}.$$

MULTIPLICATION OF FRACTIONS

179. 1. How much is 5 times $\frac{3}{4}$? $2 \times \frac{3}{4}$? $6 \times \frac{4a}{5}$? $3 \times \frac{2x}{7}$?
 $4 \times \frac{2m}{5}$? $3 \times \frac{3b}{4}$? $c \times \frac{a}{4}$? $a \times \frac{2b}{d}$?

2. Express $5 \times \frac{3}{10}$ in its lowest terms; $3 \times \frac{4}{9}$; $4 \times \frac{5a}{12}$; $2 \times \frac{7b}{6}$;
 $11 \times \frac{3x}{22}$; $7 \times \frac{a}{14}$; $10 \times \frac{5a}{20}$; $8 \times \frac{7ax}{16}$.

3. In what two ways, then, may a fraction be multiplied by an integer?

4. How much is $\frac{1}{3}$ of $\frac{10}{8}$, or $\frac{10}{8} \div 5$? $\frac{1}{4}$ of $\frac{8a}{9}$, or $\frac{8a}{9} \div 4$?

5. How much is $\frac{1}{5}$ of $\frac{7}{3}$, or $\frac{7}{3} \div 5$? $\frac{1}{4}$ of $\frac{5x}{9}$, or $\frac{5x}{9} \div 4$?

6. In what two ways, then, may a fraction be divided by an integer?

180. PRINCIPLES. — 1. *Multiplying the numerator or dividing the denominator of a fraction by any number multiplies the fraction by that number.*

2. *Dividing the numerator or multiplying the denominator of a fraction by any number divides the fraction by that number.*

EXAMPLES

1. Multiply $\frac{a}{b}$ by $\frac{c}{d}$.

PROCESS

EXPLANATION. — To multiply $\frac{a}{b}$ by $\frac{c}{d}$ is to find c times

$$\frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd}$$

the $\frac{1}{d}$ part of $\frac{a}{b}$. $\frac{1}{d}$ part of $\frac{a}{b} = \frac{a}{bd}$ (Prin. 2), and c times

$$\frac{a}{bd} = \frac{ac}{bd} \text{ (Prin. 1). Therefore, } \frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd}.$$

OR

To find the product of $\frac{a}{b} \times \frac{c}{d}$.

Let

$$x = \frac{a}{b} \times \frac{c}{d} \quad (1)$$

Multiplying each member of the equation by $b \times d$, Ax. 4,

$$x \times b \times d = \frac{a}{b} \times \frac{c}{d} \times b \times d$$

$$\S 104, 1, \quad = \frac{a}{b} \times b \times \frac{c}{d} \times d = ac.$$

$$xbd = ac.$$

$$\text{Dividing by } bd, \text{ Ax. 5,} \quad x = \frac{ac}{bd}. \quad (2)$$

$$\therefore \text{Eq. (1) and (2), Ax. 1,} \quad \frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd}.$$

$$\text{Similarly,} \quad \frac{a}{b} \times \frac{c}{d} \times \frac{e}{f} = \frac{ace}{bdf};$$

and so on for any number of fractions.

RULE. — *Multiply the numerators together for the numerator of the product, and the denominators for the denominator of the product.*

1. Cancel equal factors from the numerator and denominator.
2. Reduce integral and mixed expressions to the fractional form before multiplying.

$$2. \text{ Simplify } \frac{3x^2 + 15x}{x + 3} \times \frac{x^2 - 9}{x + 5} \times \frac{1}{2x^2 - 10x}.$$

$$\begin{aligned} \text{SOLUTION.} \quad & \frac{3x^2 + 15x}{x + 3} \times \frac{x^2 - 9}{x + 5} \times \frac{1}{2x^2 - 10x} \\ & = \frac{3x(x + 5)}{x + 3} \times \frac{(x + 3)(x - 3)}{x + 5} \times \frac{1}{2x(x - 5)} \end{aligned}$$

$$\text{Canceling,} \quad = \frac{3(x - 3)}{2(x - 5)}.$$

Multiply:

$$3. \quad \frac{3ab}{4xy} \text{ by } \frac{2y}{3a^2}.$$

$$6. \quad \frac{4mn}{3xy} \text{ by } -\frac{15bx}{16m^2}.$$

$$4. \quad \frac{5xy}{2ac} \text{ by } \frac{3ax}{10y^2}.$$

$$7. \quad \frac{2ax}{12by} \text{ by } -\frac{10b^2}{x^2}.$$

$$5. \quad \frac{4ab}{10c^2} \text{ by } \frac{3bc}{a^3}.$$

$$8. \quad \frac{a^mb^n}{4x} \text{ by } \frac{6x^2}{a^{m-1}b^{2n}}.$$

$$9. \frac{a^{m+1}}{b^{m+2}} \text{ by } \frac{b^{m+1}}{a^m}.$$

$$11. \frac{xy^2}{20-8x} \text{ by } \frac{25-10x}{x^2y}.$$

$$10. \frac{a}{a+b} \text{ by } \frac{b}{a-b}.$$

$$12. \frac{1-5x+6x^2}{2-3x+x^2} \text{ by } \frac{2-x}{1-x}.$$

Simplify the following:

$$13. \frac{(a-b)^2}{a+b} \times \frac{b}{a^2-ab} \times \frac{(a+b)^2}{a^2-b^2}.$$

$$14. \frac{a^4-x^4}{a^2+x^2} \times \frac{a+x}{a^2-x^2} \times \frac{a^2-ax+x^2}{(a+x)^2}.$$

$$15. \frac{4a-b}{2x+y} \times \frac{2a}{4a^2-ab} \times \frac{4x^2-y^2}{4}.$$

$$16. \frac{p+2}{x-3} \times \frac{3x^2-27}{2p^2-8} \times \frac{4}{px+3p}.$$

$$17. \frac{p^4-q^4}{(p-q)^2} \times \frac{p-q}{p^2+pq} \times \frac{p^2}{p^2+q^2}.$$

$$18. \frac{a^2+8}{a^2-8} \times \frac{a^2+2a+4}{a^2-2a+4}.$$

$$19. \frac{a^4+a^2x^2+x^4}{a^4-ax^3} \times \frac{x}{a^3-ax+x^2}.$$

$$20. \frac{a^4+4}{a^4+a^2+1} \times \frac{a^2+a+1}{a^2+2a+2}.$$

$$21. \frac{x^2+5x+6}{x^2+6x+5} \times \frac{x^2+7x+10}{x^2+7x+12}.$$

$$22. \frac{x^2+3x-10}{x^2-4x-21} \times \frac{x^2-10x+21}{x^2+7x+10}.$$

$$23. \frac{a^2+ab+ac+bc}{ax-ay-x^2+xy} \times \frac{a^2-ax+ay-xy}{a^2+ac+ax+cx} \times \frac{x^2-x(y-a)-ay}{a^2-a(y-b)-by}.$$

$$24. \frac{x^3-5x^2+8x-4}{x^3-8x^2+19x-12} \times \frac{x^3-10x^2+33x-36}{x^3-6x^2+11x-6}.$$

$$25. \frac{x^4-3x^3-23x^2+75x-50}{x^4-5x^3-21x^2+125x-100} \times \frac{x^3-10x^2+29x-20}{x^3-12x^2+45x-50}.$$

DIVISION OF FRACTIONS

- 181.** 1. How many times is $\frac{1}{4}$ contained in 1? $\frac{1}{5}$ in 1? $\frac{1}{6}$ in 1?
 2. If the numerator of a fraction is 1, how does the number of times the fraction is contained in 1 compare with the denominator?
 3. How many times is $\frac{1}{d}$ contained in 1? $\frac{1}{x+y}$ in 1?
 4. How many times is $\frac{2}{6}$ contained in 1? $\frac{2}{d}$ in 1? $\frac{c}{d}$ in 1?

182. The reciprocal of a fraction is the fraction inverted.

For, since $\frac{b}{a}$ multiplied by $\frac{a}{b}$ produces 1, § 102, if 1 is the dividend and $\frac{a}{b}$ the divisor, $\frac{b}{a}$ is the quotient.

EXAMPLES

1. Divide $\frac{a}{b}$ by $\frac{c}{d}$.

PROCESS **EXPLANATION.**—The fraction $\frac{1}{d}$ is contained in 1,
 $\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \times \frac{d}{c} = \frac{ad}{bc}$ d times; and $\frac{c}{d}$ is, therefore, contained in $1, \frac{1}{c}$ part
of d times, or $\frac{d}{c}$ times.

Since $\frac{c}{d}$ is contained $\frac{d}{c}$ times in 1, it will be contained $\frac{a}{b}$ times $\frac{d}{c}$, or $\frac{ad}{bc}$ times in $\frac{a}{b}$.

OR

To find the quotient of $\frac{a}{b} \div \frac{c}{d}$.

Let (1)

$$x = \frac{a}{b} \div \frac{c}{d}$$

§ 102, $x \times \frac{c}{d} = \frac{a}{b}$

Multiplying each member of this equation by $\frac{d}{c}$,

$$x \times \frac{c}{d} \times \frac{d}{c} = \frac{a}{b} \times \frac{d}{c}$$

That is, (2)

$$x = \frac{a}{b} \times \frac{d}{c}$$

Hence, eq. (1) and (2), Ax. 1, $\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \times \frac{d}{c}$

RULE. — *Multiply the dividend by the reciprocal of the divisor.*

1. Change integral and mixed expressions to the fractional form.
2. An integer may be expressed as a fraction by writing 1 for its denominator.
3. When possible, use cancellation.

Divide:

2. 1 by $\frac{ab}{x}$.

4. 1 by $\frac{a-b}{a+b}$.

3. 1 by $\frac{x^3}{x^2}$.

5. 1 by $\frac{a-3}{a+3}$.

Write the reciprocal of

6. $\frac{a}{b}$.

8. $\frac{m}{n}$.

10. $\frac{a-x}{b-y}$.

7. $\frac{3m}{p}$.

9. $\frac{1}{3m}$.

11. $\frac{4}{ab}$.

Simplify:

12. $\frac{5mn}{6bx} \div \frac{10m^2n}{3ax^2}$.

21. $\frac{x^2+x-2}{x^2-5x+4} \div \frac{x^2-x-6}{x^2+x-20}$.

13. $\frac{12a^4b}{25ac} \div \frac{4ax}{24c^2}$.

22. $\frac{a^4-b^4}{a^2-2ab+b^2} \div \frac{a^2+b^2}{a^2-ab}$.

14. $\frac{3abm}{7} \div abx$.

23. $\frac{x^2+y^2}{x^2-y^2} \div \frac{x^2+xy+y^2}{x-y}$.

15. $\frac{5ab}{3a^2c^2} \div \frac{25b^2}{15a^2}$.

24. $\frac{a^3+b^3}{a^2-4b^2} \div \frac{a^2-ab+b^2}{a-2b}$.

16. $\frac{7x^3}{4y^3} \div \frac{21x^2y^2}{14a}$.

25. $\frac{m^3-y^3}{m^2y^2-y^4} \div \frac{m^3+m^2y+my^2}{my^2+y^3}$.

17. $\frac{my-y^2}{(m+y)^2} \div \frac{y^2}{m^2-y^2}$.

26. $\frac{m^4x+m^5}{m^3x-mx^3} \div \frac{m^3x^2-mx^4}{m^3x^3+x^6}$.

18. $\frac{(a-b)^3}{a+b} \div \frac{a^2-ab}{b}$.

27. $\left(x + \frac{1}{y}\right) \div \left(y^2 + \frac{1}{x^2}\right)$.

19. $(4a+2) \div \frac{2a+1}{5a}$.

28. $\left(\frac{a^3}{b} \div b^2\right) \div \left(\frac{a^2}{b^2} \times ab\right)$.

20. $\frac{x^2-y^2}{x+2y} \div (x^2-3xy+2y^2)$.

29. $(a+c) \div \left(\frac{a^2-c^2}{1+x} + \frac{a-c}{1-x^2}\right)$.

$$30. \frac{x^3 - 6x^2 + 11x - 6}{x^3 + 2x^2 - 19x - 20} \div \frac{x^3 - 13x + 12}{x^3 + 10x^2 + 29x + 20}.$$

$$31. \frac{x^3 - 15x^2 + 74x - 120}{x^3 - 5x^2 - x + 5} \div \frac{x^3 - 9x^2 + 26x - 24}{x^3 - 6x^2 + 11x - 6}.$$

$$32. \frac{a^2 + b^2 - c^2 + 2ab}{a^2 - b^2 - c^2 + 2bc} \div \frac{a^2 - b^2 + c^2 - 2ac}{a^2 - b^2 + c^2 + 2ac}.$$

$$33. \frac{a^2 + x^2 - y^2 + 2ax}{a^2 - x^2 + y^2 - 2ay} \div \frac{a^2 - x^2 + y^2 + 2ay}{a^2 - x^2 - y^2 - 2xy}.$$

$$34. \text{Simplify } \left(y - x + \frac{x^2}{y}\right) \div \left(\frac{x}{y^2} + \frac{y}{x^2}\right).$$

SOLUTION

$$\begin{aligned} & \left(y - x + \frac{x^2}{y}\right) \div \left(\frac{x}{y^2} + \frac{y}{x^2}\right) \\ &= \frac{y^2 - xy + x^2}{y} \div \frac{x^2 + y^2}{x^2 y^2} \\ &= \frac{x^2 - xy + y^2}{y} \times \frac{x^2 y^2}{(x + y)(x^2 - xy + y^2)} \\ &= \frac{x^2 y}{x + y}. \end{aligned}$$

Simplify:

$$35. \left(x - 4 + \frac{9}{x + 2}\right) \div \left(1 - \frac{4x - 7}{x^2 - 4}\right).$$

$$36. \left(x + \frac{3x + 6}{x^2 - 1} + 2\right) \div \left(x + 3 + \frac{1}{x + 1}\right).$$

$$37. \left(x^3 - \frac{1}{x^3}\right) \div \left(x - \frac{1}{x}\right).$$

$$38. \left(1 + \frac{1}{y^2} + \frac{1}{y^4}\right) \div \left(1 + \frac{1}{y} + \frac{1}{y^3}\right).$$

$$39. \left(1 - \frac{y^2}{x^2}\right) \div \left(1 - \frac{2x}{y} + \frac{x^2}{y^2}\right).$$

$$40. \left(1 - \frac{2y^3}{x^3 + y^3}\right) \div \left(1 - \frac{2y}{x + y}\right).$$

COMPLEX FRACTIONS

183. A fraction one or both of whose terms contains a fraction is called a **Complex Fraction**.

It is simply an expression of unexecuted division.

EXAMPLES

1. Simplify the expression $\frac{\frac{a}{b}}{\frac{x}{y}}$.

PROCESS

$$\frac{\frac{a}{b}}{\frac{x}{y}} = \frac{a}{b} \div \frac{x}{y} = \frac{a}{b} \times \frac{y}{x} = \frac{ay}{bx}$$

Simplify the following expressions:

2. $\frac{\frac{x+y}{ab}}{\frac{x^2-y^2}{ab^2}}$

5. $\frac{2 + \frac{3a}{4b}}{a + \frac{8b}{3}}$

8. $\frac{b - \frac{c}{2}}{\frac{b}{2} - c}$

3. $\frac{a + \frac{b}{c}}{b + \frac{c}{a}}$

6. $\frac{1 - \frac{y^2}{x^2}}{1 + \frac{y^2}{x^2}}$

9. $\frac{ax - \frac{x^2}{2}}{\frac{a^2}{2} - ax}$

4. $\frac{m - \frac{3m}{x}}{x - \frac{x}{m}}$

7. $\frac{x - \frac{1}{x}}{1 + \frac{1}{x}}$

10. $\frac{\frac{x+y}{y} - \frac{x+y}{x}}{\frac{1}{y} - \frac{1}{x}}$

11. Simplify the expression $\frac{\frac{x^3}{y^2} - \frac{x}{y} + 1}{\frac{x^2}{y^2} + \frac{x}{y} + 1}$.

SOLUTION. — Multiplying the numerator and denominator of the fraction by y^2 , the L. C. D. of the fractional parts of the numerator and denominator, the expression becomes $\frac{x^2 - xy + 1}{x^2 + xy + 1}$.

Simplify the following:

$$12. \frac{\frac{x^2-1}{x}}{\frac{x+1}{x^2}}$$

$$14. \frac{\frac{x^3+y^3}{xy}}{\frac{x^2-xy+y^2}{xy}}$$

$$16. \frac{\frac{x^2+y^2}{2y}-x}{\frac{x}{y}-\frac{y}{x}}$$

$$13. \frac{\frac{1}{x}+\frac{1}{y+z}}{\frac{1}{x}-\frac{1}{y+z}}$$

$$15. \frac{\frac{1}{a+1}}{1-\frac{1}{a+1}}$$

$$17. \frac{\frac{1}{1-a}}{\frac{1}{1+a}}$$

$$18. \frac{x-2+\frac{1}{x+2}}{x+2+\frac{1}{x-2}}$$

$$20. \frac{6a-1-\frac{1}{a}}{\frac{2a-1}{3a}}$$

$$19. \frac{\frac{1}{x}+\frac{4}{x^2}+\frac{4}{x^3}}{1+\frac{5}{x}+\frac{6}{x^2}}$$

$$21. \frac{\frac{x-5}{2}-7+\frac{24}{x}}{\frac{3x-9}{x}}$$

$$22. \frac{\frac{1}{x+1}}{1-\frac{1}{1+x}} + \frac{\frac{1}{x+1}}{\frac{x}{1-x}} + \frac{\frac{1}{1-x}}{\frac{x}{1+x}}$$

$$23. \frac{3xyz}{yz+zx+xy} - \frac{\frac{x-1}{x} + \frac{y-1}{y} + \frac{z-1}{z}}{\frac{1}{x} + \frac{1}{y} + \frac{1}{z}}$$

$$24. \frac{\frac{1}{x+y} + \frac{2}{x-y} - \frac{9}{3x-y}}{\frac{-8y}{y^2-9x^2}}$$

$$25. \frac{\frac{x^2+(a+b)x+ab}{x^2-(a+b)x+ab}}{\frac{x^2-b^2}{x^2-a^2}}$$

$$26. \frac{\frac{1}{a} + \frac{1}{b+c}}{\frac{1}{a} - \frac{1}{b+c}} \div \frac{1}{1 + \frac{b^2+c^2-a^2}{2bc}}$$

$$27. \frac{\frac{x+y}{x-y} + \frac{x^2+y^2}{x^2-y^2}}{\frac{x-y}{x+y} - \frac{x^3-y^3}{x^3+y^3}} \times \frac{(x-y)^2}{x^4+x^2y^2+y^4}$$

$$28. \frac{\frac{a-b}{1+ab} + \frac{b-c}{1+bc}}{1 - \frac{(a-b)(b-c)}{(1+ab)(1+bc)}} \div \frac{1-\frac{c}{a}}{\frac{1}{a}+c}$$

$$29. \frac{x^2 - \frac{(x^2+y^2-z^2)^2}{4y^2}}{\frac{(x+y)^2-z^2}{y^2}} \times \frac{(x-y+z)^2}{4}$$

184. An expression of the form $\frac{a}{b + \frac{c}{d + \frac{e}{f + \dots}}}$ is called a Continued Fraction.

30. Simplify $\frac{1}{1 + \frac{1}{1 + \frac{1}{x}}}$.

SOLUTION

$$\begin{aligned} \frac{1}{1 + \frac{1}{1 + \frac{1}{x}}} &= \frac{1}{1 + \frac{1}{\frac{x+1}{x}}} \\ &= \frac{1}{1 + \frac{x}{x+1}} \\ &= \frac{x+1}{x+1+x} \\ &= \frac{x+1}{2x+1} \end{aligned}$$

SUGGESTION.—In the above example, the part first simplified is the *last complex part* $\frac{1}{1 + \frac{1}{x}}$, which is reduced to a simple fraction.

Every continued fraction may be simplified by successively reducing its *last complex part* to a simple fraction.

Simplify the following:

$$31. \frac{1}{x + \frac{1}{1 + \frac{x+1}{3-x}}}$$

$$34. \frac{x-2}{x-2 - \frac{x}{x - \frac{x-1}{x-2}}}$$

$$32. \frac{a}{a+1 + \frac{a}{a+1 - \frac{1}{a}}}$$

$$35. \frac{1}{a + \frac{1}{a + \frac{1}{a}}}$$

$$33. \frac{2}{2 - \frac{2}{2 - \frac{2}{2-x}}}$$

$$36. 1 + \frac{c}{1+c + \frac{2c}{1+\frac{1}{c}}}$$

REVIEW OF FRACTIONS

185. Reduce to their lowest terms:

$$1. \frac{x^3 + x^2 + x - 3}{x^3 + 3x^2 + 5x + 3}$$

$$5. \frac{x^3 + x^2 - 22x - 40}{x^3 - 7x^2 + 2x + 40}$$

$$2. \frac{x^3 - x^2 - x - 2}{x^3 + 3x^2 + 3x + 2}$$

$$6. \frac{x^3 + 10x^2 + 7x - 18}{x^3 - 8x^2 - 11x + 18}$$

$$3. \frac{x^3 + 4x^2 + 8x + 5}{x^3 + 3x^2 + 7x + 5}$$

$$7. \frac{2x^3 + 7x^2 - 9x - 9}{2x^3 + 9x^2 + x - 3}$$

$$4. \frac{x^3 + 3x^2 + 4x + 2}{x^3 - 3x^2 - 8x - 10}$$

$$8. \frac{8x^3 - 22x^2 + 17x - 3}{6x^3 - 17x^2 + 14x - 3}$$

Simplify:

$$9. \frac{x}{2y-1} + \frac{y}{2y+1} - \frac{y-x}{1-4y^2}$$

$$10. \frac{a^2}{4(1-a)^2} - \left(\frac{3}{8(1-a)} + \frac{1}{8(a+1)} - \frac{1-a}{4(a+1)} \right)$$

$$11. \frac{2x-1}{x-1} - \frac{3x+1}{x+1} + \frac{3x-1}{x-2} - \frac{2x+1}{x+2}$$

$$12. \frac{a^3}{(a-b)(a-c)} + \frac{b^3}{(b-c)(b-a)} + \frac{c^3}{(c-a)(c-b)}$$

13. $\left(1 + \frac{2}{m-1}\right)\left(\frac{m^2 + m - 2}{m^2 + m}\right).$
14. $\left(1 - a + a^2\right)\left(\frac{1}{a^2} + \frac{1}{a} + 1\right).$
15. $\left(\frac{x^2}{y^2} + \frac{x}{y} + 1\right)\left(\frac{x^2}{y^2} - \frac{x}{y} + 1\right).$
16. $\left(\frac{a+b}{a-b} - \frac{a-b}{a+b} - \frac{4ab}{a^2+b^2}\right)\frac{a^2-b^2}{8b^3}.$
17. $\left(1 + \frac{x}{y}\right)\left(\frac{x}{y^2} - \frac{1}{y} + \frac{1}{x}\right).$
18. $\left(x + 1 + \frac{1}{x} + \frac{1}{x^2}\right) + \left(x + 1 - \frac{1}{x} - \frac{1}{x^2}\right).$
19. $\left(\frac{a+b}{a-b} + \frac{a^2+b^2}{a^2-b^2}\right)\left(\frac{a+b}{a-b} - \frac{a^3+b^3}{a^3-b^3}\right).$
20. $\left(1 - \frac{2xy}{x^2+y^2}\right) + \left(\frac{x^3-y^3}{x-y} - 3xy\right).$
21. $(x^2 - y^2 - z^2 - 2yz) \div \frac{x+y+z}{x+y-z}.$
22. $\left(\frac{1+a}{1-a} + \frac{4a}{1+a^2} + \frac{8a}{1-a^2} - \frac{1-a}{1+a}\right) + \left(\frac{1+a^2}{1-a^2} + \frac{4a^2}{1+a^2} - \frac{1-a^2}{1+a^2}\right).$
23. $\frac{\frac{a^2x^2}{bd} + \frac{abx^2}{c^2d} - \frac{acx^2}{d^2} - \frac{b^2x}{cd^2} + \frac{a^2x}{bc} - \frac{a}{d}}{\frac{ax}{c} - \frac{b}{d}}.$
24. $\left(x^2 - 3xy - 2y^2 + \frac{12y^3}{x+3y}\right) \div \left(3x - 6y - \frac{2x^2}{x+3y}\right).$
25. $\left(\frac{m-3n}{m+n}\right)\left(1 + \frac{4n}{m+n}\right) \div \left(\frac{m}{n} + 2 - \frac{15n}{m}\right).$
26. $1 - \frac{2x+5x^2}{2(x+1)^2} - \left[\frac{x^2+x}{2x-\frac{2}{x}}\right]\left(\frac{3-3x}{(x+1)^2}\right).$
27. $\left(1 + \frac{x}{a-x}\right)\left(\frac{x}{x+a} - \frac{2x^2+2ax-a^2}{x^2+3ax+2a^2}\right).$

Expand:

28. $\left(\frac{x}{y} + \frac{y}{x}\right)\left(\frac{x}{y} - \frac{y}{x}\right).$

31. $\left(\frac{a}{b} + 1 + \frac{b}{a}\right)\left(\frac{a}{b} - 1 + \frac{b}{a}\right).$

29. $\left(\frac{x}{y} + \frac{y}{x}\right)\left(\frac{x}{y} + \frac{y}{x}\right).$

32. $\left(x^2y + xy^2\right)\left(\frac{1}{x^3} - \frac{2y}{x^4} + \frac{y^2}{x^5}\right).$

30. $\left(2x - \frac{1}{2x}\right)\left(2x - \frac{1}{2x}\right).$

33. $\left(x^2 + x + 1\right)\left(1 - \frac{1}{x} + \frac{1}{x^2}\right).$

Simplify:

34. $\frac{1 + \frac{1}{x^2} + \frac{1}{x^4}}{1 + \frac{1}{x} + \frac{1}{x^2}}.$

38. $\frac{\frac{6a}{x+y} - \frac{4}{x+y}}{\frac{1}{(x+y)^2}}.$

35. $\frac{\left(\frac{m^2+n^2}{n} - m\right) + \left(\frac{1}{n} - \frac{1}{m}\right)}{\frac{m^3+n^3}{m^2-n^2}}.$

39. $1 - \frac{\frac{a^3+3a+2}{a^2+2a+1}}{\frac{a^2+7a+12}{a^2+5a+4}}.$

36. $\frac{\frac{c}{(a+1)^2} + \frac{d}{(a+1)^2}}{\frac{a}{(a+1)^4} + \frac{1}{(a+1)^4}}.$

40. $\frac{\frac{m^3-n^3}{m^3+n^3}\left(1 - \frac{2n}{m+n}\right)}{1 + \frac{2mn}{m^2-mn+n^2}}.$

37. $\frac{1}{1 - \frac{1}{1 - \frac{1}{1-x}}}.$

41. $\frac{1}{2 - \frac{3}{4 - \frac{5}{6-x}}}.$

42. $\left(\frac{x+y}{x-y} + \frac{x-y}{x+y}\right) + \left(\frac{x+y}{2(x-y)} - \frac{x-y}{2x+2y}\right).$

43. $\frac{\left(\frac{a}{x^2} + \frac{1}{x} + \frac{1}{a} + \frac{x}{a^2}\right)\left(\frac{a^2}{x^3} - \frac{1}{x} + \frac{x}{a^2}\right)}{\frac{a^3}{x^5}\left(1 + \frac{x}{a}\right)}.$

44. $\frac{x^3 - \frac{8}{y^3}}{x^3y^3 - x^2y^3} \times \frac{\frac{1}{xy} + \frac{1}{x^2y^3}}{1 + \frac{2}{xy} + \frac{4}{x^2y^3}} \times \frac{xy-1}{xy+1}.$

SIMPLE EQUATIONS

ONE UNKNOWN NUMBER

186. An *Equation* has been defined, § 2, as an expression of equality between two numbers or quantities.

187. An equation all of whose known numbers are expressed by figures is called a **Numerical Equation**.

188. An equation one or more of whose known numbers is expressed by letters is called a **Literal Equation**.

189. An equation that does not involve an unknown number in any denominator is called an **Integral Equation**.

$x + 5 = 8$ and $\frac{2x}{3} + 5 = 8$ are integral equations. The second equation is integral; for though it contains a fraction, the unknown number x does not appear in the denominator.

190. An equation that involves an unknown number in any denominator is called a **Fractional Equation**.

$x + 5 = \frac{8}{x}$ and $\frac{2x}{x-1} = 7$ are fractional equations.

191. An equation whose members are identical, or such that they may be reduced to the same form, is called an **Identical Equation**, or an **Identity**.

$a + b = a + b$; $a^2 - b^2 = (a + b)(a - b)$ are identical equations.

An equation whose members are numerical is evidently an identical equation.

$10 = 6 + 4$; $8 \times 2 = 6 + 12 - 2$ are identical equations.

A literal equation that is true for all values of the letters involved is an identical equation, or an identity.

$(x + y)^2 = x^2 + 2xy + y^2$ is an identity, because it is true for all values of x and y .

192. An equation that is true for only certain values of its letters is called an **Equation of Condition**.

An equation of condition is usually termed simply an *Equation*.

$x + 4 = 10$ is an equation of condition, because it is true only when the value of x is 6. $x^2 = 9$ is an equation of condition, because it is true only when the value of x is $+3$ or -3 .

193. When an equation is reduced to an identity by the substitution of certain numbers for the unknown numbers, the equation is said to be *satisfied*.

When $x = 2$, the equation $3x + 4 = 10$ becomes $6 + 4 = 10$, an identity; consequently, the equation is satisfied.

Any number that satisfies an equation is called a **Root** of the equation.

2 is a root of the equation $3x + 4 = 10$.

Finding the roots of an equation is called **solving the equation**.

194. An integral equation that involves only the first power of one unknown number in any term when the similar terms have been united is called a **Simple Equation**, or an **Equation of the First Degree**.

$3x + 4 = 10$ and $x + 2y - z + 8$ are simple equations.

195. Two equations that have the same roots, each equation having all the roots of the other, are called **Equivalent Equations**.

By the axioms in § 74, if the members of an equation are equally increased or diminished or are multiplied or divided by the same or equal numbers, the resulting numbers are *equal* and, § 186, form an equation. But it does not necessarily follow that the equation so formed is *equivalent* to the given equation.

For example, if both members of the equation $x + 2 = 5$, whose only root is $x = 3$, are multiplied by $x - 1$, the resulting numbers $(x + 2)(x - 1)$ and $5(x - 1)$ are *equal* and form an equation $(x + 2)(x - 1) = 5(x - 1)$; but this equation is not *equivalent* to the given equation, since it is satisfied by $x = 1$ as well as by $x = 3$.

196. PRINCIPLES.—1. *If the same expression is added to or subtracted from both members of an equation, the resulting equation is equivalent to the given equation.*

2. *If both members of an equation are multiplied or divided by the same known number, except zero, the resulting equation is equivalent to the given equation.*

3. *If both members of an integral equation are multiplied by the same unknown integral expression, the resulting equation has all the roots of the given equation and also the roots of the equation formed by placing the multiplier equal to zero.*

It follows from Prin. 3 that if the same unknown factor is removed from both members of an equation, the resulting equation has all the roots of the given equation except those obtained from the equation formed by placing the factor removed equal to zero.

Principle 1 may be established as follows:

$$\text{Let} \qquad A = B \qquad (1)$$

be any equation and C any expression to be added or subtracted.

$$\text{It is to be proved that} \quad A \pm C = B \pm C \qquad (2)$$

is equivalent to (1), the given equation.

All the values of the unknown number or numbers that satisfy (1), that is, make A identical with B , make $A + C$ identical with $B + C$ and $A - C$ identical with $B - C$, that is, satisfy (2). Hence, (2) has all the roots of (1).

$$\text{For the same reason} \quad A \pm C \mp C = B \pm C \mp C \qquad (3)$$

has all the roots of (2). But by the Associative Law for addition (3) may be written $A + 0 = B + 0$, or $A = B$. Hence, (1) has all the roots of (2).

Since (2) has all the roots of (1) and (1) has all the roots of (2), (2) is equivalent to (1), the given equation.

Principles 2 and 3 may be established as follows:

$$\text{Let} \qquad A = B \qquad (1)$$

$$\text{and} \qquad MA = MB. \qquad (2)$$

$$\text{From (1), by Prin. 1,} \quad A - B = 0. \qquad (3)$$

$$\text{From (2), by Prin. 1,} \quad M(A - B) = 0. \qquad (4)$$

Since the first member of (4) can reduce to zero only when one or both of its factors become 0, (4) is satisfied by those values of the unknown number that make $A - B = 0$, that is, by the roots of (3), or, Prin. 1, of (1); and also by those values of the unknown number that make $M = 0$, that is, by the roots of $M = 0$, but by no other values.

If M is any known number, not zero, M cannot be placed equal to zero and then (4), or (2), is equivalent to (3), or to (1).

If M is an unknown expression, (4), or (2), has the roots of $M = 0$ in addition to the roots of $A - B = 0$, or of (1).

197. By § 196, Prin. 1 and 2, every simple equation involving one unknown number may be reduced to an equivalent equation having the form $x = a$, a being a fixed known number. Hence,

Every simple equation involving one unknown number has one root and only one; also, by § 196, Prin. 3,

The equation $(x - a)(x - b)(x - c) \cdots (x - r) = 0$ is equivalent to the simple equations $x - a = 0$, $x - b = 0$, $x - c = 0$, $\cdots x - r = 0$, and has as many roots as factors of the first degree involving x .

CLEARING EQUATIONS OF FRACTIONS

198. 1. If one third of a number is 10, what is the number?

2. If $\frac{1}{4}x = 7$, what is the value of x ? If $\frac{1}{5}x = 5$, what is the value of x ? If $\frac{1}{8}x = 6$, what is the value of x ?

3. If $\frac{2}{3}x = 6$, what is the value of $2x$? If $\frac{5}{7}x = 10$, what is the value of $5x$?

4. If $\frac{x}{3} = 5$, what is the form of the resulting equation when both members are multiplied by 3? by 6? by 9? by any multiple of the denominator?

199. The process of changing an equation containing fractions to an equation without fractions is called **Clearing the Equation of Fractions**.

200. PRINCIPLES.—1. *An equation may be cleared of fractions by multiplying both members by some common multiple of the denominators of the fractions.* (Ax. 4.)

2. *If both members of a fractional equation are multiplied by the expression of lowest degree required to clear the equation of fractions, the resulting equation is equivalent to the given equation.*

Principle 2 may be established as follows:

By § 196, Prin. 1, all the terms of the second member may be transposed to the first member. Hence, uniting the terms of the first member into one and reducing this to its lowest terms, any fractional equation may be reduced to an equivalent equation of the form

$$\frac{A}{B} = 0, \quad (1)$$

in which A is prime to B , and B is the expression of lowest degree required to clear the equation of fractions.

Since A and B have no common factors, A and B cannot reduce to zero at the same time for any value of the unknown number. Hence, eq. (1) is satisfied by every value of the unknown number that makes $A = 0$, and by no other values; that is, the equation $A = 0$, obtained by multiplying both members of the given equation by the expression of lowest degree required to clear it of fractions, is equivalent to the given equation.

EXAMPLES

1. Given $\frac{x-4}{4} = 6 - \frac{x}{3}$, to find the value of x .

PROCESS

$$\frac{x-4}{4} = 6 - \frac{x}{3}$$

Clearing of fractions, $3x - 12 = 72 - 4x$

$$\therefore x = 12$$

EXPLANATION. — Since the first fraction will become an integer if the members of the equation are multiplied by 4 or any multiple of 4, and since the second fraction will become an integer if the members of the equation are multiplied by 3 or any multiple of 3, the equation may be cleared of fractions in a single operation by multiplying its members by some *common* multiple of 4 and 3 (§ 196, Prin. 2, § 200, Prin. 1).

Since the terms derived from the numerators will be least or lowest when the multiplier is the least or lowest common multiple of the denominators, the members of the equation should be multiplied by the least common multiple of 4 and 3, which is 12.

The resulting equation is $3x - 12 = 72 - 4x$; $\therefore x = 12$.

2. Find the value of x in $\frac{x-1}{2} - \frac{x-2}{3} = \frac{2}{3} - \frac{x-3}{4}$.

SOLUTION.
$$\frac{x-1}{2} - \frac{x-2}{3} = \frac{2}{3} - \frac{x-3}{4}$$

Clearing of fractions, $6(x-1) - 4(x-2) = 8 - 3(x-3)$.

$$6x - 6 - 4x + 8 = 8 - 3x + 9.$$

$$\therefore x = 3.$$

RULE. — *Multiply both members of the equation by the least or lowest common multiple of the denominators.*

1. The multiplier of lowest degree required to clear an equation of fractions so that the resulting equation is equivalent to the given equation is usually the L. C. D. But if fractions having like denominators have not been united and every fraction reduced to its lowest terms, the multiplier required may be of lower degree than the L. C. D.

Thus, given
$$\frac{2x^2}{x-2} = \frac{3x+2}{x-2} + \frac{5x+8}{3}.$$

Uniting terms, $\frac{2x^2 - 3x - 2}{x-2}$, or $2x + 1 = \frac{5x+8}{3}.$

Multiplying by 3,
$$6x + 3 = 5x + 8.$$

$$\therefore x = 5.$$

Had the given equation been cleared of fractions by multiplying by $3(x-2)$ instead of by 3, the resulting equation simplified, which is $x^2 - 7x + 10 = 0$, or $(x-5)(x-2) = 0$, would have been satisfied both by $x = 5$ and by $x = 2$. The given equation is satisfied by $x = 5$ but not by $x = 2$. Hence, the latter value is not a root of the given equation, but has been introduced by using $(x-2)$ times the necessary multiplier, 3. Roots so introduced may be discovered by verification and rejected.

2. If a fraction has the minus sign before it, the signs of all the terms of the numerator must be changed when the denominator is removed.

Find the value of x , and verify the result:

3. $2x + \frac{x}{3} = \frac{35}{3}.$

7. $\frac{x}{2} + \frac{x}{6} = \frac{10}{3}.$

4. $\frac{x}{4} + 10 = 13.$

8. $7\frac{1}{2} - \frac{3x}{14} = \frac{x}{7}.$

5. $\frac{x}{6} + 2x = 26.$

9. $\frac{x}{5} + \frac{x}{7} = 24.$

6. $3x - \frac{x}{5} = 14.$

10. $\frac{2x}{3} - \frac{5}{6} = \frac{x}{4}.$

11. $\frac{x}{2} + \frac{x}{3} - \frac{x}{4} + \frac{3x}{10} - \frac{5x}{12} = 7.$

12. $\frac{25x}{18} - \frac{5x}{9} + \frac{2x}{3} - \frac{5x}{6} = 2.$

13. $\frac{2x}{7} - \frac{x}{4} - \frac{x}{2} + \frac{4x}{7} = \frac{3}{14}.$

$$14. \frac{2x}{3} - \frac{7x}{8} + \frac{5x}{18} + \frac{x}{24} = \frac{4}{9}.$$

$$15. \frac{3x}{4} + \frac{7x}{16} - \frac{x}{2} - \frac{9x}{16} = \frac{1}{8}.$$

$$16. \frac{15x}{7} + \frac{5x}{6} - \frac{11x}{3} + \frac{19x}{14} = 2.$$

$$17. \frac{2x}{15} + \frac{5x}{25} - \frac{4x}{9} + \frac{x}{6} = \frac{1}{9}.$$

$$18. \frac{3x}{4} - \frac{7x}{12} = \frac{11x}{36} - \frac{8x}{9} + \frac{3}{2}.$$

$$19. \frac{3x}{4} + \frac{1}{20} = \frac{3x}{10} + \frac{x}{5} + \frac{4x}{15}.$$

$$20. \frac{7x+1}{5} - 2x + \frac{4x+7}{7} = 1.$$

$$21. \frac{x-1}{2} + \frac{x-2}{3} + \frac{x-3}{4} = \frac{5x-1}{6}.$$

$$22. \frac{x+1}{3} - \frac{x+4}{5} + \frac{x+3}{4} = 16.$$

$$23. \frac{7x+2}{6} - \frac{12-x}{4} + \frac{x+2}{2} = 6.$$

$$24. \frac{x-3}{7} + \frac{x+5}{3} - \frac{x+2}{6} = 4.$$

$$25. \frac{3x-5}{4} - \frac{7x-13}{6} = 3 - \frac{x+3}{2}.$$

$$26. \frac{2x-5}{5} - \frac{3x-2}{7} = 1 - \frac{x+2}{6}.$$

$$27. \frac{1-2x}{3} - \frac{7-2x}{4} + \frac{11-2x}{6} = -\frac{7}{12}.$$

$$28. \frac{x+4}{3} + \frac{2-2x}{6} = \frac{x+1}{2} - 3\frac{1}{8}.$$

$$29. \frac{9x+5}{14} + \frac{8x-7}{6x+2} = \frac{36x+15}{56} + \frac{10\frac{1}{2}}{14}.$$

SUGGESTION. — The equation may be written

$$\S 104, 3, \quad \frac{9x}{14} + \frac{5}{14} + \frac{8x-7}{6x+2} = \frac{36x}{56} + \frac{15}{56} + \frac{41}{56}.$$

Simplify as much as possible before clearing of fractions.

$$30. \frac{3x-2}{2x-5} + \frac{3x-21}{5} = \frac{6x-22}{10}.$$

$$31. \frac{4x+3}{9} = \frac{8x+19}{18} - \frac{7x-29}{5x-12}.$$

$$32. \frac{6x+1}{15} - \frac{2x-4}{7x-13} = \frac{2x-1}{5}.$$

$$33. \frac{10x+17}{18} - \frac{5x-2}{9} = \frac{12x-1}{11x-8}.$$

$$34. \frac{6x+3}{15} - \frac{3x-1}{5x-25} = \frac{2x-9}{5}.$$

$$35. \frac{2x+1}{2x-1} - \frac{8}{4x^2-1} = \frac{2x-1}{2x+1}.$$

$$36. \text{ Solve the equation } \frac{x-1}{x-2} + \frac{x-6}{x-7} = \frac{x-5}{x-6} + \frac{x-2}{x-3}.$$

SOLUTION. — It will be observed that if the fractions in each member were connected by the sign $-$, and the members were simplified, the numerators of the resulting fractions would be simple. The fractions can be made to meet this condition by transposing one fraction in each member.

Consequently, it is sometimes expedient to defer clearing of fractions.

$$\text{Transposing,} \quad \frac{x-1}{x-2} - \frac{x-2}{x-3} = \frac{x-5}{x-6} - \frac{x-6}{x-7}.$$

$$\text{Uniting terms,} \quad \frac{-1}{x^2-5x+6} = \frac{-1}{x^2-13x+42}.$$

Since the fractions are equal and their numerators are equal, their denominators must be equal.

$$x^2-5x+6 = x^2-13x+42.$$

$$\therefore x = 4\frac{1}{2}.$$

$$37. \frac{x-1}{x-2} + \frac{x-7}{x-8} = \frac{x-5}{x-6} + \frac{x-3}{x-4}.$$

$$38. \frac{x-3}{x-4} + \frac{x-7}{x-8} = \frac{x-6}{x-7} + \frac{x-4}{x-5}.$$

$$39. \frac{x+2}{x+1} - \frac{x+3}{x+2} = \frac{x+5}{x+4} - \frac{x+6}{x+5}.$$

$$40. \frac{x+1}{x+2} + \frac{x+6}{x+7} = \frac{x+2}{x+3} + \frac{x+5}{x+6}.$$

$$41. \frac{x-5}{x+5} - \frac{x-10}{x+10} = \frac{x-4}{x+4} - \frac{x-9}{x+9}.$$

$$42. \frac{\frac{7x-2}{3} - \frac{5x-\frac{1}{2}}{3}}{4} = \frac{\frac{x-2}{4}}{5} - 3\frac{1}{8}.$$

$$43. \frac{x-1}{x-2} + \frac{x-2}{x-3} + \frac{x-6}{x-4} - 3 = 0.$$

$$44. \frac{x^3+1}{x-1} - \frac{x^3-1}{x+1} = \frac{8}{x^2-1} + 2x.$$

$$45. \frac{x^3+2}{x+1} - \frac{x^3-2}{x-1} = \frac{10}{x^2-1} - 2x.$$

$$46. \frac{\frac{2x+4}{3}}{2} = \frac{7\frac{1}{2}-x}{3} + \frac{x(\frac{6}{x}-1)}{2}.$$

$$47. \frac{\frac{4}{5x}-16}{24} - \frac{\frac{2}{5x}+6}{60} = \frac{4\frac{1}{5}}{5}.$$

$$48. \frac{x}{2}(2-x) - \frac{x}{4}(3-2x) = \frac{x+10}{6}.$$

$$49. \frac{2x\left(1-\frac{5}{x}\right)}{3} + \frac{3x\left(1-\frac{4}{x}\right)}{4} = \frac{x-4}{\frac{4}{5}}.$$

$$50. \frac{1}{2}x - 2\left(\frac{4x}{5} - 3\right) = 4 - \frac{3}{2}\left(\frac{x}{2} + 1\right).$$

$$51. \frac{(2x+1)^2}{5} - \frac{(4x-1)^2}{20} = \frac{15}{8} + \frac{3(4x+1)}{40}.$$

$$52. \frac{1}{2} \cdot \frac{x+1}{x-1} + 2 = \frac{1}{2} + \frac{x-1}{x+1} - \frac{x^2}{1-x^2}.$$

$$53. \frac{17 + \frac{3}{x}}{3} + \frac{1 + \frac{18}{x}}{5} = \frac{\frac{21}{x} - 1}{9} + \frac{\frac{100}{x} + \frac{5}{3}}{15}.$$

$$54. \frac{\frac{1}{2}(x-4)}{\frac{3}{2}} - \frac{4x-16}{6} = \frac{3}{5} - \frac{\frac{2x}{5} + 5}{\frac{5}{2}}.$$

$$55. 1 + \frac{1}{1 + \frac{1}{x}} = \frac{2 + \frac{10}{x}}{1 + \frac{6}{x}}.$$

$$56. \frac{1}{\frac{x-1}{\frac{x}{3} + \frac{3}{3x-1}}} = \frac{3}{x+5}.$$

LITERAL EQUATIONS

1. Solve the equation $\frac{x-b^2}{a} = \frac{x-a^2}{b}.$

SOLUTION

$$\frac{x-b^2}{a} = \frac{x-a^2}{b}.$$

Clearing of fractions, $bx - b^3 = ax - a^3.$

Transposing, etc., $ax - bx = a^3 - b^3.$

$$(a-b)x = a^3 - b^3.$$

Dividing by $(a-b),$ $x = a^2 + ab + b^2.$

Solve the following equations:

$$2. \frac{c^2 - x}{nx} + \frac{n^2}{cx} = \frac{1}{c}.$$

$$6. \frac{x-2ab}{cx} - \frac{1}{x} = \frac{x-3c}{abx}.$$

$$3. 1 - \frac{ab}{x} = \frac{7}{ab} - \frac{49}{abx}.$$

$$7. \frac{x-a}{b} + \frac{2x}{a} = 5 + \frac{6b}{a}.$$

$$4. \frac{a^3}{ab^2} - \frac{2a^2}{b^2x} = 1 - \frac{2b^3}{a^2x}.$$

$$8. \frac{a^2}{bx} + \frac{b^2}{ax} = \frac{a+b}{ab} - \frac{3(a+b)}{x}.$$

$$5. \frac{x}{b} - \frac{x+2b}{a} = \frac{a}{b} - 3.$$

$$9. \frac{a^2+b^2}{2bx} - \frac{a-b}{2bx^2} = \frac{b}{x}.$$

$$10. a + \frac{b(x+a)}{x-a} = \frac{2ab}{x-a}. \quad 11. \frac{x-2a}{a} + \frac{x}{b} = \frac{a^2+b^2}{ab}.$$

$$12. 6x + 18\left(1 - \frac{a}{2}\right) = a(x-a).$$

$$13. b(2x - 9c - 14b) = c(c-x).$$

$$14. a(x-a-2b) + b(x-b) + c(x+c) = 0.$$

$$15. (a-x)(x-b) + (a+x)(x-b) = (a-b)^2.$$

$$16. \frac{a-b+c}{x+a} = \frac{b-a+c}{x-a}.$$

$$17. \frac{1}{a(b-x)} + \frac{1}{b(c-x)} - \frac{1}{a(c-x)} = 0.$$

$$18. \frac{x-1}{a-1} - \frac{a-1}{x-1} = \frac{x^2-a^2}{(a-1)(x-1)}.$$

$$19. \frac{a+x}{a} - \frac{2x}{a+x} + \frac{x^2(x-a)}{a(x^2-x^2)} = \frac{1}{3}.$$

$$20. \frac{x+a}{b} + \frac{x+c}{a} + \frac{x+b}{c} = \frac{a}{b} + \frac{b}{c} + \frac{c}{a} + 1.$$

$$21. \frac{x^2-ax-bx+ab}{x-a} = \frac{x^2-2bx+2b^2}{x-b} - \frac{c^2}{x-c}.$$

$$22. \frac{1}{m+n} - \frac{2mn}{(m+n)^3} - \frac{m}{(m+n)^2} = \frac{x-n}{(m+n)^2}.$$

$$23. \frac{x}{a+b+c} + \frac{x}{a+b-c} = a^2 + b^2 + c^2 + 2ab.$$

$$24. \frac{x+2b}{x-b} + \frac{x+3b}{x+b} = \frac{x+b}{x-3b} + \frac{x+2b}{x+5b}.$$

$$25. \frac{2x+3a}{x+a} + \frac{3x+7a}{x+2a} = \frac{2a}{x+4a} + 5.$$

$$26. \frac{x+7a}{x+6a} + \frac{x-a}{x-3a} = \frac{x+7a}{x+a} + \frac{x-a}{x+2a}.$$

$$27. (a-b)(x-c) - (b-c)(x-a) = (c-a)(x-b).$$

PROBLEMS

DIRECTIONS FOR SOLVING. — *Represent one of the unknown numbers by x , and from the conditions of the problem find an expression for each of the other unknown numbers.*

Find from the problem two expressions that are equal, and write them as an equation.

Solve the equation.

1. A man bought a farm, a house, and a barn for \$12,600. If the house cost twice as much as the barn, and the farm twice as much as the house and barn together, how much did each cost?

SUGGESTION. — There are three unknown quantities—the cost of the house, the cost of the barn, and the cost of the farm. It is evident that, if the cost of the barn were known, the cost of the house could be found from it, and from both the cost of the farm.

Accordingly, represent the cost of the barn as x dollars, and express the cost of the house and the cost of the farm in terms of x . Discover two expressions for the total cost, and equate them.

2. If A is twice as old as B, and B twice as old as C, how old is each, if the sum of their ages is 140 years?

3. Mr. Henry bought three building lots for \$36,000. If the third cost twice as much as the second, and the second 3 times as much as the first, what was the cost of each?

4. A man left \$63,000 to his wife, two sons, and a daughter. If each son received twice as much as the daughter, and half as much as the wife, what was the share of each?

5. A man left $\frac{1}{2}$ of his property to his wife, $\frac{1}{4}$ to his son, $\frac{1}{8}$ to his daughter, and the rest, which was \$2000, to a hospital. What was the value of his property?

6. A owed B, C, and D \$27,000 in all. If he owed B 4 times as much as C, and D $\frac{3}{4}$ as much as B, what sum did he owe each?

7. A person spends $\frac{1}{5}$ of his annual income for board, $\frac{1}{4}$ for clothes, and \$260 for other expenses. If he saves $\frac{1}{3}$ of his income, what is his income?

8. A table and a chair cost \$11. The table and a picture cost \$14. If the chair and the picture together cost 3 times as much as the table, what was the cost of each article?

9. A and B together had \$50, and A and C had \$60. After each had spent \$5, A had $\frac{1}{4}$ as much as B and C together. How much had each at first?

10. A gave his age as follows: " $\frac{2}{3}$ of my age less $\frac{1}{3}$ of what it will be a year hence is equal to $\frac{1}{3}$ of my age 5 years ago." What was his age?

11. James is 5 years older than his sister, and 5 years hence he will be $3\frac{1}{2}$ times as old as his sister was 5 years ago. What is the age of each?

12. A's daily wages are $\frac{2}{3}$ of B's, and C's are $\frac{3}{4}$ of A's. If A and C together earn 25 cents more a day than B, what are the daily wages of each?

13. A man paid a debt of \$8.00 with an equal number of 5, 10, and 25-cent pieces. How many of each were there?

14. A man bought equal quantities of white and brown sugar, paying $6\frac{1}{4}$ cents a pound for the former and 5 cents a pound for the latter. How many pounds of each did he buy, if the whole quantity cost him \$1.80?

15. A field is twice as long as it is wide. By increasing its length 20 rods and its width 30 rods, the area will be increased 2200 square rods. What are its dimensions?

16. Three fifths of a certain number exceeds $\frac{1}{2}$ of it by 7. What is the number?

17. One third of a number added to 3 times the number is equal to 50. What is the number?

18. After spending $\frac{2}{3}$ of my income and \$300 more, I had $\frac{1}{4}$ of it left. What was my income?

19. The sum of $\frac{1}{4}$ of a number, $\frac{1}{3}$ of it, and $\frac{2}{3}$ of it is 4 more than $\frac{1}{4}$ of the number. What is the number?

20. The sum of a number, its half, its third, and its fourth, and 16 is 66. What is the number?

21. A man deposited $\frac{2}{3}$ of his month's wages in a bank, and paid out $\frac{1}{2}$ of the remainder for groceries and \$9 for dry goods. If he had \$3 left, how much money had he at first?

22. Divide 500 into two parts, such that the greater decreased by $\frac{1}{4}$ of the smaller is 5 times as much as the smaller decreased by $\frac{1}{2}$ of the larger.

23. Divide 61 into two parts, such that, if the greater is divided by the less, the quotient will be 3 and the remainder 1.

24. Divide 111 into two parts, such that the first diminished by 3 is equal to the second divided by 3.

25. Divide 40 into two parts, such that the first divided by 3 is equal to the second divided by 5.

26. Divide 40 into two parts, such that the first is 10 greater than twice the second.

27. Divide 44 into two parts, such that, if the greater is divided by 5 and the less by 7, the difference of the quotients will be 4.

28. Divide 54 into two parts, such that $\frac{1}{4}$ of the first part is equal to $\frac{1}{5}$ of the second.

SOLUTION

Let	$x = \frac{1}{4}$ of the first part or $\frac{1}{5}$ of the second part.
Then,	$4x =$ the first part,
and	$5x =$ the second part;
\therefore	$4x + 5x = 54;$
whence,	$x = 6,$
	$4x = 24,$ the first part,
and	$5x = 30,$ the second part.

29. Divide 40 into three parts, such that $\frac{1}{2}$ of the first, $\frac{1}{3}$ of the second, and $\frac{1}{4}$ of the third are equal.

30. Find three parts of 60, such that the first divided by 5, the second multiplied by 2, and the third increased by 5 are equal.

31. Divide 72 into four parts, such that the first divided by 2, the second diminished by 2, the third multiplied by 2, and the fourth increased by 2 are equal.

32. A can do a piece of work in 8 days. If B can do it in 10 days, in how many days can both working together do it?

SOLUTION

Let x = the required number of days.

Then, $\frac{1}{x}$ = the part of the work both can do in 1 day,

$\frac{1}{8}$ = the part of the work A can do in 1 day,

$\frac{1}{10}$ = the part of the work B can do in 1 day ;

$$\therefore \frac{1}{x} = \frac{1}{8} + \frac{1}{10} = \frac{9}{40}.$$

Solving, $x = \frac{40}{9}$, or $4\frac{4}{9}$, the required number of days.

33. A can do a piece of work in 10 days, and B can do it in 15 days. How long will it take both to do it?

34. Three pipes empty into a cistern. One can fill the cistern in 5 hours, another in 6 hours, and the third in 10 hours. How long will it take the three pipes together to fill it?

35. A can do a piece of work in 10 days, B can do it in 12 days, and C can do it in 8 days. In how many days can all together do it?

36. A can pave a walk in 6 days, and B can do it in 8 days. How long will it take A to finish the job after both have worked 3 days?

37. A can build a wall in 15 days, but with the aid of B and C, the wall can be built in 6 days. If B does as much work in 1 day as C does in 2 days, in how many days can B and C separately build the wall?

38. A and B can dig a ditch in 10 days, B and C can dig it in 6 days, and A and C in $7\frac{1}{2}$ days. In what time can each man do the work?

SUGGESTION. — Since A and B can dig $\frac{1}{10}$ of the ditch in 1 day, B and C $\frac{1}{6}$ of it in 1 day, and A and C $\frac{1}{7\frac{1}{2}}$ of it in 1 day, $\frac{1}{10} + \frac{1}{6} + \frac{1}{7\frac{1}{2}}$ is twice the part they can all dig in 1 day.

39. A and B can load a car in $2\frac{3}{4}$ hours, B and C in $3\frac{1}{2}$ hours, and A and C in $3\frac{1}{3}$ hours. How long will it take each alone to load it?

40. A boy bought some oranges at the rate of 30 cents a dozen. He sold $\frac{3}{4}$ of them for 4 cents each, and the rest for 3 cents each. If he gained 90 cents, how many oranges did he buy?

41. Find a fraction whose value is $\frac{4}{7}$ and whose denominator is 15 greater than its numerator.

42. Find a fraction whose value is $\frac{3}{8}$ and whose numerator is 3 greater than half of its denominator.

43. The numerator of a certain fraction is 8 less than the denominator; and if each term of the fraction is decreased by 5, the value of the fraction becomes $\frac{1}{3}$. What is the fraction?

44. The units' digit of a number expressed by two digits exceeds the tens' digit by 5. If the number increased by 63 is divided by the sum of its digits, the quotient is 10. What is the number?

SOLUTION

Let x = the digit in tens' place.

Then, $x + 5$ = the digit in units' place,

and $10x + (x + 5)$, or $11x + 5$ = the number;

$$\therefore \frac{11x + 5 + 63}{2x + 5} = 10;$$

whence, $x = 2$,

and $x + 5 = 7$.

Therefore, the number is 27.

45. The tens' digit of a number expressed by two digits is 3 times the units' digit. If the number diminished by 33 is divided by the difference of the digits, the quotient is 10. What is the number?

46. The tens' digit of a number expressed by two digits is $\frac{1}{2}$ of the units' digit. If the number increased by 27 is divided by the sum of its digits, the quotient is $6\frac{1}{4}$. What is the number?

47. In a purse containing \$1.45 there are $\frac{1}{2}$ as many quarters as 5-cent pieces and $\frac{3}{4}$ as many dimes as 5-cent pieces. How many pieces are there of each kind?

48. A woman spent \$10 more than $\frac{2}{3}$ of her money; then \$10 more than $\frac{2}{3}$ of the remainder. If she had \$2 left, how much money had she at first?

49. A man spent \$1 less than $\frac{2}{3}$ of his money and had left \$1 less than $\frac{2}{3}$ of it. How much money had he at first?

50. A girl found that she could buy 12 apples with her money and have 5 cents left, or 10 oranges and have 6 cents left, or 6 apples and 6 oranges and have 2 cents left. How much money had she?

51. A boy spent $\frac{1}{2}$ of his money and $\frac{1}{2}$ a cent more, then $\frac{1}{2}$ of the remainder and $\frac{1}{2}$ a cent more, then $\frac{1}{2}$ of what he had left and $\frac{1}{2}$ a cent more, when he found that he had 2 cents remaining. How much had he at first?

52. Five boys bought a boat, agreeing to share the expense equally. But one of them having left \$1 of his share unpaid, each of the others had to pay $\frac{1}{4}$ more than his share. What was the cost of the boat?

53. A sum of money was divided among A, B, C, and D so that A received $\frac{1}{2}$ as much as all the others, B received $\frac{1}{3}$ as much as all the others, C received $\frac{1}{4}$ as much as all the others, and D received \$2800 less than A. What sum did each receive?

54. In an alloy of 90 ounces of silver and copper there are 6 ounces of silver. How much copper must be added that 10 ounces of the new alloy may contain $\frac{2}{3}$ of an ounce of silver?

55. If 80 pounds of sea water contain 4 pounds of salt, how much fresh water must be added that 45 pounds of the new solution may contain $1\frac{1}{2}$ pounds of salt?

56. An officer, attempting to arrange his men in a solid square, found that with a certain number of men on a side he had 34 men over, but with 1 man more on a side he needed 35 men to complete the square. How many men had he?

SUGGESTION. — With x men on a side, the square contained x^2 men; with $x + 1$ men on a side, there were places for $(x + 1)^2$ men. Since the number of men was the same under both arrangements, $x^2 + 34 = (x + 1)^2 - 35$.

57. A regiment drawn up in the form of a solid square lost 60 men in battle. Afterward, when the men were arranged in a solid square with 1 man less on a side, it was found that there was 1 man over. How many men were there in the regiment at first?

58. A regiment drawn up in the form of a solid square was reinforced by 240 men. When the regiment was formed again in a solid square, there were 4 more men on a side. How many men were there in the regiment at first?

59. A man was hired for 40 days under the following conditions: for every day he worked he was to receive \$3 besides his board, while for every day he was idle he was to receive nothing, but was to be charged \$1.20 for his board. If at the end of the period he received \$57, how many days did he work?

60. A man invested \$800, a part at 6% and the rest at 5%. If the total annual interest was \$45, how much did he invest at each rate?

SUGGESTION. — Let x = the number of dollars invested at 6%.

Then, $800 - x$ = the number of dollars invested at 5%;

$$\therefore \frac{6}{100}x + \frac{5}{100}(800 - x) = 45.$$

61. A man has $\frac{2}{3}$ of his property invested at 4%, $\frac{1}{4}$ at 3%, and the remainder at 2%. How much property has he, if his annual income is \$860?

62. A man put out \$4330 in two investments. On one of them he gained 12%, and on the other he lost 5%. If his net gain was \$251, what was the amount of each investment?

63. There were distributed among 20 men and 25 women \$160 in such a way that the sum of what a man and a woman received was \$7. How much did the men receive, and how much did the women receive?

64. At what time between 5 and 6 o'clock will the hands of a clock be together?

SOLUTION

Let x = the number of minute spaces that the minute hand travels after 5 o'clock before they come together.

Then, $\frac{x}{12}$ = the number of minute spaces that the hour hand travels in the same time.

Since they are 25 minute spaces apart at 5 o'clock,

$$x - \frac{x}{12} = 25;$$

$$\therefore x = 27\frac{1}{3}, \text{ the number of minutes after 5 o'clock.}$$

65. At what time between 1 and 2 o'clock will the hands of a clock be together?

66. At what time between 6 and 7 o'clock will the hands of a clock be together?

67. At what time between 10 and 11 o'clock will the hands of a clock point in opposite directions?

68. At what times between 4 and 5 o'clock will the hands of a clock be 15 minute spaces apart?

69. When after 9 o'clock and before 10 o'clock will the hands of a clock be at right angles to each other?

70. A man rows downstream at the rate of 6 miles an hour and returns at the rate of 3 miles an hour. How far downstream can he go and return within 9 hours?

71. At the rate of 3 miles an hour uphill and 4 miles an hour downhill a woman can walk 60 miles in 17 hours. How much of the distance is uphill, and how much is downhill?

72. A hare pursued by a hound takes 4 leaps while the hound takes 3; but 2 of the hound's leaps are equal to 3 of the hare's. If the hare has a start equal to 60 of her own leaps, how many leaps must the hound take to catch the hare?

SOLUTION

Let $3x$ = the number of leaps taken by the hound.

Then, $4x$ = the number of leaps taken by the hare.

Suppose a = the number of feet in one leap of the hare.

Then, $\frac{3a}{2}$ = the number of feet in one leap of the hound,

$$\frac{3a}{2} \times 3x = \frac{9ax}{2}, \text{ the number of feet the hound runs,}$$

and $a \times 4x = 4ax$, the number of feet the hare runs.

Since the hare has a start equal to 60 times a feet, or $60a$ feet, the hare runs $60a$ feet less than the hound.

$$\text{Therefore, } 4ax = \frac{9ax}{2} - 60a.$$

$$\text{Dividing by } a, \quad 4x = \frac{9x}{2} - 60.$$

$$\text{Therefore, } x = 120,$$

and $3x = 360$, the number of leaps taken by the hound.

73. A fox is 70 leaps ahead of a hound and takes 5 leaps while the hound takes 3; but 3 of the hound's leaps equal 7 of the fox's. How many leaps must the hound take to catch the fox?

74. A rabbit makes 5 leaps while a dog makes 4; but 3 of the dog's leaps are equal to 4 of the rabbit's. If the rabbit has a start of 20 leaps, how many leaps will each take before the rabbit is caught?

75. A hound is 39 of his leaps behind a rabbit that takes 7 leaps while the hound takes 8. If 6 leaps of the rabbit are equal to 5 leaps of the hound, how many leaps must the hound take to catch the rabbit?

76. A wheelman and a pedestrian leave the same place at the same time to go to a point 54 miles distant, the former traveling 3 times as fast as the latter. The wheelman makes the trip and returning meets the pedestrian in $6\frac{3}{4}$ hours from the time they started. What is the rate of each?

77. If 1 pound of lead loses $\frac{5}{7}$ of a pound, and 1 pound of iron loses $\frac{2}{5}$ of a pound when weighed in water, how many pounds of lead and of iron are there in a mass of lead and iron that weighs 159 pounds in air and 143 pounds in water?

78. If 97 ounces of gold weighs 92 ounces when it is weighed in water, and 21 ounces of silver weighs 19 ounces when it is weighed in water, how many ounces of gold and of silver are there in a mass of gold and silver that weighs 320 ounces in air and 298 ounces in water?

79. A merchant increases his capital annually by $\frac{1}{4}$ of it, and at the end of each year takes out \$800 for expenses. At the end of three years, after taking out his expenses, he finds that his capital is \$6325. What was his original capital?

80. A merchant added annually to his capital $\frac{1}{4}$ of it, and at the end of each year took out \$1000 for expenses. If at the end of the third year, after taking out the last \$1000, he had $\frac{3}{4}$ of his original capital, what was his original capital?

81. A cistern can be filled by one pipe in 20 minutes, by another in 15 minutes, and it can be emptied by a third in 10

minutes. If the three pipes are running at the same time, how long will it take to fill the cistern?

82. A man walked from A to B at the rate of 2 miles an hour, and rode back at the rate of $3\frac{1}{2}$ miles an hour, being gone 13 hours. How far is it from A to B?

83. An express train whose rate is 40 miles an hour starts 1 hour and 4 minutes after a freight train and overtakes it in 1 hour and 36 minutes. How many miles does the freight train run per hour?

84. The distance from Albany to Syracuse is 148 miles. A canal boat leaves Albany for Syracuse, moving at the rate of 3 miles in 2 hours; at the same time another leaves Syracuse for Albany, moving at the rate of 5 miles in 4 hours. How far from Albany do they meet?

85. A steamer goes 5 miles downstream in the same time that it goes 3 miles upstream; but if its rate each way is diminished 4 miles an hour, its rate downstream will be twice its rate upstream. How fast does it go in each direction?

86. A, B, and C together can do a piece of work in $3\frac{1}{2}$ days; B can do $\frac{1}{2}$ as much as A, and C can do $\frac{3}{4}$ as much as B in a day. In how many days can each do the work alone?

87. A can do a piece of work in 6 days that B can do in 14 days. A began the work, and after a certain number of days B took his place and finished the work in 10 days from the time it was begun by A. How many days did B work?

88. In a certain weight of gunpowder the niter composed 10 pounds more than $\frac{3}{4}$ of the weight, the sulphur 3 pounds more than $\frac{1}{12}$, and the charcoal 3 pounds less than $\frac{1}{10}$ of the niter. What was the weight of the gunpowder?

89. A library containing 16,000 volumes was divided into five departments. In the department of history there were twice as many volumes as in the department of science, and 500 more than $\frac{1}{3}$ as many volumes as in the juvenile department. Of fiction there were $7\frac{1}{2}$ times as many volumes as of science, and 500 less than 8 times as many as in the reference department. How many volumes were there in each department?

90. An estate was divided among four heirs, A, B, C, and D. If the value of the estate had been \$1000 less, what A received would have been $\frac{1}{3}$ of it, and what B received $\frac{1}{4}$ of it; if the value of the estate had been $\frac{1}{2}$ greater, what C received would have been $\frac{1}{3}$ of it, and what D received $\frac{1}{4}$ of it. What sum did each receive?

91. A father takes 3 steps while his son takes 5; but 2 of the father's steps are equal to 3 of the son's. How many steps will the son require to overtake his father, who is 36 steps ahead?

92. A purse contained some money and a ring worth \$10 more than the money. If the purse was worth $\frac{1}{3}$ as much as the money it contained, and the purse and the money together were worth $\frac{1}{2}$ as much as the ring, what was the value of each?

93. Brass is $8\frac{1}{2}$ times as heavy as water, and iron $7\frac{1}{2}$ times as heavy as water. A mixed mass weighs 57 pounds, and when immersed displaces 7 pounds of water. How many pounds of each metal does the mass contain?

94. A man began business with \$4725, and annually added to his capital $\frac{1}{3}$ of it. At the end of each year he put aside a certain sum for expenses. If at the end of the third year, after taking out the sum for expenses, his capital was \$3800, what were his annual expenses?

95. The sum of two numbers is s , and their difference d . What are the numbers?

SOLUTION

Let x = the greater number.

Then, $x - d$ = the less number,

and $x + x - d = s$;

$$\therefore x = \frac{s+d}{2}, \text{ the greater number,} \quad (1)$$

$$\text{and} \quad x - d = \frac{s-d}{2}, \text{ the less number.} \quad (2)$$

If the sum of two numbers is 30, and their difference is 6, what are the numbers?

By (1), the greater number is $\frac{30+6}{2}$, or 18;

by (2), the less number is $\frac{30-6}{2}$, or 12.

A problem in which the numbers assumed to be known are represented by letters to which any values may be assigned is called a **General Problem**.

Problem 95 is a general problem.

The results obtained in solving a *general problem* may be considered *formulae* for solving similar problems.

96. Divide c cents between two boys so that one shall have d cents more than the other. If $c = 50$ and $d = 10$, how much will each receive?

97. A horse and a saddle are together worth a dollars, and the horse is worth m times as much as the saddle. What is the value of each? What, if $a = 160$ and $m = 3$?

98. Divide b into two parts one of which represents m times the other. What will they be, if b represents 100, and $m, 4$?

99. An estate of a dollars is divided between two heirs in the proportion of m to n . What is the share of each? What is the share of each, if $a = 40,000$, $m = 5$, and $n = 3$?

100. If A can do a piece of work in a days, and B in b days, in what time can both do it working together? Give the result, if $a = 10$ and $b = 15$.

101. An alloy of two metals is composed of m parts of one to n parts of the other. How many pounds of each are required in the composition of a pounds of the alloy?

Bell metal is an alloy of 5 parts of tin and 16 parts of copper. How many pounds of tin and of copper are there in a bell which weighs 4200 pounds?

102. A wheelman set out from B at the rate of r miles an hour. a hours later another started in pursuit at the rate of p miles an hour. How far from B will the second wheelman overtake the first? What will be the distance, if $r = 10$, $p = 12$, and $a = 8$?

103. A man traveled from home at the rate of a miles an hour and returned at the rate of b miles an hour. If he made the entire journey in h hours, how far from home did he go? How far, if $a = 4$, $b = 3\frac{1}{2}$, and $h = 15$?

SIMULTANEOUS SIMPLE EQUATIONS

TWO UNKNOWN NUMBERS

201. 1. If $x + y = 12$, what is the value of x ? of y ? How many values may x have? How many may y have?

2. In the expression $x + y = 12$, x and y each may have an indefinite number of values, but if, *at the same time*, $x - y = 4$, what is the value of x ? of y ?

3. Although one equation containing two unknown numbers has an indefinite number of values for each unknown number, or is *indeterminate*, what can be said about the values of the unknown numbers, when *two* equations are given involving the same values of the unknown numbers, but in different relations, that is, when two *independent equations* are given?

202. Two or more equations in which the unknown numbers have the same values are called **Simultaneous Equations**.

If x and y represent the same numbers in $2x + 3y = 19$ as they represent in $5x - y = 22$, $2x + 3y = 19$ and $5x - y = 22$ are simultaneous equations.

203. Equations that represent different relations between the unknown numbers, and so cannot be reduced to the same form, are called **Independent Equations**.

$3x + 3y = 18$ and $2x + 2y = 12$ really express but one relation between x and y ; viz., that their sum is 6. Hence, both equations may be reduced to the same form, as $x + y = 6$. But $3x + 3y = 18$ and $x + 3y = 14$ express different relations between x and y , and cannot be reduced to the same form. Hence, they are independent equations.

204. An equation whose unknown numbers may have an infinite number of values is called an **Indeterminate Equation**.

$x + y = 6$ is an indeterminate equation, because, if $x = 2$, $y = 4$; if $x = 3$, $y = 3$; if $x = 2\frac{1}{2}$, $y = 3\frac{1}{2}$; etc.

205. PRINCIPLE. — *Every single equation involving two or more unknown numbers is indeterminate.*

206. The process of deriving from a set of simultaneous equations, equations involving a less number of unknown numbers than is found in the given equations is called **Elimination**.

207. Two sets of simultaneous equations each having all the roots of the other set are called **Equivalent Systems** of equations.

$$\text{Thus,} \quad \left\{ \begin{array}{l} x + y = 8 \\ 3x + 2y = 21 \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{l} 2x + 2y = 16 \\ 3x + 2y = 21 \end{array} \right\}$$

are equivalent systems, for each system is satisfied by the same set of values, $x = 5$ and $y = 3$, and neither is satisfied by any other set of values.

208. Any equation in a set of simultaneous equations may be transformed into an equivalent equation by employing the principles of equivalency stated in § 196, since these principles apply to all equations. Hence, it remains to seek the principle by which simultaneous equations are combined in the process of elimination without introducing or losing roots.

This **Principle of Elimination** may be stated as follows, a and b being known multipliers, not zero, and either positive or negative:

If any equation of a system is replaced by the sum or difference of a times that equation and b times another equation of the system, the resulting system is equivalent to the given system.

The proof is as follows:

By § 196, Prin. 1, all the terms of the second member of an equation may be transposed to the first member.

Then, let

$$\left. \begin{array}{l} A = 0 \\ B = 0 \\ \dots \end{array} \right\} \quad (1)$$

be the given system, and let a and b be any known multipliers except zero and either positive or negative.

It is to be proved that the system

$$\left. \begin{array}{l} aA + bB = 0 \\ B = 0 \\ \dots \end{array} \right\} \quad (2)$$

is equivalent to the given system (1).

Since a and b are known multipliers, not zero, by § 196, Prin. 2, every set of values of the unknown numbers that makes $A = 0$ makes $aA = 0$, and every set of values that makes $B = 0$ makes $bB = 0$. Hence, every set of

values that satisfies (1) makes $aA + bB$ equal to zero and thus satisfies (2), since all of the equations of (1) and (2) except the first are the same.

Again, since every set of values that satisfies (2) makes B , or bB equal to zero and also makes $aA + bB = 0$, every such set of values makes $aA = 0$ and therefore, by § 196, Prin. 2, makes $A = 0$ and satisfies (1).

Since all the roots of (1) are roots of (2) and all the roots of (2) are roots of (1), (1) and (2) are equivalent systems.

209. Elimination by Addition and Subtraction.

1. If $2x + 2y = 10$ and $3x - 2y = 5$ are added, what is the resulting equation? When may an unknown number be eliminated by *addition*?

2. If $x + 2y = 5$ is subtracted from $3x + 2y = 11$, what is the resulting equation? When may a number be eliminated by *subtraction*?

PRINCIPLE. — *A literal number having the same coefficient in two equations may be eliminated by adding the equations, if the coefficients have unlike signs, or by subtracting one equation from the other, if the coefficients have like signs.*

EXAMPLES

1. Find the value of x and of y in $2x + 3y = 7$ and $3x + 4y = 10$.

PROCESS	
$2x + 3y = 7$	(1)
$3x + 4y = 10$	(2)
$6x + 9y = 21$	(3)
$6x + 8y = 20$	(4)
$y = 1$	(5)
$2x + 3 = 7$	(6)
$x = 2$	(7)

EXPLANATION. — Since x has not the same coefficients in both equations, the equations are multiplied by such numbers as will make the coefficients of x alike. Multiplying eq. (1) by 3 and eq. (2) by 2 gives equations (3) and (4). From these x is eliminated by subtraction, and the value of y is found.

The value of x may be found by multiplying eq. (1) and (2) by such numbers as will make the coefficients of y equal and then subtracting the resulting equations.

Or, the value of x may be found by substituting the value of y for y in one of the given equations, as eq. (1).

PROOF OF THE EQUIVALENCE. — By the principle of elimination, the system (1, 5), obtained by subtracting (4) from (3), is equivalent to the given system (1, 2). Hence, the only value of y in the given system is 1.

Since y represents 1 in the given system, (1) is equivalent to (6), which by § 196 is equivalent to (7). Hence, the given system, being equivalent to (1, 5), is equivalent to (7, 5), which is satisfied by one and only one set of values, $x = 2$ and $y = 1$.

RULE. — *If necessary, multiply the equations by such numbers as will cause the coefficients of one letter to be numerically equal in the resulting equations.*

When the signs of these coefficients are unlike, add the equations; when the signs are alike, subtract one equation from the other.

Solve by addition or subtraction:

$$2. \begin{cases} 7x - 5y = 52, \\ 2x + 5y = 47. \end{cases}$$

$$3. \begin{cases} 3x + 2y = 23, \\ x + y = 8. \end{cases}$$

$$4. \begin{cases} 3x - 4y = 7, \\ x + 10y = 25. \end{cases}$$

$$5. \begin{cases} 2x - 10y = 15, \\ 2x - 4y = 18. \end{cases}$$

$$6. \begin{cases} 3x - y = 4, \\ x + 3y = -2. \end{cases}$$

$$7. \begin{cases} 4x - y = 19, \\ x + 3y = 21. \end{cases}$$

$$8. \begin{cases} x + 2y = 5, \\ 2x + y = 1. \end{cases}$$

$$9. \begin{cases} 2x + 3y = 17, \\ 3x + 2y = 18. \end{cases}$$

$$10. \begin{cases} 3x + 4y = 25, \\ 4x + 3y = 31. \end{cases}$$

$$11. \begin{cases} 5x + 6y = 32, \\ 7x - 3y = 22. \end{cases}$$

$$12. \begin{cases} 3x + 6y = 39, \\ 9x - 4y = 51. \end{cases}$$

$$13. \begin{cases} 7x - 9y = 6, \\ x + 2y = 14. \end{cases}$$

$$14. \begin{cases} 13x - y = 20, \\ 4x + 2y = 20. \end{cases}$$

$$15. \begin{cases} 8x - 3y = 44, \\ 7x - 5y = 29. \end{cases}$$

$$16. \begin{cases} 6x - 5y = 33, \\ 4x + 4y = 44. \end{cases}$$

$$17. \begin{cases} x + 14y = 38, \\ 14x + y = 142. \end{cases}$$

$$18. \begin{cases} 5x + y = 12, \\ x + 5y = 36. \end{cases}$$

$$19. \begin{cases} 3x + 11y = 67, \\ 5x - 3y = 5. \end{cases}$$

$$20. \begin{cases} \frac{x}{4} + \frac{y}{2} = 12, \\ \frac{x}{4} - \frac{y}{2} = -2. \end{cases}$$

$$21. \begin{cases} \frac{x}{3} + \frac{y}{3} = 7, \\ \frac{x}{6} + \frac{y}{2} = 6\frac{1}{2}. \end{cases}$$

210. Elimination by Comparison.

1. If, in the simultaneous equations $x - y = 3$ and $x + 4y = 13$, the terms containing y are transposed, what will be the resulting equations?

2. Since the second members of these derived equations are each equal to x , how do they compare with each other?

3. If these second members, then, are placed equal to each other, how many unknown numbers will this equation contain?

4. How may an unknown number be eliminated from two simple simultaneous equations by *comparison*?

EXAMPLES

1. Find the value of x and of y in $2x - 3y = 10$ and $5x + 2y = 6$.

PROCESS

$$2x - 3y = 10 \quad (1)$$

$$5x + 2y = 6 \quad (2)$$

$$x = \frac{10 + 3y}{2} \quad (3)$$

$$x = \frac{6 - 2y}{5} \quad (4)$$

$$\therefore \frac{10 + 3y}{2} = \frac{6 - 2y}{5} \quad (5)$$

$$\therefore y = -2 \quad (6)$$

$$x = \frac{10 - 6}{2} \quad (3)$$

$$x = 2 \quad (7)$$

EXPLANATION. — Solving eq. (1) as if x were the only unknown number, the value of x is given as in eq. (3). In like manner, solving eq. (2) for x , another expression is found for the value of x , as in eq. (4).

Since the equations are simultaneous, the unknown numbers have the same values in each, and the two values of x form an equation, as eq. (5), from which x has been eliminated. Solving eq. (5), the value of y is found to be -2 , eq. (6).

By a similar process the value of x may be found; or, substituting the value of y for y in one of the preceding equations, as (3), the value of x is found to be 2, eq. (7).

PROOF OF THE EQUIVALENCE. — By § 196, (3) is equivalent to (1) and (4) is equivalent to (2). Hence, the system (3, 4) is equivalent to the given system.

Since (5), which by § 196 is equivalent to (6), may be obtained by subtracting (3) from (4) and transposing, by the principle of elimination, § 208, (6, 1), or (6, 3), is equivalent to (4, 3), or to (2, 1), the given system.

But since -2 is the only value of y in the given system, by § 196, (7) is equivalent to (3), or to (1). Hence, (7, 6), which is satisfied by one and only one set of values, $x = 2$ and $y = -2$, is equivalent to the given system. Therefore, the given system is satisfied by $x = 2$, $y = -2$ and by no other values of x and y .

RULE. — Find an expression for the value of the same unknown number in each equation, equate the two expressions, and solve the equation thus formed.

Solve by comparison :

$$2. \quad \begin{cases} 3x - 2y = 10, \\ x + y = 70. \end{cases}$$

$$9. \quad \begin{cases} 4x + 3y = 34, \\ 11x + 5y = 87. \end{cases}$$

$$3. \quad \begin{cases} 5x + y = 22, \\ x + 5y = 14. \end{cases}$$

$$10. \quad \begin{cases} 4x - 13y = 5, \\ 3x + 11y = -17. \end{cases}$$

$$4. \quad \begin{cases} 2x + 3y = 24, \\ 5x - 3y = 18. \end{cases}$$

$$11. \quad \begin{cases} 18x - 3y = 4y, \\ 1 - 4x + 3y = 27. \end{cases}$$

$$5. \quad \begin{cases} 3x + 5y = 14, \\ 2x - 3y = 3. \end{cases}$$

$$12. \quad \begin{cases} 7y - x = 0, \\ x + 2y = 18. \end{cases}$$

$$6. \quad \begin{cases} 3x + 2y = 36, \\ 5x - 9y = 23. \end{cases}$$

$$13. \quad \begin{cases} 3y + 9 = 5x, \\ 16 - 2x = 5y. \end{cases}$$

$$7. \quad \begin{cases} 2x + 7y = 8, \\ 3x + 9y = 9. \end{cases}$$

$$14. \quad \begin{cases} 5x - 40 = y, \\ 5y - 60 = x. \end{cases}$$

$$8. \quad \begin{cases} 4x + 6y = 19, \\ 3x - 2y = \frac{1}{2}. \end{cases}$$

$$15. \quad \begin{cases} 2y - 11x = 67, \\ 2x + 5y = 20. \end{cases}$$

211. Elimination by Substitution.

1. In an equation containing two unknown numbers, if the value of one is found, how is the value of the other obtained ?

2. Express the value of x in the first of the simultaneous equations $x + y = 5$ and $x + 2y = 7$ by transposing y to the second member.

3. When this expression for the value of x is substituted for x in the second equation, how many unknown numbers does the resulting equation contain ?

4. How may an unknown number be eliminated from simultaneous equations by *substitution* ?

EXAMPLES

1. Find the value of
- x
- and of
- y
- in
- $x+5y=9$
- and
- $3x-2y=10$
- .

PROCESS

$$x + 5y = 9 \quad (1)$$

$$3x - 2y = 10 \quad (2)$$

$$\underline{x = 9 - 5y} \quad (3)$$

$$3(9 - 5y) - 2y = 10 \quad (4)$$

$$\therefore y = 1 \quad (5)$$

$$x = 9 - 5 \quad (3)$$

$$x = 4 \quad (6)$$

EXPLANATION. — Solving eq. (1)

for x , $x = 9 - 5y$.Since the given equations are simultaneous, x has the same value in eq. (2) as in eq. (1).If $9 - 5y$ is substituted for x in eq. (2), the resulting equation will be true. Substituting and solving, $y = 1$.Substituting the value of y for y in eq. (3), $x = 4$, eq. (6).

PROOF OF THE EQUIVALENCE. — By § 196, (3) is equivalent to (1). In the process, (4) is obtained by substituting the expression equal to x in (3) for x in (2), or by substituting $3(9 - 5y)$ for $3x$ in (2). Since the substitution of $3(9 - 5y)$ for $3x$ may be performed by subtracting $3x = 3(9 - 5y)$ from (2), by the principle of elimination (4), or the equivalent equation (5), may take the place of (2) in the system (3, 2) equivalent to the given system. Hence, (3, 5) is equivalent to (1, 2).

Since the only value of y in the system (3, 5) is 1, (6) is equivalent to (3); and (6, 5), which is satisfied by only one set of values, $x = 4$, $y = 1$, is equivalent to (3, 5) and therefore to (1, 2).

RULE. — Find an expression for the value of one of the unknown numbers in one of the equations.

Substitute this value for that unknown number in the other equation, and solve the resulting equation.

Solve by substitution :

2.
$$\begin{cases} x - y = 4, \\ 4y - x = 14. \end{cases}$$

3.
$$\begin{cases} x + y = 10, \\ 6x - 7y = 34. \end{cases}$$

4.
$$\begin{cases} 3x - 4y = 26, \\ x - 8y = 22. \end{cases}$$

5.
$$\begin{cases} 6y - 10x = 14, \\ y - x = 3. \end{cases}$$

6.
$$\begin{cases} y + 1 = 3x, \\ 5x + 9 = 3y. \end{cases}$$

7.
$$\begin{cases} 17 = 3x + z, \\ 7 = 3z - 2x. \end{cases}$$

8.
$$\begin{cases} 4y = 10 - x, \\ y - x = 5. \end{cases}$$

9.
$$\begin{cases} 7z - 3x = 18, \\ 2z - 5x = 1. \end{cases}$$

10.
$$\begin{cases} 3 - 15y = -x, \\ 3 + 15y = 4x. \end{cases}$$

11.
$$\begin{cases} 1 - x = 3y, \\ 3(1 - x) = 40 - y. \end{cases}$$

Solve by any method, eliminating without clearing of fractions, when possible:

$$12. \begin{cases} x + \frac{y}{3} = 11, \\ \frac{x}{3} + 3y = 21. \end{cases}$$

$$19. \begin{cases} \frac{x+y}{2} - \frac{x-y}{3} = 8, \\ \frac{x+y}{3} + \frac{x-y}{4} = 11. \end{cases}$$

$$13. \begin{cases} \frac{3x}{4} + \frac{4y}{5} = 21, \\ \frac{2x}{3} + \frac{3y}{5} = 17. \end{cases}$$

$$20. \begin{cases} \frac{x}{2} - \frac{y}{3} - 1 = 0, \\ \frac{2x-1}{2} - \frac{3y-1}{3} = \frac{5}{6}. \end{cases}$$

$$14. \begin{cases} \frac{x}{3} = 11 - \frac{y}{2}, \\ \frac{x}{3} + \frac{2y}{7} = 8. \end{cases}$$

$$21. \begin{cases} \frac{7+x}{5} - \frac{2x-y}{4} = 3y-5, \\ \frac{5y-7}{2} + \frac{4x-3}{6} = 18-5x. \end{cases}$$

$$15. \begin{cases} \frac{x}{8} + 4y = 15, \\ \frac{x}{6} + \frac{2y}{3} = 6. \end{cases}$$

$$22. \begin{cases} \frac{x}{2} - 12 = \frac{y}{4} + 8, \\ \frac{x+y}{5} + \frac{x}{3} = \frac{2y-x}{4} + 35. \end{cases}$$

$$16. \begin{cases} \frac{x-1}{4} + y = 3, \\ \frac{x-1}{4} + 4y = 9. \end{cases}$$

$$23. \begin{cases} \frac{1}{x-1} - \frac{3}{x+y} = 0, \\ \frac{3}{x-y} + 3 = 0. \end{cases}$$

$$17. \begin{cases} \frac{3x}{4} + \frac{2y}{3} = 20, \\ \frac{x}{2} + \frac{3y}{4} = 17. \end{cases}$$

$$24. \begin{cases} \frac{x}{2} - 12 = \frac{y+32}{4}, \\ \frac{y}{8} + \frac{3x-2y}{5} = 25. \end{cases}$$

$$18. \begin{cases} \frac{x}{3} = \frac{y}{2}, \\ \frac{x}{3} - \frac{y}{3} = 1. \end{cases}$$

$$25. \begin{cases} \frac{.2y + .5}{1.5} = \frac{.49x - .7}{4.2}, \\ \frac{.5x - .2}{1.6} = \frac{41}{16} - \frac{1.5y - 11}{8}. \end{cases}$$

$$26. \begin{cases} \frac{x+y}{5} + \frac{x-y}{5} = \frac{2x-y}{8} + \frac{4y-x}{5} + 1, \\ \frac{3x}{4} - \frac{y}{5} = y + 12. \end{cases}$$

$$27. \begin{cases} x + \frac{1}{2}(3x - y - 1) = \frac{1}{4} + \frac{3}{4}(y - 1), \\ \frac{1}{5}(4x + 3y) = \frac{1}{10}(7y + 24). \end{cases}$$

$$28. \begin{cases} \frac{6x+9}{4} + \frac{3x+5y}{4x-6} = 3\frac{1}{4} + \frac{3x+4}{2}, \\ \frac{8y+7}{10} + \frac{6x-3y}{2(y-4)} = 4 + \frac{4y-9}{5}. \end{cases}$$

$$29. \begin{cases} \frac{3x-5y}{3} - \frac{2x-8y-9}{12} = \frac{31}{12}, \\ \left(\frac{x}{7} + \frac{y}{4} + 1\frac{1}{8}\right) - \left(4x - \frac{y}{8} - 25\right) = \frac{5}{6}. \end{cases}$$

$$30. \begin{cases} x - 20 - \frac{2y-x}{23-x} = \frac{2x-59}{2}, \\ y - \frac{3-y}{x-18} - 30 = \frac{3y-73}{3}. \end{cases}$$

$$31. \begin{cases} \frac{3x+6}{7} - \frac{x+5}{3y-5} = \frac{6x-2}{14}, \\ \frac{2y-3}{6} + \frac{3y+4}{5x-7} = \frac{3y+5}{9}. \end{cases}$$

$$32. \begin{cases} \frac{x}{2} + \frac{16-x}{2} = 30 + \frac{5y+2x}{40-x}, \\ \frac{2y-3}{y+8} + \frac{83-8y}{8} = 10 - y. \end{cases}$$

$$33. \begin{cases} 4x + \frac{2y-\frac{x}{2}}{17-3x} = 5 + \frac{16x-1}{4}, \\ 50 - \frac{y-1}{\frac{5x-10}{3}} = 8y + \frac{147-24y}{3}. \end{cases}$$

34. Solve
$$\begin{cases} \frac{4}{x} - \frac{3}{y} = \frac{14}{5}, \\ \frac{2}{x} + \frac{5}{y} = \frac{25}{3}. \end{cases}$$

SOLUTION

$$\frac{4}{x} - \frac{3}{y} = \frac{14}{5} \quad (1)$$

$$\frac{2}{x} + \frac{5}{y} = \frac{25}{3} \quad (2)$$

$$(2) \times 2, \quad \frac{4}{x} + \frac{10}{y} = \frac{50}{3} \quad (3)$$

$$(3) - (1), \quad \frac{13}{y} = \frac{208}{15} \quad (4)$$

$$\frac{1}{y} = \frac{16}{15} \quad (5)$$

$$\therefore y = \frac{15}{16} \quad (6)$$

Substituting the value of $\frac{1}{y}$ in equation (1),

$$\frac{4}{x} - \frac{48}{15} = \frac{14}{5} \quad (7)$$

$$\frac{1}{x} = \frac{3}{2} \quad (8)$$

$$\therefore x = \frac{2}{3} \quad (9)$$

In fractional equations in which the denominators are simple expressions like the above, an unknown number should be eliminated before clearing of fractions.

35.
$$\begin{cases} \frac{x}{2} - \frac{2y}{3} = -2, \\ \frac{5x}{2} + \frac{y}{3} = 12. \end{cases}$$

37.
$$\begin{cases} \frac{5}{x} + \frac{6}{y} = 64, \\ \frac{6}{x} + \frac{5}{y} = 73\frac{1}{2}. \end{cases}$$

36.
$$\begin{cases} \frac{x}{3} + \frac{y}{3} = 13, \\ \frac{4y}{3} - \frac{x}{3} = -8. \end{cases}$$

38.
$$\begin{cases} \frac{10}{x} + \frac{5}{y} = 20, \\ \frac{5}{x} + \frac{10}{y} = 57\frac{1}{2}. \end{cases}$$

$$39. \begin{cases} \frac{5}{x} - \frac{3}{y} = -2, \\ \frac{25}{x} + \frac{1}{y} = 6. \end{cases}$$

$$42. \begin{cases} \frac{7}{x} + \frac{8}{y} = 30, \\ \frac{7}{y} + \frac{8}{x} = 30. \end{cases}$$

$$40. \begin{cases} \frac{2}{x} - \frac{3}{y} = 5, \\ \frac{5}{x} - \frac{2}{y} = 7. \end{cases}$$

$$43. \begin{cases} \frac{3}{2x} - \frac{1}{y} = -3, \\ \frac{5}{2x} + \frac{3}{y} = 23. \end{cases}$$

$$41. \begin{cases} \frac{4}{x} + \frac{3}{y} = \frac{9}{8}, \\ \frac{3}{x} + \frac{4}{y} = \frac{11}{12}. \end{cases}$$

$$44. \begin{cases} \frac{7}{8x} - \frac{2}{3y} = 10, \\ \frac{5}{6x} - \frac{2}{11y} = 17. \end{cases}$$

LITERAL SIMULTANEOUS EQUATIONS

$$1. \text{ Solve } \begin{cases} ax + by = m, \\ cx + dy = n. \end{cases}$$

SOLUTION

$$ax + by = m \quad (1)$$

$$cx + dy = n \quad (2)$$

$$(1) \times d, \quad adx + bdy = dm \quad (3)$$

$$(2) \times b, \quad bcx + bdy = bn \quad (4)$$

$$(3) - (4), \quad (ad - bc)x = dm - bn$$

$$\therefore x = \frac{dm - bn}{ad - bc} \quad (5)$$

$$(1) \times c, \quad acx + bcy = cm \quad (6)$$

$$(2) \times a, \quad acx + ady = an \quad (7)$$

$$(7) - (6), \quad (ad - bc)y = an - cm$$

$$\therefore y = \frac{an - cm}{ad - bc} \quad (8)$$

In literal simultaneous equations, elimination is usually performed by the method of addition and subtraction.

$$2. \begin{cases} ax + by = m, \\ bx - ay = c. \end{cases}$$

$$3. \begin{cases} ax - by = m, \\ cx - dy = r. \end{cases}$$

$$4. \begin{cases} ax = by, \\ x + y = ab. \end{cases}$$

$$5. \begin{cases} x - ay = n, \\ bx + y = p. \end{cases}$$

$$6. \begin{cases} a(x - y) = 5, \\ bx - cy = n. \end{cases}$$

$$7. \begin{cases} a(a - x) = b(y - b), \\ ax = by. \end{cases}$$

$$8. \begin{cases} x + y = b - a, \\ bx - ay + 2ab = 0. \end{cases}$$

$$9. \begin{cases} \frac{1}{x} + \frac{1}{y} = \frac{1}{a}, \\ \frac{1}{x} - \frac{1}{y} = \frac{1}{b}. \end{cases}$$

$$10. \begin{cases} \frac{a}{x} - \frac{b}{y} = -1, \\ \frac{b}{x} - \frac{a}{y} = -1. \end{cases}$$

$$11. \begin{cases} \frac{x}{a} + \frac{y}{b} = 2ab, \\ \frac{x}{ab} + \frac{y}{ab} = a + b. \end{cases}$$

$$12. \begin{cases} \frac{x}{a} + \frac{y}{b} - 2 = 0, \\ bx - ay = 0. \end{cases}$$

$$13. \begin{cases} \frac{x}{a} + \frac{y}{b} = 1, \\ \frac{x}{b} - \frac{y}{a} = \frac{1}{2}. \end{cases}$$

$$14. \begin{cases} \frac{1}{ax} + \frac{1}{by} = c, \\ \frac{1}{bx} - \frac{1}{ay} = d. \end{cases}$$

$$15. \begin{cases} \frac{a}{x} + \frac{b}{y} = c, \\ \frac{m}{x} + \frac{n}{y} = e. \end{cases}$$

$$16. \begin{cases} \frac{x+1}{y+1} = \frac{a+b+1}{a-b+1}, \\ x-y = 2b. \end{cases}$$

$$17. \begin{cases} \frac{x+y}{a} = \frac{x-y}{b}, \\ \frac{x-y}{a} = \frac{y-a}{b}. \end{cases}$$

$$18. \begin{cases} \frac{1}{x-a} = \frac{1}{a-y}, \\ \frac{x+y}{x-y} = a. \end{cases}$$

$$19. \begin{cases} \frac{x}{a} + \frac{y}{b} = c, \\ \frac{x}{b} + \frac{y}{c} = d. \end{cases}$$

$$20. \begin{cases} \frac{a}{a+x} + \frac{b}{b-y} = \frac{a}{b}, \\ \frac{b}{a+x} + \frac{a}{b-y} = \frac{b}{a}. \end{cases}$$

PROBLEMS

1. There are two numbers such that if twice the first is added to 3 times the second, the sum will be 130; but if 5 times the first is diminished by the second, the remainder will be 70. What are the numbers?

SOLUTION

Let	$x =$ the first number,
and	$y =$ the second number.
Then,	$2x + 3y = 130,$
and	$5x - y = 70.$
Eliminating y ,	$17x = 340,$
	$x = 20.$
Whence, by substitution,	$y = 30.$

2. A drover sold 3 cows and 7 horses to one person for \$600, and to another person, at the same prices, 3 cows and 3 horses for \$300. How much per head did he get for each?

3. With \$30 a man can buy 20 yards of one kind of cloth and 50 yards of another; with \$23 he can buy 30 yards of the first kind and 20 yards of the second kind. What is the price of each per yard?

4. If 45 bushels of wheat and 37 bushels of rye together cost \$62.70, and 37 bushels of wheat and 25 bushels of rye, at the same prices, cost \$48.30, what is the price of each per bushel?

5. Henry expended 95 cents for apples and oranges, paying 5 cents for each orange and 4 cents for each apple. If he had 22 of both, how many of each did he buy?

6. Five years ago A was $\frac{1}{3}$ as old as B, and 10 years hence he will be $\frac{1}{2}$ as old as B. What are their ages?

7. A said to B, "If you were twice as old, and I were $\frac{1}{2}$ as old, or if you were $\frac{1}{3}$ as old, and I were 3 times as old, the sum of our ages would be 70." How old was each?

8. A boy is given 28 cents to buy a dozen cakes. He finds that some cost 2 cents each and some 3 cents each. How many of each kind can he purchase?

9. A said to B, "Give me \$20, and I shall have 3 times as much money as you." B replied, "Give me \$5, and I shall have twice as much money as you." How much money had each?

SOLUTION

Let x = the number of dollars A had,
 and y = the number of dollars B had.
 Then, $x + 20 = 3(y - 20)$,
 and $y + 5 = 2(x - 5)$.
 Solving, $x = 25$, the number of dollars A had,
 and $y = 35$, the number of dollars B had.

10. If A gives B \$100, B will have 4 times as much money as A; but if B gives A \$200, A will have 4 times as much money as B. What sum of money has each?

11. A said to B, "Give me 20 cents of your money, and I shall have half as much as you." B replied, "Give me 25 cents of your money, and I shall have 5 times as much as you." How much had each?

12. If A had \$300 more, he would have twice as much as B; if B had \$300 less, he would have $\frac{1}{2}$ as much as A. How much money has each?

13. If 1 is added to each term of a fraction, its value will be $\frac{2}{3}$; if 1 is subtracted from each term of the fraction, its value will be $\frac{1}{2}$. What is the fraction?

SOLUTION

Let $\frac{x}{y}$ represent the fraction.
 Then, $\frac{x+1}{y+1} = \frac{2}{3}$,
 and $\frac{x-1}{y-1} = \frac{1}{2}$.
 Solving, $x = 3$,
 and $y = 5$.
 That is, $\frac{3}{5}$ is the fraction.

14. If 1 is added to the numerator of a certain fraction, its value becomes $\frac{2}{3}$; if 2 is added to the denominator, its value becomes $\frac{1}{2}$. What is the fraction?

15. Find a fraction that is equal to $\frac{1}{3}$ when its terms are diminished by 2, and is equal to $\frac{2}{3}$ when its terms are increased by 2.

16. A certain number expressed by two digits is equal to 7 times the sum of its digits; if 27 is subtracted from the number, the difference will be expressed by reversing the order of the digits. What is the number?

SOLUTION

Let x = the digit in tens' place,
and y = the digit in units' place.
Then, $10x + y$ = the number,
and $10y + x$ = the number with its digits reversed;
 $\therefore 10x + y = 7(x + y)$,
and $10x + y - 27 = 10y + x$.
Solving, $x = 6$,
and $y = 3$.
Hence, $10x + y = 60 + 3$, or 63, the number.

17. The sum of the two digits of a certain number is 12, and the number is 3 greater than 6 times the sum of its digits. What is the number?

18. When a certain number expressed by two digits is divided by the sum of its digits, the quotient is 8; and when the first digit is diminished by 3 times the second, the remainder is 1. What is the number?

19. The sum of the two digits of a number is 12. If the order of the digits is reversed, the number will lack 12 of being doubled. What is the number?

20. A farmer bought 100 acres of land for \$3250. If part of it cost him \$40 an acre and the rest of it \$15 an acre, how many acres were there of each kind?

21. The admission to an entertainment was 50 cents for adults and 35 cents for children. If the proceeds from 100 tickets amounted to \$39.50, how many tickets of each kind were sold?

22. A man paid a bill of \$16 in 25-cent pieces and 5-cent pieces. If the number of coins was 80, how many of each kind were there?

23. A man paid \$1 for some apples at 3 cents each and some oranges at 5 cents each. He sold $\frac{2}{3}$ of the apples and $\frac{1}{4}$ of the oranges at cost for 34 cents. How many of each did he buy?

24. A and B together can do a piece of work in 12 days. After A has worked alone for 5 days, B finishes the work in 26 days. In what time can each alone do the work?

25. A blacksmith and his son had a contract to make a certain number of horseshoes. If both had worked together, they could have done the work in 6 days. But the father worked 8 days, and the son finished the work in 3 days. In how many days could each have done the work?

26. A man and his two sons can dig a ditch in 6 days; if the man and either son work 7 days, the other son can complete the ditch by working 2 days. In what time can each alone dig the ditch?

27. A certain number of persons agree to share equally the expense of hiring a coach. If each paid 75 cents, there would be \$1.25 over; but if each paid 50 cents, there would be \$2.50 lacking. What is the number of persons and the expense of hiring the coach?

28. A train ran a certain distance at a uniform rate. Had the rate been increased 5 miles an hour, the journey would have been 2 hours shorter; but had the rate been diminished 5 miles an hour, the journey would have been $2\frac{1}{2}$ hours longer. What was the distance and the rate of the train?

SUGGESTION. — Let x miles per hour be the actual rate of the train and y the number of hours required to complete the journey.

29. A sum of money was divided equally among a certain number of persons. If there had been 4 persons more, the share of each would have been \$3 less; but if there had been

2 persons less, the share of each would have been \$2 more. Among how many persons was the money divided and what was the share of each?

30. A dealer had eggs to sell and wished to buy potatoes. He found that 6 dozen eggs were worth 10 cents more than 2 bushels of potatoes; and that 10 dozen eggs were worth 10 cents less than 4 bushels of potatoes. How much were eggs and potatoes worth?

31. If a rectangular floor were 2 feet wider and 5 feet longer, its area would be 140 square feet greater; if it were 7 feet wider and 10 feet longer, its area would be 390 square feet greater. What are its dimensions?

32. If 54 is added to a certain number, expressed by two digits whose sum is 8, the order of the digits will be reversed. What is the number?

33. If 13 is added to a certain number, the sum will be equal to 5 times the sum of the two digits of the number; and if 36 is added to the number, the order of its digits will be reversed. What is the number?

34. A and B can do a piece of work in a days, or if A works m days alone, B can finish the work by working n days. In how many days can each do the work?

35. A can build a wall in c days, and B can build it in d days. How many days must each work so that, after A has done a part of the work, B can take his place and finish the wall in a days from the time A began?

36. One cask contains a mixture of 20 gallons of wine and 30 gallons of water, and another contains a mixture of 12 gallons of wine and 15 gallons of water. How many gallons must be drawn from each cask so that the mixture will contain 8 gallons of wine and 11 gallons of water?

SUGGESTION. — If x gallons of the mixture are drawn from the first cask, $\frac{2}{3}x$ gallons of it will be wine.

If y gallons of the mixture are drawn from the second cask, $\frac{4}{5}y$ gallons of it will be wine.

37. "If I had received 3 oranges more for my money," said A, "they would have cost me 1 cent less each; but if I had received 2 less, they would have cost me 1 cent more each." How many oranges had he bought, and at what price each?

38. A merchant mixes two kinds of tea. If he mixes it in parts proportional to 7 and 5, the value of the mixture is 46 cents a pound. If he mixes it in parts proportional to 5 and 7, the value of the mixture is 50 cents a pound. What is each kind of tea worth per pound?

39. A man invested \$4000, a part at 5% and the rest at 4%. If the annual income from both investments was \$175, what was the amount of each investment?

40. A man invested a dollars, a part at $r\%$ and the rest at $s\%$ yearly. If the annual income from both investments was b dollars, what was the amount of each investment?

41. A sum of money, at simple interest, amounted to b dollars in t years, and to a dollars in s years. What was the principal, and what was the rate of interest?

42. A had a certain sum invested at a certain rate per cent, and B had \$100 less invested at a rate 2% higher. B's annual income was \$56 greater than A's; but if B's rate upon his investment had been only 1% higher than A's his annual income would have been only \$25 greater than A's. How much was invested by each man, and at what rate?

43. A crew can row 8 miles downstream and back, or 12 miles downstream and half the way back in $1\frac{1}{2}$ hours. What is the rate of rowing in still water and the velocity of the stream?

44. A man rows 15 miles downstream and back in 11 hours. The current is such that he can row 8 miles downstream in the same time as 3 miles upstream. What is his rate of rowing in still water, and what is the velocity of the stream?

45. A box will hold 18 quires of paper and 18 bunches of envelopes, or 20 quires of paper and 15 bunches of envelopes. How many quires of paper will the box hold? How many bunches of envelopes will it hold?

46. A shelf will hold 20 arithmetics and 24 algebras or 15 arithmetics and 36 algebras. How many arithmetics will the shelf hold? How many algebras will it hold?

47. Two men had a certain distance to row and took turns in rowing. Whenever the first rowed, the boat moved at a rate sufficient to cover the entire distance in 10 hours, and whenever the second rowed, in 14 hours. If the journey was completed in 12 hours, how many hours did each row?

48. A train ran 1 hour and 36 minutes, and was then detained 40 minutes. It then proceeded at $\frac{3}{4}$ of its former rate and reached its destination 16 minutes late. If the detention had occurred 10 miles farther on, the train would have arrived 20 minutes late. At what rate did the train set out, and what was the whole distance traveled?

49. A certain number of people charter an excursion boat, agreeing to share the expense equally. If each pays a cents, there will be b cents lacking from the necessary amount; and if each pays c cents, d cents too much will be collected. How many persons are there, and how much should each pay?

50. A sum of money was to be divided equally among a certain number of persons, but a persons more than were expected appeared to claim a share, and in consequence each received b dollars less. If there had been c persons less than were expected, each would have received d dollars more. How many persons were there, and how much did each receive?

Give the results when $a = 5$, $b = 100$, $c = 4$, and $d = 125$.

51. A and B working together can do a piece of work in a days. But finding it impossible to work at the same time, A works b days, and later B finishes the work in c days. In how many days can each do the work alone?

If $a = 5\frac{1}{11}$, $b = 5$, and $c = 6$, in how many days can each do the work alone?

52. A purse holds c crowns and d guineas; a crowns and b guineas will fill $\frac{m}{n}$ th of it. How many will it hold of each?

How many, if $c = 12$, $d = 6$, $a = 4$, $b = 6$, $m = 1$, and $n = 2$?

53. A mine is emptied of water by two pumps which together discharge m gallons per hour. Both pumps can do the work in b hours, or the larger can do it in a hours. How many gallons per hour does each pump discharge? What is the discharge of each per hour when $a = 5$, $b = 4$, and $m = 1250$?

54. Two trains are scheduled to leave A and B, m miles apart, at the same time, and to meet in b hours. If the train that leaves B is a hours late and runs at its customary rate, it will meet the first train in c hours. What is the rate of each train?

What is the rate of each, if $m = 800$, $c = 9$, $a = 1\frac{1}{2}$, and $b = 10$?

55. If a quarts of good wine is mixed with b quarts of poorer wine, the mixture will be worth c cents a quart; if b quarts of the better wine is mixed with a quarts of the poorer, the mixture will be worth d cents a quart. What is each kind of wine worth per quart? What is each kind of wine worth per quart, if $a = 40$, $b = 20$, $c = 100$, and $d = 80$?

212. Discussion of the general solution of a system of two simultaneous simple equations involving two unknown numbers.

$$\text{Let} \quad ax + by = c \quad (1)$$

$$\text{and } * \quad a'x + b'y = c' \quad (2)$$

be any two simultaneous simple equations.

$$(1) \times b', \quad ab'x + bb'y = b'c \quad (3)$$

$$(2) \times b, \quad a'bx + bb'y = bc' \quad (4)$$

$$(3) - (4), \quad (ab' - a'b)x = b'c - bc' \quad (5)$$

$$(1) \times a', \quad aa'x + a'by = ca' \quad (6)$$

$$(2) \times a, \quad aa'x + ab'y = c'a \quad (7)$$

$$(7) - (6), \quad (ab' - a'b)y = c'a - ca' \quad (8)$$

By the principles of equivalence the given system may be replaced by the equivalent system (5, 8), in which (5) involves x alone and (8) involves y alone. By § 197, each of these simple equations involving one unknown number has one and only one root, which can be found except when the common coefficient of x and y is equal to zero. Hence, when $ab' - a'b$ is not equal to zero, the given system is satisfied by one and only one set of values

$$x = \frac{b'c - bc'}{ab' - a'b} \quad \text{and} \quad y = \frac{c'a - ca'}{ab' - a'b}.$$

* In algebraic notation a' , a'' , a''' , etc., are read ' a prime,' ' a second,' ' a third,' etc.

If $ab' - a'b = 0$, that is, if $ab' = a'b$, the first members of (3) and (4) are identical, and therefore the second members must be equal. The same is true of equations (6) and (7). Hence, (4) is only a different form of (3), and (7) is only a different form of (6); that is, (1) and (2) are not independent equations.

But if $ab' - a'b$ is not equal to zero, neither (3) and (4) nor (6) and (7) are reducible to the same form; that is, (1) and (2) are independent equations.

It is evident from the Distributive Law for multiplication that the equations (3) and (4), and also (6) and (7), cannot be combined by addition or subtraction unless x and y have the same values in (2) as in (1); that is, that (1) and (2) cannot be solved unless they are *simultaneous* equations.

It is evident that equations (5) and (8), and therefore (1) and (2), cannot be solved if $ab' - a'b = 0$, since § 196, Prin. 2, the members cannot be divided by a known expression equal to 0; and it has been shown that (1) and (2) are dependent or independent equations according as $ab' - a'b$ is or is not equal to zero.

Hence, it follows that :

Two simple equations involving two unknown numbers cannot be solved unless the equations are simultaneous and independent.

Every system of two independent simultaneous simple equations involving two unknown numbers can be solved, and is satisfied by one, and only one, set of values of its unknown numbers.

THREE OR MORE UNKNOWN NUMBERS

213. 1. In the equations $x + 2y + z = 8$ and $2x + y - z = 1$, how may z be eliminated?

2. If one of the unknown numbers in the above equations is eliminated, how many unknown numbers will be left?

3. How many independent equations are necessary before the values of two unknown numbers can be found?

4. How many independent equations containing the same two unknown numbers can be formed by combining the equations in (1)?

5. Since only one derived equation containing two unknown numbers was obtained from the given equations by eliminating z , and since we must have two such equations before we can find the values of x and y , if another independent equation involving x , y , and z is combined with either of the equations in (1), how

many independent equations containing x and y only will be available for finding the values of x and y ?

6. When the values of x and y are found, how may the value of z be found?

7. Then, how many independent equations containing three unknown numbers must be given, so that the values of the unknown numbers may be found? How many to find the values of four unknown numbers?

214. PRINCIPLE. — *Every system of independent simultaneous simple equations involving the same number of unknown numbers as there are equations can be solved, and is satisfied by one and only one set of values of its unknown numbers.*

The above principle may be established as follows:

From the given system of n equations involving n unknown numbers, a second system of $n - 1$ equations involving $n - 1$ unknown numbers may be derived by eliminating one of the unknown numbers; from the second system a third system of $n - 2$ equations involving $n - 2$ unknown numbers may be derived; and this process may be continued until the n th system, a single simple equation involving only one unknown number, is obtained.

Since, § 197, this equation has one and only one root, by substituting this value in either of the two equations of the next preceding system and solving, one and only one value of the other number in that equation is obtained; by substituting these two values in any one of the three equations of the next preceding system, one and only one value of the remaining unknown number in that equation is obtained; and by continuing this process, the value of each of the other unknown numbers is obtained.

By the principles of equivalent equations, the following system of n equations may be substituted for the given system: the single equation finally derived by elimination and composing the n th system; either of the two equations of the preceding, or $(n - 1)$ th system; any one of the three equations of the system preceding that, or of the $(n - 2)$ th system; and so on to any one of the n equations in the 1st or given system.

But each of the n equations just described has one and only one value of an unknown number. Hence, the given system can be solved, and is satisfied by one and only one set of values of its unknown numbers.

If the number of unknown numbers is greater than the number of independent simultaneous equations, the last equation obtained by repeated eliminations is indeterminate, and hence the system is indeterminate.

If the number of unknown numbers is less than the number of independent simultaneous equations, say $n - p$, any $n - p$ of the equations involving the $n - p$ unknown numbers form a determinate system.

EXAMPLES

1. Find the values of x , y , and z in $\begin{cases} x + 2y + 3z = 14, \\ 2x + y + 2z = 10, \\ 3x + 4y - 3z = 2. \end{cases}$

SOLUTION

$$x + 2y + 3z = 14 \quad (1)$$

$$2x + y + 2z = 10 \quad (2)$$

$$3x + 4y - 3z = 2 \quad (3)$$

Eliminating z by combining (1) and (3),

$$(1) + (3), \quad 4x + 6y = 16 \quad (4)$$

Eliminating z by combining (2) and (3),

$$(2) \times 3, \quad 6x + 3y + 6z = 30 \quad (5)$$

$$(3) \times 2, \quad 6x + 8y - 6z = 4 \quad (6)$$

$$\text{Adding,} \quad 12x + 11y = 34 \quad (7)$$

Eliminating x by combining (7) and (4),

$$(4) \times 3, \quad 12x + 18y = 48 \quad (8)$$

$$(8) - (7), \quad 7y = 14 \quad (9)$$

$$\therefore y = 2 \quad (10)$$

$$\text{Substituting in (4),} \quad 4x + 12 = 16 \quad (11)$$

$$\therefore x = 1 \quad (12)$$

Substituting the values of x and y in (1),

$$1 + 4 + 3z = 14 \quad (13)$$

$$\therefore z = 3 \quad (14)$$

EXPLANATION. — Eliminating z from (1) and (3) and from (2) and (3), two simultaneous equations, (4) and (7), are obtained involving x and y . By the principle of elimination, § 208, the new system (1, 4, 7), or (2, 4, 7), or (3, 4, 7), is equivalent to the given system.

Eliminating x from (4) and (7), a simple equation involving but one unknown number y is obtained, and from this equation the value of y is found, equation (10). Hence, the system (1, 4, 7) has been replaced by the equivalent system (1, 4, 10), which is, therefore, equivalent to the given system.

Substituting 2 for y in (4), the value of x is found, giving a new system (1, 12, 10) equivalent to the given system. Substituting the values of both x and y in (1), the value of z is found, giving the desired system (14, 12, 10) equivalent to the given system.

Solve the following:

$$2. \begin{cases} 2x - y + 2z = 12, \\ x + 3y + z = 41, \\ 2x + y + 4z = 22. \end{cases}$$

$$11. \begin{cases} 3x - 2y + z = 2, \\ 2x + 5y + 2z = 27, \\ x + 3y + 3z = 25. \end{cases}$$

$$3. \begin{cases} 3x + 5y - z = 8, \\ 4x + 3y + 2z = 47, \\ 6x + 5y - 2z = 11. \end{cases}$$

$$12. \begin{cases} 4x - 5y + 3z = 14, \\ x + 7y - z = 13, \\ 2x + 5y + 5z = 36. \end{cases}$$

$$4. \begin{cases} x + 3y - z = 10, \\ 2x + 5y + 4z = 57, \\ 3x - y + 2z = 15. \end{cases}$$

$$13. \begin{cases} 2x + y - 3z + 4w = 44, \\ 3x - 2y + z - w = -1, \\ 4x - y + 2z + w = 55, \\ 5x - 3y + 4z - w = 39. \end{cases}$$

$$5. \begin{cases} x + y + z = 53, \\ x + 2y + 3z = 105, \\ x + 3y + 4z = 134. \end{cases}$$

$$14. \begin{cases} 7x - 1 = 3y, \\ 11z - 1 = 7v, \\ 4z - 1 = 7y, \\ 19x - 1 = 3v. \end{cases}$$

$$6. \begin{cases} x - y + z = 30, \\ 3y - x - z = 12, \\ 7z - y + 2x = 141. \end{cases}$$

$$15. \begin{cases} x + \frac{1}{2}y + \frac{1}{3}z = 32, \\ \frac{1}{3}x + \frac{1}{4}y + \frac{1}{5}z = 15, \\ \frac{1}{4}x + \frac{1}{5}y + \frac{1}{6}z = 12. \end{cases}$$

$$7. \begin{cases} 8x - 5y + 2z = 53, \\ x + y - z = 9, \\ 13x - 9y + 3z = 71. \end{cases}$$

$$16. \begin{cases} \frac{1}{6}x - \frac{1}{5}y + \frac{1}{4}z = 3, \\ \frac{1}{5}x - \frac{1}{4}y + \frac{1}{3}z = 1, \\ \frac{1}{4}x - \frac{1}{3}y + \frac{1}{2}z = 5. \end{cases}$$

$$8. \begin{cases} x + 3y + 4z = 83, \\ x + y + z = 29, \\ 6x + 8y + 3z = 156. \end{cases}$$

$$17. \begin{cases} \frac{x+y}{3} + 3z = 29, \\ \frac{2x-y}{2} + 2z = 22, \\ 3x - y = 3(z - 1). \end{cases}$$

$$9. \begin{cases} 2x + 3y + 4z = 29, \\ 3x + 2y + 5z = 32, \\ 4x + 3y + 2z = 25. \end{cases}$$

$$10. \begin{cases} 2x - 3y + 4z - v = 4, \\ 4x + 2y - z + 2v = 13, \\ x - y + 2z + 3v = 17, \\ 3x + 2y - z + 4v = 20. \end{cases} \quad 18. \begin{cases} 3x + y - z + 2v = 0, \\ 3y - 2x + z - 4v = 21, \\ x - y + 2z - 3v = 6, \\ 4x + 2y - 3z + v = 12. \end{cases}$$

19. Solve

$$\begin{cases} u+v+x+y-z=5, \\ u+v+x-y+z=7, \\ u+v-x+y+z=9, \\ u-v+x+y+z=11, \\ v-u+x+y+z=13. \end{cases}$$

SOLUTION. — Adding the equations,

$$3u + 3v + 3x + 3y + 3z = 45.$$

Dividing by 3,

$$u + v + x + y + z = 15.$$

Subtracting each of the given equations from this equation,

$$2z = 10, \quad 2y = 8, \quad 2x = 6, \quad 2v = 4, \quad 2u = 2;$$

$$\therefore z = 5, \quad y = 4, \quad x = 3, \quad v = 2, \quad u = 1.$$

Solve the following:

20.
$$\begin{cases} x + y = 9, \\ y + z = 7, \\ z + x = 5. \end{cases}$$

24.
$$\begin{cases} x + 3y + z = 14, \\ x + y + 3z = 16, \\ 3x + y + z = 20. \end{cases}$$

21.
$$\begin{cases} v + x + y = 15, \\ x + y + z = 18, \\ y + z + v = 17, \\ z + v + x = 16. \end{cases}$$

25.
$$\begin{cases} y + z + v - x = 22, \\ z + v + x - y = 18, \\ v + x + y - z = 14, \\ x + y + z - v = 10. \end{cases}$$

22.
$$\begin{cases} \frac{1}{x} + \frac{1}{y} = 6, \\ \frac{1}{y} + \frac{1}{z} = 10, \\ \frac{1}{z} + \frac{1}{x} = 8. \end{cases}$$

26.
$$\begin{cases} \frac{1}{x} + \frac{1}{y} - 1 = 0, \\ \frac{1}{y} + \frac{1}{z} + 3 = 0, \\ \frac{1}{z} + \frac{1}{x} - 2 = 0. \end{cases}$$

23.
$$\begin{cases} \frac{xy}{x+y} = \frac{1}{5}, \\ \frac{yz}{y+z} = \frac{1}{6}, \\ \frac{zx}{z+x} = \frac{1}{7}. \end{cases}$$

27.
$$\begin{cases} \frac{xy}{x+y} = \frac{1}{a}, \\ \frac{yz}{y+z} = \frac{1}{b}, \\ \frac{zx}{z+x} = \frac{1}{c}. \end{cases}$$

SUGGESTION. — If $\frac{xy}{x+y} = \frac{1}{5}$, $\frac{x+y}{xy} = \frac{5}{1}$; whence, $\frac{1}{y} + \frac{1}{x} = 5$.

$$28. \text{ Solve } \begin{cases} \frac{a}{x} + \frac{b}{y} + \frac{c}{z} = a, & (1) \\ bzx - cxy + ayz = bxyz, & (2) \\ \frac{a}{x} - \frac{b}{y} - \frac{c}{z} = c. & (3) \end{cases}$$

SOLUTION

$$(1) + (3), \quad \frac{2a}{x} = a + c. \\ \therefore x = \frac{2a}{a+c} \quad (4)$$

$$(2) \div xyz, \quad \frac{b}{y} - \frac{c}{z} + \frac{a}{x} = b. \quad (5)$$

$$(5) - (3), \quad \frac{2b}{y} = b - c. \\ \therefore y = \frac{2b}{b-c}.$$

Substituting the values of x and y in (1), and solving,

$$z = \frac{2c}{a-b}.$$

$$29. \begin{cases} axy - x - y = 0, \\ bzx - z - x = 0, \\ cyz - y - z = 0. \end{cases} \quad 33. \begin{cases} abxyz + cxy - ayz - bzx = 0, \\ bcxyz + ayz - bzx - cxy = 0, \\ caxyz + bzx - cxy - ayz = 0. \end{cases}$$

$$30. \begin{cases} x + y - z = 0, \\ x - y = 2b, \\ x + z = 3a + b. \end{cases} \quad 34. \begin{cases} x + y + z = a + b + c, \\ x + 2y + 3z = b + 2c, \\ x + 3y + 4z = b + 3c. \end{cases}$$

$$31. \begin{cases} v + x = 2a, \\ x + y = 2a - z, \\ y + z = a + b, \\ v - z = a + c. \end{cases} \quad 35. \begin{cases} v + x + y = a + 2b + c, \\ x + y + z = 3b, \\ y + z + v = a + b, \\ z + v + x = a + 3b - c. \end{cases}$$

$$32. \begin{cases} y + z - 3x = 2a, \\ z + x - 3y = 2b, \\ x + y - 3z = 2c, \\ 2x + 2y + v = 0. \end{cases} \quad 36. \begin{cases} ax + by + cz = 3, \\ x + y = \frac{a+b}{ab}, \\ y + z = \frac{b+c}{bc}. \end{cases}$$

PROBLEMS

215. 1. Three men bought grain at the same prices. A paid \$4.80 for 2 bushels of rye, 3 bushels of wheat, and 4 bushels of oats; B paid \$6.40 for 3 bushels of rye, 5 bushels of wheat, and 2 bushels of oats; and C paid \$5.30 for 2 bushels of rye, 4 bushels of wheat, and 3 bushels of oats. What was the price of each?

2. A dealer was asked his price for 10 bushels of wheat, corn, and rye. He replied, "For 5 of wheat, 2 of corn, and 3 of rye, \$6.60; for 2 of wheat, 3 of corn, and 5 of rye, \$5.80; and for 3 of wheat, 5 of corn, and 2 of rye, \$5.60." What prices had he in mind?

3. Divide 90 into three parts such that the sum of $\frac{1}{2}$ of the first, $\frac{1}{3}$ of the second, and $\frac{1}{4}$ of the third shall be 30; and the first shall be twice the third diminished by twice the second.

4. There are three numbers such that the sum of $\frac{1}{2}$ of the first, $\frac{1}{3}$ of the second, and $\frac{1}{4}$ of the third is 12; of $\frac{1}{3}$ of the first, $\frac{1}{4}$ of the second, and $\frac{1}{5}$ of the third is 9, and the sum of the numbers is 38. What are the numbers?

5. There are three numbers whose sum is 72. If the sum of the first two is divided by the third, the quotient is $1\frac{2}{3}$; and if the third is subtracted from twice the first, the remainder will be $\frac{1}{4}$ of the second. Find the numbers.

6. A and B can do a piece of work in 10 days; A and C can do it in 8 days; and B and C can do it in 12 days. How long will it take each to do it alone?

7. A certain number is expressed by three digits whose sum is 14. If 693 is added to the number, the digits will appear in reverse order. If the units' digit is equal to the tens' digit increased by 6, what is the number?

8. The third digit of a number of three digits is as much larger than the second digit as the second is larger than the first. If the number is divided by the sum of its digits, the quotient is 15. What is the number, if the order of its digits may be reversed by adding 396?

9. Find three numbers such that the first increased by $\frac{1}{3}$ of the sum of the other two shall be 36; the second increased by $\frac{1}{3}$ of the sum of the other two shall be 40; and the third increased by $\frac{1}{3}$ of the sum of the other two shall be 44.

10. Divide 800 into three parts such that the sum of the first, $\frac{1}{2}$ of the second, and $\frac{2}{3}$ of the third shall be 400; and the sum of the second, $\frac{2}{3}$ of the first, and $\frac{1}{2}$ of the third shall be 400.

11. Three cities, A, B, and C, connected by straight roads, are situated at the vertices of a triangle. From A to B by way of C is 130 miles; from B to C by way of A is 110 miles; and from C to A by way of B is 140 miles. How far apart are the cities?

12. Find three numbers such that the first with $\frac{1}{3}$ of the sum of the second and third is 340; the second with $\frac{1}{3}$ of the sum of the first and third is 600; and the third with $\frac{1}{3}$ of the remainder when the first is subtracted from the second is 450.

13. A merchant has three kinds of tea. He can sell 2 pounds of the first kind, 3 of the second, and 4 of the third for \$4.70; or 4 of the first, 3 of the second, and 2 of the third for \$4.30. If a pound of the third kind is worth 5 cents more than $\frac{3}{4}$ of a pound of the first kind and $\frac{1}{2}$ of a pound of the second kind, what is the value of 1 pound of each kind?

14. A, B, and C have certain sums of money. If A gives B \$100, they will have the same amount; if A gives C \$100, C will have twice as much as A; and if B gives C \$100, C will have 4 times as much as B. What sum has each?

15. A quantity of water sufficient to fill three jars of different sizes will fill the smallest jar 4 times; the largest jar twice with 4 gallons to spare; or the second jar 3 times with 2 gallons to spare. What is the capacity of each jar?

16. A gave to B and C as much as each of them had; B then gave to A and C as much as each of them had; and C then gave to A and B as much as each of them had, after which each had \$8. How much had each at first?

17. Three boys, A, B, and C, each had a bag of nuts. After each boy had given each of the others $\frac{1}{3}$ of the nuts in his bag, they counted and found that A had 740, B 580, and C 380. How many had each at first?

INVOLUTION

216. 1. How many times is a number used as a factor in producing its second power? its third power? its fourth power? its fifth power? its n th power, when n is a positive integer?

2. What is the meaning of 2^3 ? of $(-2)^3$? of a^3 ? of $(ax)^3$? of x^n , when n is a positive integer?

3. What sign has $(+a)^3$? $(+a)^4$? $(+a)^n$, or any power of a ? What sign has any power of a positive number?

4. What sign has $(-a)^3$? $(-a)^4$? $(-a)^5$? $(-a)^n$?

What sign have the even powers of a negative number? What sign have the odd powers?

5. What is the fourth power of a^2 ? of a^3 ? of a^{10} ? of a^n , when n is a positive integer? What are the fifth powers of these numbers? the sixth powers? the m th powers, when m is a positive integer?

6. How does 8^3 compare in value with $2^3 \times 4^3$? with $2^3 \times 2^2 \times 2^2$? 3^3 with $6^2 \div 2^2$? 5^3 with $10^3 \div 2^3$?

217. The process of finding any required power of an expression is called **Involution**.

218. PRINCIPLES.—1. **Law of Signs.**—*All powers of a positive number are positive; even powers of a negative number are positive, and odd powers are negative.*

2. **Law of Exponents.**—*The exponent of a power of a number is equal to the exponent of the number multiplied by the exponent of the power to which the number is to be raised.*

3. *Any power of a product is equal to the product of its factors each raised to that power.*

4. *Any power of the quotient of two numbers is equal to the quotient of the numbers each raised to that power.*

The above principles may be established as follows :

PRINCIPLE 1 follows directly from the law of signs for multiplication.

PRINCIPLE 2. When m and n are positive integers,

$$\begin{aligned}\S 24, \quad (a^m)^n &= a^m \times a^m \times a^m \dots \text{to } n \text{ factors} \\ &= a^{m+m+m+\dots \text{to } n \text{ terms}} \\ &= a^{mn}.\end{aligned}$$

PRINCIPLE 3. When n is a positive integer,

$$\begin{aligned}\S 24, \quad (ab)^n &= ab \times ab \times ab \dots \text{to } n \text{ factors} \\ \S 83, \quad &= (aaa \dots \text{to } n \text{ factors}) (bbb \dots \text{to } n \text{ factors}) \\ &= a^n b^n.\end{aligned}$$

PRINCIPLE 4. When n is a positive integer,

$$\begin{aligned}\S 24, \quad \left(\frac{a}{b}\right)^n &= \frac{a}{b} \times \frac{a}{b} \times \frac{a}{b} \dots \text{to } n \text{ factors} \\ \S 180, \quad &= \frac{aaa \dots \text{to } n \text{ factors}}{bbb \dots \text{to } n \text{ factors}} \\ &= \frac{a^n}{b^n}.\end{aligned}$$

219. Involution of monomials.

EXAMPLES

1. What is the third power of $4a^2b$?

$$\text{SOLUTION.} \quad (4a^2b)^3 = 4a^2b \times 4a^2b \times 4a^2b = 64a^6b^3.$$

2. What is the fifth power of $-2ab^2$?

$$\begin{aligned}\text{SOLUTION.} \quad (-2ab^2)^5 &= -2ab^2 \times -2ab^2 \times -2ab^2 \times -2ab^2 \times -2ab^2 \\ &= -32a^5b^{10}.\end{aligned}$$

To raise an integral term to any power :

RULE. — *Raise the numerical coefficient to the required power and annex to it each letter with an exponent equal to the product of its exponent by the exponent of the required power.*

Prefix the sign + to any power of a positive number or to an even power of a negative number ; the sign - to an odd power of a negative number.

To raise a fraction to any power:

RULE. — *Raise both numerator and denominator to the required power and prefix the proper sign to the result.*

Raise to the power indicated:

- | | |
|----------------------------|---------------------------------|
| 3. $(ab^2c^3)^2$. | 15. $(abcx)^m$. |
| 4. $(a^3b^2c)^4$. | 16. $(2e^2x^6)^6$. |
| 5. $(2a^2c)^3$. | 17. $(3bc)^n$. |
| 6. $(7a^2m^5)^3$. | 18. $(2a^2x^3)^n$. |
| 7. $(-1)^3$. | 19. $(-1)^{99}$. |
| 8. $(-ab)^2$. | 20. $(-1)^{300}$. |
| 9. $(-3c)^3$. | 21. $(-1)^{2n}$. |
| 10. $(-10x^2)^3$. | 22. $(-b)^{2n+1}$. |
| 11. $(-6a^2x^3)^2$. | 23. $(-b^2)^{2n+1}$. |
| 12. $(-4c^2y^5)^3$. | 24. $(-a^2b^4c^n \cdot 1d)^2$. |
| 13. $(-2l^4m^5d^2)^3$. | 25. $(-a^{2n}y^{3p}z^4)^5$. |
| 14. $(-a^2x^ny^{n-1})^2$. | 26. $(-a^{n-1}b^{n-2}c)^3$. |

27. What is the square of $-\frac{5a^3x^2}{7b^2c}$?

SOLUTION

$$\left(-\frac{5a^3x^2}{7b^2c}\right)^2 = -\frac{5a^3x^2}{7b^2c} \times -\frac{5a^3x^2}{7b^2c} = +\frac{25a^6x^4}{49b^4c^2}.$$

Raise to the power indicated:

- | | | |
|---|--|---|
| 28. $\left(\frac{1}{4b}\right)^2$. | 33. $\left(-\frac{5}{ab}\right)^2$. | 38. $\left(-\frac{2a}{b}\right)^6$. |
| 29. $\left(\frac{2x}{7y}\right)^2$. | 34. $\left(-\frac{2}{3x}\right)^4$. | 39. $\left(-\frac{b^3c^2}{a^2x}\right)^2$. |
| 30. $\left(\frac{3x^2}{10y^3}\right)^2$. | 35. $\left(-\frac{3x}{2y}\right)^3$. | 40. $\left(\frac{a^2b^3}{xy^4}\right)^n$. |
| 31. $\left(\frac{2x^2}{3y}\right)^3$. | 36. $\left(-\frac{2a}{x^2y}\right)^5$. | 41. $\left(\frac{a^{n-1}b}{x^{2b}y^a}\right)^n$. |
| 32. $\left(\frac{a^n}{2b^{n-1}}\right)^3$. | 37. $\left(-\frac{x^ny^n}{a^2}\right)^7$. | 42. $\left(\frac{a^{n-1}c^n}{x^{m+n-1}}\right)^n$. |

220. Involution of polynomials.

$$\S 91, \quad (a + b)^2 = a^2 + 2ab + b^2.$$

$$\S 93, \quad (a - b)^2 = a^2 - 2ab + b^2.$$

$$\S 95, \quad (a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2ac + 2bc.$$

Raise the following to the second power:

- | | | |
|---------------------|-----------------|----------------------------|
| 1. $2a + b.$ | 5. $3x - 4y^2.$ | 9. $a - b + x - y.$ |
| 2. $2a - b.$ | 6. $5m^4 - 11.$ | 10. $a^m + x^n - y^{n+1}.$ |
| 3. $a^m - 3b^n.$ | 7. $1 - 3abc.$ | 11. $2a + 3b - 4c.$ |
| 4. $a^2 - 2x^{2n}.$ | 8. $4x^4 + 5.$ | 12. $5a^2 - 1 + 4n^3.$ |

Raise to the required power by multiplication:

- | | | |
|------------------|------------------|------------------|
| 13. $(x + y)^3.$ | 15. $(x + y)^4.$ | 17. $(x + y)^5.$ |
| 14. $(x - y)^3.$ | 16. $(x - y)^4.$ | 18. $(x - y)^5.$ |

221. Involution of binomials by the Binomial Theorem.

By multiplication,

$$(a + x)^3 = a^3 + 3a^2x + 3ax^2 + x^3.$$

$$(a - x)^3 = a^3 - 3a^2x + 3ax^2 - x^3.$$

$$(a + x)^4 = a^4 + 4a^3x + 6a^2x^2 + 4ax^3 + x^4.$$

$$(a - x)^4 = a^4 - 4a^3x + 6a^2x^2 - 4ax^3 + x^4.$$

$$(a + x)^5 = a^5 + 5a^4x + 10a^3x^2 + 10a^2x^3 + 5ax^4 + x^5.$$

$$(a - x)^5 = a^5 - 5a^4x + 10a^3x^2 - 10a^2x^3 + 5ax^4 - x^5.$$

Examine carefully the above powers of $(a + x)$ and $(a - x)$.

1. How does the number of terms in a power of a binomial compare with the exponent of the binomial?

2. What terms of the power contain the first term of the binomial? the second term of the binomial? both terms?

3. What is the exponent of the first term of the binomial in the first term of the power? in the second? in the third, etc.?

4. What is the exponent of the second term of the binomial in the second term of the power? in the third? in the fourth, etc.?

5. What is the coefficient of the first term of the power? How does the coefficient of the second term compare with the exponent of the binomial?

6. If the coefficient of any term is multiplied by the exponent of the first term of the binomial found in that term, and the product is divided by the number of the term, how does the quotient compare with the coefficient of the succeeding term?

7. What are the signs of the terms in any power of $(a + b)$? What terms are negative in any power of $(a - b)$?

222. PRINCIPLES.—1. *The number of terms in a positive integral power of a binomial is one greater than the index of the required power.*

2. *The first term of the power contains only the first term of the binomial; the last term of the power, only the second term of the binomial; all other terms of the power contain as factors both terms of the binomial.*

3. *The exponent of the first term of the binomial in the first term of the power is the same as the index of the required power, and it decreases 1 in each succeeding term. The exponent of the second term of the binomial in the second term of the power is 1, and it increases 1 in each succeeding term.*

4. *The coefficient of the first term of the power is 1. The coefficient of the second term is the same as the index of the required power.*

5. *The coefficient of any term may be found by multiplying the coefficient of the preceding term by the exponent of the first term of the binomial found in that term, and then dividing the result by the number of the term.*

6. *If both terms of the binomial are positive, all the terms of any power of the binomial will be positive.*

7. *If the second term of the binomial is negative and the first term positive, the terms of any power of the binomial will be alternately positive and negative.*

EXAMPLES

1. Find the fifth power of $(b - y)$ by the binomial theorem.

SOLUTION

Letters and exponents,	b^5	b^4y	b^3y^2	b^2y^3	by^4	y^5
Coefficients,	1	5	10	10	5	1
Signs,	+	-	+	-	+	-
Combined,	$b^5 - 5b^4y + 10b^3y^2 - 10b^2y^3 + 5by^4 - y^5$					

In every term of a power of a binomial the sum of the exponents of the terms of the binomial is equal to the index of the required power.

Expand :

- | | | |
|-------------------|-------------------|---------------------|
| 2. $(x + y)^5$. | 13. $(c - n)^6$. | 24. $(x - 2)^3$. |
| 3. $(m + n)^5$. | 14. $(x - a)^7$. | 25. $(x + r)^7$. |
| 4. $(m - n)^5$. | 15. $(d - y)^8$. | 26. $(b - c)^7$. |
| 5. $(a - c)^3$. | 16. $(b + y)^6$. | 27. $(p + q)^5$. |
| 6. $(a + b)^3$. | 17. $(m + n)^4$. | 28. $(a - b)^5$. |
| 7. $(b + d)^4$. | 18. $(p - q)^4$. | 29. $(a + bc)^4$. |
| 8. $(q - r)^5$. | 19. $(s + t)^3$. | 30. $(ab - c)^4$. |
| 9. $(c + d)^7$. | 20. $(x + 2)^3$. | 31. $(m - pn)^4$. |
| 10. $(x + y)^8$. | 21. $(a + 3)^3$. | 32. $(m - an)^4$. |
| 11. $(x - y)^3$. | 22. $(x + 4)^3$. | 33. $(ax - by)^3$. |
| 12. $(x - y)^4$. | 23. $(x + 5)^3$. | 34. $(ax - by)^5$. |

35. Expand $(2b^2 - 3y)^4$.

SOLUTION

Let $2b^2 = a$, and $3y = x$.

Then, $2b^2 - 3y = a - x$,

and $(2b^2 - 3y)^4 = (a - x)^4$

Restoring values,

$$\begin{aligned}
 &= a^4 - 4a^3x + 6a^2x^2 - 4ax^3 + x^4 \\
 &= (2b^2)^4 - 4(2b^2)^3(3y) + 6(2b^2)^2(3y)^2 \\
 &\quad - 4(2b^2)(3y)^3 + (3y)^4 \\
 &= 16b^8 - 96b^6y + 216b^4y^2 - 216b^2y^3 + 81y^4.
 \end{aligned}$$

36. Expand $(1 + x^2)^3$.

SOLUTION

$$\begin{aligned}(1 + x^2)^3 &= 1^3 + 3(1)^2(x^2) + 3(1)(x^2)^2 + (x^2)^3 \\ &= 1 + 3x^2 + 3x^4 + x^6.\end{aligned}$$

Expand:

- | | | |
|----------------------|------------------------|---|
| 37. $(x + 2y)^4$. | 41. $(1 - 3x^2)^4$. | 45. $(1 - x)^7$. |
| 38. $(2x - y)^3$. | 42. $(5x^2 - ab)^3$. | 46. $(1 - 2x)^6$. |
| 39. $(2x - 5)^3$. | 43. $(1 + a^2b^2)^4$. | 47. $(x - \frac{1}{2})^6$. |
| 40. $(x^2 - 10)^4$. | 44. $(2ax - b)^5$. | 48. $(\frac{1}{2}x - \frac{1}{3}y)^4$. |

Expand:

- | | | |
|---------------------------------------|--|-------------------------------|
| 49. $(2a + \frac{1}{2})^5$. | 52. $(3a^2 + \frac{b}{6})^3$. | 55. $(\frac{1}{2x} - 2x)^5$. |
| 50. $(\frac{x}{y} - \frac{y}{x})^4$. | 53. $(1 + \frac{3x}{2})^5$. | 56. $(\frac{1}{a} - a)^6$. |
| 51. $(\frac{x}{y} - \frac{y}{x})^6$. | 54. $(\frac{3}{5} + \frac{5x}{3})^4$. | 57. $(x + \frac{1}{x})^7$. |

58. Expand $(a - b - c)^3$.

SOLUTION

 $(a - b - c)^3 = (\overline{a - b - c})^3$, a binomial form.

$$\begin{aligned}(\overline{a - b - c})^3 &= (a - b)^3 - 3(a - b)^2c + 3(a - b)c^2 - c^3 \\ &= a^3 - 3a^2b + 3ab^2 - b^3 - 3c(a^2 - 2ab + b^2) + 3ac^2 - 3bc^2 - c^3 \\ &= a^3 - 3a^2b + 3ab^2 - b^3 - 3a^2c + 6abc - 3b^2c + 3ac^2 - 3bc^2 - c^3.\end{aligned}$$

59. Expand $(a + b - c - d)^3$.SUGGESTION. $(a + b - c - d)^3 = (\overline{a + b - c + d})^3$, a binomial form.

Expand:

- | | |
|-----------------------|---------------------------|
| 60. $(a + x - y)^3$. | 66. $(a + 2b - 3c)^3$. |
| 61. $(a - m - n)^3$. | 67. $(a + b + x + y)^3$. |
| 62. $(a - x + y)^3$. | 68. $(a + b - x - y)^3$. |
| 63. $(a - x - y)^3$. | 69. $(a - b + x - y)^3$. |
| 64. $(a + x + 2)^3$. | 70. $(a - b - x + y)^3$. |
| 65. $(a - x - 2)^3$. | 71. $(a - b - x - y)^3$. |

EVOLUTION

223. 1. Of what two equal numbers is 16 the product? What is the square root of 16? Since 16 is equal also to -4×-4 , what other square root may 16 have? What is the square root of 25? of 64? What is the fourth root of 16? of 81?

2. What is the sign of an *even* root of a *positive* number?

3. Can the square root of -16 be found? of -25 ? of -64 ? the fourth root of -16 ? of -81 ? Can an even root of any negative number be found?

4. What is the cube root of 8? of 27? of 64? of -8 ? of -27 ? of -64 ? What is the fifth root of 32? of -32 ?

5. How does the sign of an *odd* root of a number compare with the sign of the number?

6. Since $a^3 = a^2 \times a^2 \times a^2$, what power of a^2 is a^6 ? What is the cube root of a^6 ? of a^9 ? of a^{12} ? How is the exponent of a in the cube, or third, root of any power of a found? What is the fourth root of a^8 ? of a^{12} ?

7. How is the exponent of a root of a power obtained from the index of the power and the index of the root?

8. How does $\sqrt{4 \times 25}$ compare in value with $\sqrt{4} \times \sqrt{25}$? $\sqrt{4 \times 9}$ with $\sqrt{4} \times \sqrt{9}$? $\sqrt[3]{8 \times 1000}$ with $\sqrt[3]{8} \times \sqrt[3]{1000}$? In each case how does the root of the product compare in value with the product of the roots of the factors?

9. How does $\sqrt{100 \div 4}$ compare in value with $\sqrt{100} \div \sqrt{4}$? $\sqrt{36 \div 9}$ with $\sqrt{36} \div \sqrt{9}$? $\sqrt[3]{64 \div 8}$ with $\sqrt[3]{64} \div \sqrt[3]{8}$? In each case how does the root of the quotient compare in value with the quotient of the roots of the dividend and the divisor?

224. The process of finding any required root of an expression is called **Evolution**.

225. Since the product of two numbers having like signs is positive, *every positive number has two square roots, numerically equal, but with opposite signs*. It will be seen later that every number has two square roots, three cube roots, four fourth roots, five fifth roots, and, in general, q q th roots. Of these roots the positive roots of positive numbers and the negative odd roots of negative numbers are called **Principal Roots**.

$\sqrt{25} = +5$ or -5 , and $+5$ is the principal square root. $\sqrt[4]{16} = +2$ or -2 or, as will be seen later, $+\sqrt{-4}$ or $-\sqrt{-4}$, but $+2$ is the principal root. The principal cube root of 8 is $+2$ and of -8 is -2 .

226. A number that is or can be expressed as an integer or as a fraction with integral terms is called a **Rational Number**.

a , 3, $5\frac{1}{2}$, $a^2 + b^2$, $\sqrt{25}$, and $.33\bar{3}$ are rational numbers.

A number that cannot be expressed as an integer or as a fraction with integral terms is called an **Irrational Number**.

When the indicated root of a number cannot be exactly obtained, the root is irrational.

The indicated roots $\sqrt{2}$, $\sqrt[3]{4}$, $\sqrt{a^2 + b^2}$, $\sqrt[3]{x^2}$, $\sqrt[4]{x^5}$, and, in general, the q th root of a number that is not the q th power of some rational number, are irrational numbers.

227. A rational arithmetical number is called a **Commensurable Number**, and an irrational arithmetical number is called an **Incommensurable Number**.

2 , $\frac{3}{4}$, $.54$, and $.66\bar{6}$ are commensurable, but $\sqrt{2}$ is incommensurable.

In algebra, commensurable and incommensurable numbers may be either positive or negative.

The terms rational and irrational applied to algebraic numbers relate to their forms, while the terms commensurable and incommensurable relate to their arithmetical values.

$3a$, $a + b$, $x - 3$, x^3 , are rational but not necessarily commensurable. For a may represent $\sqrt{2}$, b may represent $\sqrt[3]{5}$, etc. Again, \sqrt{x} is irrational, but if $x = 16$, \sqrt{x} is commensurable.

Incommensurable numbers obey the Commutative, Associative, and Distributive Laws, but the proof is too complicated to be given here.

228. Since the n th power of a number is the product of n equal factors, and since one of these factors is a root of the power, it follows that $(\sqrt[n]{a})^n = a$, and if the principal root is meant, $\sqrt[n]{a^n} = a$, when n is a positive integer.

In the statement of the following principles and in Ax. 7 the term root means principal root.

229. PRINCIPLES. — 1. Law of Signs. — *An odd root of a number has the same sign as the number ; an even root of a positive number is positive ; an even root of a negative number is impossible, or imaginary.*

2. Law of Exponents. — *The exponent of any root of a number is equal to the exponent of the given number divided by the index of the root.*

3. *Any root of a product is equal to the product of that root of each of the factors.*

4. *Any root of the quotient of two numbers is equal to the quotient of that root of each of the numbers.*

Even roots of negative numbers will be discussed later.

The above principles may be established as follows :

PRINCIPLE 1 follows from the Law of Signs for multiplication.

PRINCIPLE 2. When m and n are positive integers,

§ 218, Prin. 2,

$$a^{mn} = (a^m)^n.$$

Taking the n th root, § 228,

$$\sqrt[n]{a^{mn}} = a^m ;$$

$$\therefore \sqrt[n]{a^{mn}} = a^{mn \div n} = a^m.$$

Thus,

$$\sqrt[5]{a^{10}} = a^2.$$

PRINCIPLE 3. When n is a positive integer,

$$\text{§ 228, § 218, Prin. 3, } ab = (\sqrt[n]{a})^n \times (\sqrt[n]{b})^n = (\sqrt[n]{a} \times \sqrt[n]{b})^n.$$

Taking the n th root,

$$\sqrt[n]{ab} = \sqrt[n]{a} \times \sqrt[n]{b}.$$

Thus,

$$\sqrt[3]{27 a^9} = \sqrt[3]{27} \times \sqrt[3]{a^9} = 3 a^3.$$

PRINCIPLE 4. When n is a positive integer,

§ 228, § 218, Prin. 4,

$$\frac{a}{b} = \frac{(\sqrt[n]{a})^n}{(\sqrt[n]{b})^n} = \left(\frac{\sqrt[n]{a}}{\sqrt[n]{b}} \right)^n.$$

Taking the n th root,

$$\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}.$$

230. Evolution of monomials.

EXAMPLES

1. What is the square root of $36 a^6 b^2$?

SOLUTION. — Since, in squaring a monomial, § 218, the coefficient is squared and the exponents of the letters are multiplied by 2, to extract the square root, the square root of the coefficient must be found, and to it must be annexed the letters each with its exponent divided by 2.

The square root of 36 is 6, and the square root of the literal factors is $a^3 b$. Therefore, the *principal* square root of $36 a^6 b^2$ is $6 a^3 b$.

The square root may also be $-6 a^3 b$, since $-6 a^3 b \times -6 a^3 b = 36 a^6 b^2$.

$$\therefore \sqrt{36 a^6 b^2} = \pm 6 a^3 b.$$

2. What is the cube root of $-125 x^6 y^{21}$?

SOLUTION. $\sqrt[3]{-125 x^6 y^{21}} = -5 x^2 y^7.$

To find the root of an integral term:

RULE. — *Extract the required root of the numerical coefficient, annex to it the letters each with its exponent divided by the index of the root, sought, and prefix the proper sign to the result.*

To find the root of a fractional term:

RULE. — *Find the required root of both numerator and denominator and prefix the proper sign to the resulting fraction.*

Find the indicated root:

3. $\sqrt[3]{a^9 b^9 c^{15}}.$

4. $\sqrt{a^6 b^{16} c^{14}}.$

5. $\sqrt[5]{a^{10} x^5 y^{30}}.$

6. $\sqrt[4]{a^{4n} b^8 c^{12}}.$

7. $\sqrt{x^{4n} y^8 z^{2m}}.$

8. $\sqrt[3]{-8 a^6 b^{15}}.$

9. $\sqrt[5]{-32 x^{10} y^{10}}.$

10. $\sqrt{16 x^4 y^2}.$

11. $\sqrt[7]{-a^{21} b^{35} c^{14}}.$

12. $\sqrt[5]{-243 y^{25}}.$

13. $\sqrt[4]{16 m^4 n^8}.$

14. $\sqrt[7]{\frac{(x-y)^{14}}{128 x^{14}}}.$

15. $\sqrt[5]{\frac{-32 a^5 x^{10}}{243 y^{15}}}.$

16. $\sqrt[8]{\frac{256 x^8}{6561}}.$

17. $\sqrt[3]{-\frac{125 x^{12} y^{12}}{1728 c^3}}.$

18. $\sqrt[2n]{\frac{a^{2n^2-2n} b^{4n}}{x^{4n} y^{2n}}}.$

19. $\sqrt[n]{\frac{b^{4n} c^n x^{2n}}{2^n x^{2n} y^{2n}}}.$

231. To extract the square root of a polynomial.

1. Since $a^2 + 2ab + b^2$ is the square of $(a + b)$, what is the square root of $a^2 + 2ab + b^2$?

2. How may the first term of the square root be found from $a^2 + 2ab + b^2$?

3. How may the second term of the square root be found from $2ab$, the second term of the power?

4. What are the factors of $a^2 + 2ab + b^2$?

5. Since $2ab + b^2$ is equal to $b(2a + b)$, what are the factors of the last two terms of the square of a binomial?

6. By what divisor, then, must the last two terms of $a^2 + 2ab + b^2$ be divided so that the quotient may be the second term of the square root?

EXAMPLES

1. Find the process for extracting the square root of $a^2 + 2ab + b^2$.

PROCESS

$$\begin{array}{r} a^2 + 2ab + b^2 \quad | \quad a + b \\ \underline{a^2} \end{array}$$

Trial divisor, $2a$

Complete divisor, $2a + b$

$$\begin{array}{r} 2ab + b^2 \\ \underline{2ab + b^2} \end{array}$$

EXPLANATION. — Since $a^2 + 2ab + b^2$ is the square of $(a + b)$, we know that the square root of $a^2 + 2ab + b^2$ is $a + b$.

Since the first term of the root is a , it may be found by taking the square root of a^2 , the first term of the power. Subtracting a^2 , there is a remainder of $2ab + b^2$.

The second term of the root is known to be b , and that may be found by dividing the first term of the remainder by twice the part of the root already found. This divisor is called a *trial* divisor.

Since $2ab + b^2$ is equal to $b(2a + b)$, the complete divisor which multiplied by b produces the remainder $2ab + b^2$ is $2a + b$; that is, the complete divisor is found by adding the second term of the root to twice the root already found.

Multiplying the complete divisor by the second term of the root and subtracting, there is no remainder, consequently, $a + b$ is the required root.

2. Find the square root of $9x^2 - 30xy + 25y^2$.

PROCESS

$$\begin{array}{r}
 9x^2 - 30xy + 25y^2 \quad | \quad 3x - 5y \\
 \underline{9x^2} \\
 6x \quad | \quad -30xy + 25y^2 \\
 \underline{6x - 5y} \quad | \quad -30xy + 25y^2
 \end{array}$$

Since, in squaring $a + b + c$, $a + b$ may be represented by x , and the square of the number by $x^2 + 2xc + c^2$, it is obvious that the square root of a number whose root consists of *more than two terms* may be extracted in the same way as in Example 1, by *considering the terms already found as one term*.

3. Find the square root of $4x^4 + 12x^3 - 3x^2 - 18x + 9$.

PROCESS

$$\begin{array}{r}
 4x^4 + 12x^3 - 3x^2 - 18x + 9 \quad | \quad 2x^2 + 3x - 3 \\
 \underline{4x^4} \\
 4x^2 \quad | \quad 12x^3 - 3x^2 \\
 \underline{4x^2 + 3x} \quad | \quad 12x^3 - 9x^2 \\
 4x^2 + 6x \quad | \quad -12x^2 - 18x + 9 \\
 \underline{4x^2 + 6x - 3} \quad | \quad -12x^2 - 18x + 9
 \end{array}$$

EXPLANATION. — Proceeding as in the previous example, the first two terms of the root are found to be $2x^2 + 3x$.

Considering $(2x^2 + 3x)$ as the first term of the root, the next term of the root is found as the second term of a root is found, by dividing the remainder by twice the part of the root already found. Hence, the trial divisor is $4x^2 + 6x$, and the next term of the root is -3 . Annexing this, as before, to the trial divisor already found, the entire divisor is $2x^2 + 3x - 3$. This multiplied by -3 and the product subtracted from $-12x^2 - 18x + 9$, leaves no remainder. Hence, the square root of the number is $2x^2 + 3x - 3$.

RULE. — *Arrange the terms of the polynomial with reference to the consecutive powers of some letter.*

Extract the square root of the first term, write the result as the first term of the root, and subtract its square from the given polynomial.

Divide the first term of the remainder by twice the root already found, as a trial divisor, and the quotient will be the next term of the root. Write this result in the root, and annex it to the trial divisor to form the complete divisor.

Multiply the complete divisor by this term of the root, and subtract the product from the first remainder.

Find the next term of the root by dividing the first term of the remainder by the first term of the trial divisor.

Form the complete divisor as before and continue in this manner until all the terms of the root are found.

Find the square root of the following :

4. $4x^2 + 12x + 9.$

7. $x^2 + xy + \frac{1}{4}y^2.$

5. $4x^2 + 20x + 25.$

8. $4x^2 - 52x + 169.$

6. $25x^2 + 40x + 16.$

9. $(a+b)^2 - 4(a+b) + 4.$

10. $x^6 + 4x^5 + 2x^4 + 9x^3 - 4x + 4.$

11. $9x^4 - 12x^3 + 10x^2 - 4x + 1.$

12. $x^4 - 6x^3y + 13x^2y^2 - 12xy^3 + 4y^4.$

13. $x^8 + 2a^2x^2 - a^4x^4 - 2a^2x^6 + a^8.$

14. $25x^4 + 4 - 12x - 30x^3 + 29x^2.$

15. $1 + 6x + 15x^2 + 20x^3 + 15x^4 + 6x^5 + x^6.$

16. $1 - 2x + 3x^2 - 4x^3 + 3x^4 - 2x^5 + x^6.$

17. $a^4 - 2a^2b + 2a^2c^2 - 2bc^2 + b^2 + c^4.$

18. $4a^2 - 12ab + 16ac + 9b^2 + 16c^2 - 24bc.$

19. $9x^2 + 25y^2 + 9z^2 - 30xy + 18xz - 30yz.$

20. $x^2 + 2x - 1 - \frac{2}{x} + \frac{1}{x^2}.$

21. $x^4 + x^3 + \frac{13x^2}{20} + \frac{x}{5} + \frac{1}{25}.$

22. $\frac{a^4}{4} + a^3x + \frac{4a^2x^2}{3} + \frac{2ax^3}{3} + \frac{x^4}{9}.$

23. $x^8 + 4x^7 - 2x^6 - 20x^5 - 3x^4 + 32x^3 + 4x^2 - 16x + 4.$

24. Find four terms of the square root of $1 + x$.

SOLUTION

$$\begin{array}{r}
 1 + x \mid 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{16}x^3 \\
 \underline{1} \\
 2 + \frac{1}{2}x \mid x \\
 \underline{x + \frac{1}{4}x^2} \\
 2 + x - \frac{1}{4}x^2 \mid -\frac{1}{4}x^2 \\
 \underline{-\frac{1}{4}x^2 - \frac{1}{8}x^3 + \frac{1}{16}x^4} \\
 2 + x - \frac{1}{4}x^2 + \frac{1}{16}x^3 \mid \frac{1}{8}x^3 - \frac{1}{16}x^4
 \end{array}$$

Find the square root of the following to four terms:

25. $1 - a$.

27. $x^2 - 1$.

29. $y^2 + 3$.

26. $a^2 + 1$.

28. $4 - a$.

30. $a^2 + 2b$.

SQUARE ROOT OF ARITHMETICAL NUMBERS

$1^2 = 1$

$10^2 = 100$

$100^2 = 10000$

$9^2 = 81$

$99^2 = 9801$

$999^2 = 998001$

232. 1. How many figures are required to express the square of a number expressed by 1 figure? 2 figures? 3 figures? 4 figures?

2. How does the number of figures in the square of a number compare with the number of figures in the number?

3. How many figures are there in the square root of a number that is expressed by 4 figures? by 3 figures? by 5 figures? by 6 figures? by 7 figures?

4. How, then, may the number of figures in the square root of a given number be found?

PRINCIPLES. — 1. *The square of a number is expressed by twice as many figures as there are in the number itself, or by one less than twice as many.*

2. *The orders of units in the square root of a number correspond to the number of periods of two figures each into which the number can be separated, beginning at units.*

233. If the number of units expressed by the tens' digit is represented by t and the number of units expressed by the units' digit by u , the square of a number consisting of tens and units will be represented by $(t + u)^2$, or $t^2 + 2tu + u^2$.

Thus, $25 = 2 \text{ tens} + 5 \text{ units, or } 20 + 5 \text{ units,}$
and $25^2 = 20^2 + 2(20 + 5) + 5^2 = 625.$

EXAMPLES

1. What is the square root of 2809?

FIRST PROCESS

$$\begin{array}{r}
 28\text{-}09 \quad | \quad 50 + 3 \\
 t^2 = \quad 25 \quad 00 \\
 \hline
 2t = 100 \quad | \quad 3 \quad 09 \\
 u = \quad 3 \\
 \hline
 2t + u = 103 \quad | \quad 3 \quad 09
 \end{array}$$

EXPLANATION. — According to Prin. 2, § 231, the orders of units in the square root of a number may be determined by separating the number into periods of two figures each, beginning at units. Separating 2809 thus, there are found to be two orders of units in the root; that is, it is composed of tens and units.

Since the square of tens is hundreds, and the hundreds of the power are less than 36, or 6^2 , and more than 25, or 5^2 , the tens' figure of the root must be 5. 5 tens, or 50, squared is 2500, and 2500 subtracted from 2809 leaves 309, which is equal to 2 times the tens \times the units + the units².

Since two times the tens multiplied by the units is much greater than the square of the units, 309 is a little more than 2 times the tens multiplied by the units. Therefore, if 309 is divided by 2 times the tens, or 100, the *trial divisor*, the units are found to be 3. And since the complete divisor is found by adding the units to twice the tens, the complete divisor is $100 + 3$, or 103. This multiplied by 3 gives as a product 309, which subtracted from 309 leaves no remainder. Therefore, the square root of 2809 is 53.

SECOND PROCESS

$$\begin{array}{r}
 28\text{-}09 \quad | \quad 53 \\
 t^2 = \quad 25 \\
 \hline
 2t = 100 \quad | \quad 309 \\
 u = \quad 3 \\
 \hline
 2t + u = 103 \quad | \quad 309
 \end{array}$$

EXPLANATION. — In practice it is usual to place the figures of the same order in a column, and to disregard the ciphers on the right of the products.

Since any number may be regarded as composed of tens and units, the processes given above have a general application.

Thus, $346 = 34 \text{ tens} + 6 \text{ units; } 2377 = 237 \text{ tens} + 7 \text{ units.}$

2. Find the square root of 104976.

SOLUTION

		10-49-76 <u>324</u>
		9
Trial divisor	$= 2 \times 30 = 60$	<u>1 49</u>
Complete divisor	$= 60 + 2 = 62$	<u>1 24</u>
Trial divisor	$= 2 \times 320 = 640$	25 76
Complete divisor	$= 640 + 4 = 644$	<u>25 76</u>

RULE. — *Separate the number into periods of two figures each, beginning at units.*

Find the greatest square in the left-hand period and write its root for the first figure of the required root.

Square this root, subtract the result from the left-hand period, and annex to the remainder the next period for a new dividend.

Double the root already found, with a cipher annexed, for a trial divisor, and by it divide the dividend. The quotient or the quotient diminished will be the second figure of the root. Add to the trial divisor the figure last found, multiply this complete divisor by the figure of the root found, subtract the product from the dividend, and to the remainder annex the next period for the next dividend.

Proceed in this manner until all the periods have been used. The result will be the square root sought.

1. When the number is not a perfect square, annex periods of decimal ciphers and continue the process.

2. Decimals are pointed off from the decimal point toward the right.

3. The square root of a fraction may be found by extracting the square root of both numerator and denominator separately or by reducing it to a decimal and then extracting its root.

Extract the square root of the following:

- | | | |
|-----------|-------------|--------------|
| 3. 529. | 9. 57121. | 15. 125316. |
| 4. 2209. | 10. 42025. | 16. 455625. |
| 5. 4761. | 11. 95481. | 17. 992016. |
| 6. 7921. | 12. 186624. | 18. 2480.04. |
| 7. 17424. | 13. 165649. | 19. 10.9561. |
| 8. 19321. | 14. 134689. | 20. .001225. |

21. 10201.	24. 1332.25.	27. 540.5625.	
22. 95481.	25. 101.0025.	28. 1.018081.	
23. 363609.	26. 111.0916.	29. 13003236.	
30. $\frac{525}{125}$.	32. $\frac{192}{125}$.	34. $\frac{288}{125}$.	36. $\frac{576}{125}$.
31. $\frac{576}{125}$.	33. $\frac{192}{125}$.	35. $\frac{288}{125}$.	37. $\frac{576}{125}$.

Extract the square root to four decimal places :

38. $\frac{3}{4}$.	40. $\frac{5}{8}$.	42. $\frac{5}{8}$.	44. $\frac{7}{8}$.
39. $\frac{4}{5}$.	41. .6.	43. $\frac{2}{3}$.	45. $\frac{5}{16}$.

234. To extract the cube root of a polynomial.

1. Since $a^3 + 3a^2b + 3ab^2 + b^3$ is the cube of $(a + b)$, what is the cube root of $a^3 + 3a^2b + 3ab^2 + b^3$?

2. How may the first term of the root be found from $a^3 + 3a^2b + 3ab^2 + b^3$?

3. How may the second term of the root be found from the second term of the power, $3a^2b$?

4. What are the factors of $3a^2b + 3ab^2 + b^3$?

5. Since $3a^2b + 3ab^2 + b^3$ is equal to $b(3a^2 + 3ab + b^2)$, by what number must the last three terms of $a^3 + 3a^2b + 3ab^2 + b^3$ be divided so that the quotient may be the second term of the cube root?

EXAMPLES

1. Find the process for extracting the cube root of $a^3 + 3a^2b + 3ab^2 + b^3$.

PROCESS

$$\begin{array}{r}
 a^3 + 3a^2b + 3ab^2 + b^3 \quad | \quad a + b \\
 \underline{a^3} \\
 \text{Trial divisor, } 3a^2 \\
 \text{Complete divisor, } 3a^2 + 3ab + b^2 \quad | \quad \begin{array}{l} 3a^2b + 3ab^2 + b^3 \\ 3a^2b + 3ab^2 + b^3 \end{array}
 \end{array}$$

EXPLANATION. — Since $a^3 + 3a^2b + 3ab^2 + b^3$ is the cube of $(a + b)$, we know that the cube root of $a^3 + 3a^2b + 3ab^2 + b^3$ is $a + b$.

Since the first term of the root is a , it may be found by taking the cube root of a^3 , the first term of the power. Subtracting, there is a remainder of $3a^2b + 3ab^2 + b^3$.

The second term of the root is known to be b , and that may be found by dividing the first term of the remainder by 3 times the square of the part of the root already found. This divisor is called a *trial* divisor.

Since $3a^2b + 3ab^2 + b^3$ is equal to $b(3a^2 + 3ab + b^2)$, the complete divisor, which multiplied by b produces the remainder $3a^2b + 3ab^2 + b^3$, is $3a^2 + 3ab + b^2$; that is, the complete divisor is found by adding to the trial divisor 3 times the product of the first and second terms of the root and the square of the second term of the root.

Multiplying the complete divisor by the second term of the root, and subtracting, there is no remainder; consequently, $a + b$ is the required root.

Since, in cubing $a + b + c$, $a + b$ may be expressed by x , its cube will be $x^3 + 3x^2c + 3xc^2 + c^3$. Hence, it is obvious that the cube root of an expression whose root consists of *more than two terms* may be extracted in the same way as in example 1, by *considering the terms already found as one term*.

2. Find the cube root of $b^6 - 3b^4 + 5b^3 - 3b - 1$.

PROCESS

$$\begin{array}{r}
 b^6 - 3b^4 + 5b^3 - 3b - 1 \quad | \quad b^2 - b - 1 \\
 \underline{b^6} \\
 \text{Trial divisor,} \quad 3b^4 \quad | \quad -3b^2 + 5b^3 \\
 \text{Complete divisor,} \quad 3b^4 - 3b^2 + b^2 \quad | \quad -3b^2 + 3b^4 - b^3 \\
 \text{Trial divisor,} \quad 3b^4 - 6b^2 + 3b^2 \quad | \quad -3b^4 + 6b^3 - 3b - 1 \\
 \text{Complete divisor,} \quad 3b^4 - 6b^2 + 3b + 1 \quad | \quad -3b^4 + 6b^3 - 3b - 1
 \end{array}$$

EXPLANATION. — The first two terms are found in the same manner as in the previous example. In finding the next term, $b^2 - b$ is considered as one term, which we square and multiply by 3 for a trial divisor. Dividing the remainder by this trial divisor, the next term of the root is found to be -1 . Adding to the trial divisor 3 times $(b^2 - b)$ multiplied by -1 , and the square of -1 , we obtain the complete divisor. This multiplied by -1 , and the product subtracted from $-3b^4 + 6b^3 - 3b - 1$, leaves no remainder. Hence, the cube root of the polynomial is $b^2 - b - 1$.

RULE. — *Arrange the polynomial with reference to the consecutive powers of some letter.*

Extract the cube root of the first term, write the result as the first term of the root, and subtract its cube from the given polynomial.

Divide the first term of the remainder by 3 times the square of the root already found, as a trial divisor, and the quotient will be the next term of the root.

Add to this trial divisor 3 times the product of the first and second terms of the root, and the square of the second term. The result will be the complete divisor.

Multiply the complete divisor by the last term of the root found, and subtract this product from the dividend.

Find the next term of the root by dividing the first term of the remainder by the first term of the trial divisor.

Form the complete divisor as before, and continue in this manner until all the terms of the root are found.

Find the cube root of

3. $x^3 - 3x^2y + 3xy^2 - y^3$.

4. $m^3 - 9m^2 + 27m - 27$.

5. $8m^3 - 60m^2n + 150mn^2 - 125n^3$.

6. $27x^3 - 189x^2y + 441xy^2 - 343y^3$.

7. $125a^3 + 675a^2x + 1215ax^2 + 729x^3$.

8. $1000p^6 - 300p^4q + 30p^2q^2 - q^3$.

9. $m^6 + 6m^5 + 15m^4 + 20m^3 + 15m^2 + 6m + 1$.

10. $x^6 - 6x^5 + 15x^4 - 20x^3 + 15x^2 - 6x + 1$.

11. $x^6 + 3x^5 + 9x^4 + 13x^3 + 18x^2 + 12x + 8$.

12. $x^6 + 12x^5 + 63x^4 + 184x^3 + 315x^2 + 300x + 125$.

13. $x^6 + 6x^5 - 18x^4 - 1000 + 180x^2 - 112x^3 + 600x$.

14. $x^3 - 12x^2 + 54x - 112 + \frac{108}{x} - \frac{48}{x^2} + \frac{8}{x^3}$.

15. $1 - 6a + 21a^2 - 44a^3 + 63a^4 - 54a^5 + 27a^6$.

16. $64 - 144p + 156p^2 - 99p^3 + 39p^4 - 9p^5 + p^6$.

17. $\frac{a^3b^3x^9}{c^3} - \frac{c^3x^6}{b^3} + \frac{3acx^7}{b} - \frac{3a^2bx^5}{c}$.

18. $x^6 + 15x^2 + \frac{15}{x^2} + 20 + \frac{6}{x^4} + \frac{1}{x^6} + 6x^4$.

CUBE ROOT OF ARITHMETICAL NUMBERS

$1^3 = 1.$

$10^3 = 1000.$

$100^3 = 1000000.$

$3^3 = 27.$

$30^3 = 27000.$

$300^3 = 27000000.$

$9^3 = 729.$

$99^3 = 970299.$

$999^3 = 997002999.$

235. 1. How many figures are there in the cubes of numbers that have 1 figure? 2 figures? 3 figures? 4 figures?

2. What places, then, belong to the cube of units? of tens? of hundreds? of thousands?

3. How many figures are there in the cube root of a number expressed by 4 figures? 5 figures? 6 figures? 7 figures?

4. How may the number of figures in the cube root of a number be found?

PRINCIPLES. — 1. *The cube of a number is expressed by three times as many figures as the number itself, or by one or two less than three times as many.*

2. *The orders of units in the cube root of a number correspond to the number of periods of three figures each, into which the number can be separated, beginning at units.*

236. If the number of units expressed by the tens' digit is represented by t , and the number of units expressed by the units' digit by u , the cube of a number consisting of tens and units will be represented by $(t + u)^3$, or $t^3 + 3t^2u + 3tu^2 + u^3$.

Thus, $25 = 2 \text{ tens} + 5 \text{ units, or } 20 + 5 \text{ units,}$
and $25^3 = 20^3 + 3(20^2 \times 5) + 3(20 \times 5^2) + 5^3 = 15625.$

EXAMPLES

1. What is the cube root of 12167?

FIRST PROCESS

		12·167	20 + 3
	$t^3 =$	8 000	
Trial divisor,	$3t^2 = 1200$	4 167	
	$3tu = 180$		
	$u^2 = 9$		
Complete divisor,	= 1389	4 167	

EXPLANATION. — By Prin. 2, § 235, the orders of units may be determined by separating the number into periods of three figures each, beginning at units. Separating 12167 thus, there are found to be two orders of units in the root; that is, the root is composed of tens and units.

Since the cube of tens is thousands, and the thousands of the power are less than 27, or 3^3 , and more than 8, or 2^3 , the tens' figure of the root is 2. 2 tens, or 20, cubed is 8000, and 8000 subtracted from 12167 leaves 4167, which is equal to 3 times the $\text{tens}^2 \times \text{the units} + 3 \text{ times the tens} \times \text{the units}^2 + \text{the units}^3$.

Since 3 times the $\text{tens}^2 \times \text{the units}$ is much greater than 3 times the tens \times the units^2 , 4167 is a little more than 3 times the $\text{tens}^2 \times \text{the units}$. If, then, 4167 is divided by 3 times the tens^2 , or by 1200, the trial divisor, the quotient 3 will be the units of the root, provided proper allowance has been made for the additions necessary to obtain the complete divisor.

Since the complete divisor is found by adding to 3 times the tens^2 3 times the tens \times the units and the units^2 , the complete divisor is $1200 + 180 + 9$, or 1389. This multiplied by 3 gives 4167, which subtracted from 4167 leaves no remainder. Therefore, the cube root of 12167 is $20 + 3$, or 23.

SECOND PROCESS

$$\begin{array}{r}
 12\text{-}167 \quad | \quad 23 \\
 t^3 = 8 \\
 3t^2 = 1200 \quad | \quad 4\ 167 \\
 3tu = 180 \\
 u^3 = 9 \\
 \hline
 1389 \quad | \quad 4\ 167
 \end{array}$$

EXPLANATION. — In practice it is usual to place figures of the same order in a column, and to disregard the ciphers on the right of the products.

2. What is the cube root of 1740992427 ?

SOLUTION

$$\begin{array}{r}
 1\text{-}740\text{-}992\text{-}427 \quad | \quad 1203 \\
 \begin{array}{l} \text{Complete} \\ \text{divisor} \end{array} \left\{ \begin{array}{l} t^3 = \\ 3t^2 = 3(10)^2 = 300 \\ 3tu = 3(10 \times 2) = 60 \\ u^2 = 2^2 = 4 \end{array} \right. \\
 \hline
 364 \quad | \quad 740 \\
 \begin{array}{l} \text{Complete} \\ \text{divisor} \end{array} \left\{ \begin{array}{l} 3t^2 = 3(120)^2 = 43200 \\ 3t^2 = 3(1200)^2 = 4320000 \\ 3tu = 3(1200 \times 3) = 10800 \\ u^2 = 3^2 = 9 \end{array} \right. \\
 \hline
 4330809 \quad | \quad 12992427
 \end{array}$$

Since a root expressed by any number of figures may be regarded as composed of tens and units, the processes of example 1 have a general application. Thus, $120 = 12 \text{ tens} + 0 \text{ units}$; and $1203 = 120 \text{ tens} + 3 \text{ units}$.

Since the third figure of the root is 0, it is not necessary to form the complete divisor, inasmuch as the product to be subtracted will be 0.

RULE. — *Separate the numbers into periods of three figures each, beginning at units.*

Find the greatest cube in the left-hand period, and write its root for the first term of the required root. Cube the root, subtract the result from the left-hand period, and annex to the remainder the next period for a new dividend.

Take 3 times the square of the root already found, with two ciphers annexed, for a trial divisor, and by it divide the dividend. The quotient, or quotient diminished, will be the second figure of the root.

To this trial divisor add 3 times the product of the first part of the root with a cipher annexed, multiplied by the second part, and also the square of the second part. Their sum will be the complete divisor.

Multiply the complete divisor by the second part of the root, and subtract the product from the dividend.

Continue thus until all the figures of the root have been found.

1. When there is a remainder after subtracting the last product, annex decimal ciphers, and continue the process.

2. Decimals are pointed off from the decimal point toward the right.

3. The cube root of a common fraction may be found by extracting the cube root of both numerator and denominator separately, or by reducing it to a decimal and then extracting its root.

Extract the cube root of

3. 29791.	9. 2406104.	15. .000024389.
4. 54872.	10. 69426531.	16. .001906624.
5. 110592.	11. 28372625.	17. .000912673.
6. 300763.	12. 48.228544.	18. .259694072.
7. 681472.	13. 17173.512.	19. 926.859375.
8. 941192.	14. 95.443993.	20. 514500.058197.

Extract the cube root to three decimal places :

21. 2.	23. .8.	25. $\frac{5}{64}$.	27. $\frac{7}{8}$.
22. 5.	24. .16.	26. $\frac{2}{3}$.	28. $\frac{2}{16}$.

237. To extract any root of a polynomial.

To find a formula for obtaining the complete divisor in extracting the *fourth*, *fifth*, *sixth*, or any required root of a polynomial, raise $(a + b)$ to the required power, and separate all the terms after the first into two factors one of which shall be the second term of the root. The other factor will be the formula for the complete divisor.

$$(a + b)^5 = a^5 + 5 a^4 b + 10 a^3 b^2 + 10 a^2 b^3 + 5 a b^4 + b^5.$$

Trial divisor, $5 a^4$.

Complete divisor, $5 a^4 + 10 a^3 b + 10 a^2 b^2 + 5 a b^3 + b^4$.

$$(a + b)^7 = a^7 + 7 a^6 b + 21 a^5 b^2 + 35 a^4 b^3 + 35 a^3 b^4 + 21 a^2 b^5 + 7 a b^6 + b^7.$$

Trial divisor, $7 a^6$.

Complete divisor, $7 a^6 + 21 a^5 b + 35 a^4 b^2 + 35 a^3 b^3 + 21 a^2 b^4 + 7 a b^5 + b^6$.

Since the *fourth* power is the *square* of the *second* power, the *sixth* power the *cube* of the *second* power, etc., any root whose index is 4, 6, 8, 9, etc., may be found by extracting successively the roots corresponding to the factors of the index.

The fourth root may be obtained by extracting the square root of the square root; the sixth root, by extracting the cube root of the square root, or the square root of the cube root; the eighth root, by extracting the square root of the square root of the square root.

EXAMPLES

1. Find the fourth root of $16 - 32x + 24x^2 - 8x^3 + x^4$.
2. Find the fourth root of $x^4 + 12x^3y + 54x^2y^2 + 108xy^3 + 81y^4$.
3. Find the fourth root of $16m^4 - 32m^3 + 24m^2 - 8m + 1$.
4. Find the fifth root of $32x^5 + 80x^4 + 80x^3 + 40x^2 + 10x + 1$.
5. Find the fifth root of $a^{10} + 15a^8 + 90a^6 + 270a^4 + 405a^2 + 243$.
6. Find the sixth root of $x^6 - 12x^5 + 60x^4 - 160x^3 + 240x^2 - 192x + 64$.
7. Find the sixth root of $64x^6 - 576x^5 + 2160x^4 - 4320x^3 + 4860x^2 - 2916x + 729$.
8. Find the sixth root of $x^6 + 6acx^5 + 15a^2c^2x^4 + 20a^3c^3x^3 + 15a^4c^4x^2 + 6a^5c^5x + a^6c^6$.

238. To extract any root of an arithmetical number.**EXAMPLES**

1. Find the cube root of 42875.

SOLUTIONBy factoring, $42875 = 5 \times 5 \times 5 \times 7 \times 7 \times 7$. \therefore §§ 26, 229, Prin. 3, $\sqrt[3]{42875} = 5 \times 7 = 35$.

Find, by factoring, the roots indicated :

- | | | |
|-----------------------|-------------------------|--------------------------|
| 2. $\sqrt[3]{3375}$. | 4. $\sqrt[6]{531441}$. | 6. $\sqrt[5]{4084101}$. |
| 3. $\sqrt[4]{1296}$. | 5. $\sqrt[5]{759375}$. | 7. $\sqrt[6]{262144}$. |

8. Find the sixth root of 1771561.

SOLUTION. — The square root of 1771561 is 1331.

The cube root of 1331 is 11.

 \therefore § 237, $\sqrt[6]{1771561} = 11$.

Find the roots indicated :

- | | | |
|-------------------------|---------------------------|-------------------------------|
| 9. $\sqrt[4]{50625}$. | 11. $\sqrt[6]{531441}$. | 13. $\sqrt[6]{24137569}$. |
| 10. $\sqrt[6]{46656}$. | 12. $\sqrt[8]{5764801}$. | 14. $\sqrt[9]{10604499373}$. |

239. Factoring by evolution.**EXAMPLES**

1. Factor
- $x^4 + 4x^3 + 8x^2 + 8x - 5$
- .

SOLUTION. — Extracting the square root as far as possible, the root obtained is $x^2 + 2x + 2$ with a remainder of -9 .Therefore, $x^4 + 4x^3 + 8x^2 + 8x - 5$

$$= (x^2 + 2x + 2)^2 - 9$$

§ 128, $= (x^2 + 2x + 2 + 3)(x^2 + 2x + 2 - 3)$

$$= (x^2 + 2x + 5)(x^2 + 2x - 1).$$

Factor the following polynomials:

- $x^4 + 6x^3 + 11x^2 + 6x - 8$.
- $x^6 + 2x^5 + 5x^4 + 8x^3 + 8x^2 + 8x + 3$.
- $x^6 - 4x^5 + 6x^4 + 6x^3 - 19x^2 + 10x + 9$.
- $4x^6 + 12x^5 + 25x^4 + 40x^3 + 40x^2 + 32x + 15$.

THEORY OF EXPONENTS

240. Thus far the exponents used have been *positive integers* only, and consequently the laws relating to exponents have been obtained in the following restricted forms:

1. $a^m \times a^n = a^{m+n}$ when m and n are positive integers.
2. $a^m \div a^n = a^{m-n}$ when m and n are positive integers and $m > n$.
3. $(a^m)^n = a^{mn}$ when m and n are positive integers.
4. $\sqrt[n]{a^m} = a^{m/n}$ when m and n are positive integers, and m is a multiple of n .
5. $(ab)^n = a^n b^n$ when n is a positive integer.

If all restrictions are removed from m and n , we may then have expressions like a^{-2} and $a^{\frac{1}{3}}$. But such expressions are, as yet, unintelligible, because -2 and $\frac{1}{3}$ cannot show how many times a number is used as a factor.

Since, however, these forms may occur in algebraic processes, it is important to discover meanings for them that will allow their use in accordance with the laws already established, for otherwise great complexity and confusion would arise in the processes involving them.

Assuming that the law of exponents for multiplication,

$$a^m \times a^n = a^{m+n}, \tag{I}$$

is true for *all* values of m and n , the meanings of zero, negative, and fractional exponents may be readily discovered.

Then, having verified the remaining laws of exponents for these exponents, all the laws will have been shown to be of general application for all commensurable exponents.

241. The meaning of a zero exponent.

1. What is the exponent of a in the product of $a^3 \times a^2$? of $a^3 \times a^1$? of $a^3 \times a^0$, if the law of exponents for multiplication is true for *all* values of m and n ?

2. Since $a^3 \times a^0 = a^3$, what is the value of a^0 ? What is the value of x^0 ? of b^0 ? of y^0 ? of m^0 ?

3. What, then, is the value of the zero power of any number?

242. PRINCIPLE. — *Any number with a zero exponent is equal to 1.*

The above principle may be established as follows:

It is to be proved that $a^0 = 1$.

Since Law I is to be true for all values of m and n , § 240, when $n = 0$,

$$a^m \times a^0 = a^{m+0} = a^m.$$

Dividing by a^m ,

$$a^0 = \frac{a^m}{a^m} = 1.$$

243. The meaning of a negative exponent.

1. What is the exponent of a in the product of $a^4 \times a^2$? of $a^4 \times a$? of $a^4 \times a^0$? of $a^4 \times a^{-1}$, if the law of exponents for multiplication is true for all values of m and n ? of $a^4 \times a^{-2}$?

2. Since $a^4 \times a^{-1} = a^3$ and since $a^4 \times \frac{1}{a} = a^3$, to what fraction is a^{-1} equal? Since $a^4 \times a^{-2} = a^2$, to what fraction is a^{-2} equal?

3. What is another expression for a^{-3} ? for x^{-3} ? for y^{-4} ?

4. What is the equivalent of any number with a negative exponent?

5. Since $a^{-2} = \frac{1}{a^2}$, to what is $a^{-2}b$ equal? $\frac{a^{-2}b}{c}$?

6. Since $\frac{1}{a^{-2}} = 1 + \frac{1}{a^2}$, or a^2 , to what is $\frac{1}{a^{-2}} \times \frac{b}{c}$, or $\frac{b}{a^{-2}c}$, equal?

7. Without changing the value of the fraction, transfer a from the numerator to the denominator in $\frac{a^{-3}b}{c}$; in $\frac{a^2b}{c}$; from the denominator to the numerator in $\frac{b}{a^{-2}c}$; in $\frac{b}{a^2c}$.

8. How may a factor be transferred from one term of a fraction to the other without changing the value of the fraction?

244. PRINCIPLES. — 1. *Any number with a negative exponent is equal to the reciprocal of the same number with a numerically equal positive exponent.*

2. *Any factor may be transferred from one term of a fraction to the other, without changing the value of the fraction, if the sign of the exponent is changed.*

The above principles may be established as follows:

PRINCIPLE 1. It is to be proved that $a^{-n} = \frac{1}{a^n}$.

Since § 240, Law I, $a^m \times a^n = a^{m+n}$, is to hold for all values of m and n , when $m = -n$,

$$a^{-n} \times a^n = a^{-n+n} = a^0;$$

but, § 242,

$$a^0 = 1;$$

∴ Ax. 1,

$$a^{-n} \times a^n = 1.$$

Dividing by a^n ,

$$a^{-n} = \frac{1}{a^n}.$$

PRINCIPLE 2. It is to be proved that $\frac{a^{-m}}{b^{-n}} = \frac{b^n}{a^m}$.

By Prin. 1, $a^{-m} = \frac{1}{a^m}$ and $b^{-n} = \frac{1}{b^n}$.

Therefore,

$$\frac{a^{-m}}{b^{-n}} = \frac{\frac{1}{a^m}}{\frac{1}{b^n}} = \frac{1}{a^m} \times \frac{b^n}{1} = \frac{b^n}{a^m}.$$

EXAMPLES

Write the following with negative exponents:

1. $1 + a.$

3. $1 + a^n.$

5. $c + a^2x^2.$

2. $1 + a^2.$

4. $a + x^3.$

6. $am^3 + bx^n.$

7. Write $5x^3y^2$ with positive exponents.

SOLUTION. — Prin. 1, $5x^3y^2 = 5y^2 \frac{1}{x^3} = \frac{5y^2}{x^3}.$

Write the following with positive exponents:

8. $2x^{-1}.$

11. $a^{-1}b^{-1}.$

14. $4a^2c^{-2}.$

9. $5a^{-3}.$

12. $x^{-3}y^{-2}.$

15. $3ax^{-2}.$

10. $3b^{-2}.$

13. $a^{-1}b^2c^{-3}.$

16. $a^nb^{-2n}.$

17. Write $\frac{3a^2}{x^2}$ without a denominator.

SOLUTION. — Prin. 2, $\frac{3a^2}{x^2} = 3a^2x^{-2}$.

Write the following without denominators:

18. $\frac{ax}{by}$

23. $\frac{1}{a^{-2}b^3}$

28. $\left(\frac{x}{y}\right)^2$

19. $\frac{mn}{a^2}$

24. $\frac{1}{m^{-3}n}$

29. $\left(\frac{3}{m}\right)^3$

20. $\frac{c^2a^2}{a^2b^2}$

25. $\frac{3a^2c^{-2}}{x^2}$

30. $\left(\frac{a}{b}\right)^3$

21. $\frac{a^2m^3}{b^3n^2}$

26. $\frac{4}{x^{-4}}$

31. $\frac{a^{-3}}{b^{-3}}$

22. $\frac{b}{x^{-6}}$

27. $\frac{x^{-1}}{y}$

32. $\frac{1}{(ab)^3}$

245. The meaning of a fractional exponent.

1. What is the cube root of a^6 ? How is its exponent obtained? Express the root with this division indicated.

2. In $a^{\frac{6}{3}}$ what does 6 express with reference to a ? What does 3 express with reference to a^6 ?

3. What is the fifth root of b^{15} ? Express the root with a fractional exponent.

4. In $b^{\frac{15}{5}}$ what does 15 express with reference to b ? What does 5 express with reference to b^{15} ?

5. Express the cube root of a with a fractional exponent; the fourth root of a^5 ; the third root of the seventh power of a .

6. What does the numerator of a fractional exponent indicate? What, the denominator?

7. Since $a^{\frac{1}{3}} \times a^{\frac{1}{3}} \times a^{\frac{1}{3}} = a^1$, if the law of exponents for multiplication holds true for all values of m and n , of what is $a^{\frac{1}{3}}$ the cube root?

Since $a^{\frac{1}{6}} \times a^{\frac{1}{6}} \times a^{\frac{1}{6}} \times a^{\frac{1}{6}} \times a^{\frac{1}{6}} \times a^{\frac{1}{6}} = a^1$, of what is $a^{\frac{1}{6}}$ the sixth power? What two meanings may $a^{\frac{1}{3}}$ have? $a^{\frac{2}{3}}$? $a^{\frac{4}{3}}$?

8. What does a fractional exponent indicate?

246. PRINCIPLES.—1. *The numerator of a positive fractional exponent indicates a power and the denominator a root.*

2. *A positive fractional exponent indicates a root of a power or a power of a root.*

Prin. 2 refers to principal roots only (§ 225).

The above principles may be established as follows:

Let p and q be any positive integers, and a any positive number.

Since $\frac{-p}{-q} = \frac{p}{q}$, $\frac{p}{q}$ may represent any positive fraction.

1. It is to be proved that in $a^{\frac{p}{q}}$, p indicates a power and q a root.

Since Law I, or $a^m \times a^n = a^{m+n}$, is to hold for all values of m and n ,

when

$$m = n = \frac{p}{q},$$

$$a^{\frac{p}{q}} \times a^{\frac{p}{q}} = a^{\frac{p}{q} + \frac{p}{q}} = a^{\frac{2p}{q}},$$

also

$$a^{\frac{p}{q}} \times a^{\frac{p}{q}} \times a^{\frac{p}{q}} = a^{\frac{2p}{q}} \times a^{\frac{p}{q}} = a^{\frac{3p}{q}}, \text{ etc.}$$

Hence, $a^{\frac{p}{q}} \times a^{\frac{p}{q}} \times a^{\frac{p}{q}} \dots$ to q factors $= a^{\frac{qp}{q}} = a^p$.

Taking the q th root, § 26 and Ax. 7,

$$a^{\frac{p}{q}} = \sqrt[q]{a^p}.$$

Hence, in $a^{\frac{p}{q}}$, p indicates a power and q a root, and $a^{\frac{p}{q}}$ indicates the q th root of the p th power of a .

2. It is to be proved that $a^{\frac{p}{q}} = \sqrt[q]{a^p}$, or $(\sqrt[q]{a})^p$.

By the previous proof, $a^{\frac{p}{q}} = \sqrt[q]{a^p}$. (1)

If $p = 1$, $a^{\frac{1}{q}} = \sqrt[q]{a}$.

Raising to the p th power,

$$a^{\frac{1}{q}} \times a^{\frac{1}{q}} \times a^{\frac{1}{q}} \dots \text{ to } p \text{ factors} = (\sqrt[q]{a})^p,$$

or, Law I,

$$a^{\frac{p}{q}} = (\sqrt[q]{a})^p. \quad (2)$$

Hence, (1) and (2), $a^{\frac{p}{q}}$ indicates the q th root of the p th power of a , or the p th power of the q th root of a .

It follows from § 244 that $a^{-\frac{p}{q}} = \frac{1}{a^{\frac{p}{q}}}$.

EXAMPLES

1. Express $\sqrt[3]{a^2bc^{-4}}$ with positive fractional exponents.

SOLUTION.
$$\sqrt[3]{a^2bc^{-4}} = a^{\frac{2}{3}}b^{\frac{1}{3}}c^{-\frac{4}{3}} = \frac{a^{\frac{2}{3}}b^{\frac{1}{3}}}{c^{\frac{4}{3}}}.$$

Express with positive fractional exponents:

- | | | |
|----------------------|-------------------------|-----------------------------|
| 2. $\sqrt{ab^3}$. | 5. $(\sqrt{x})^3$. | 8. $(\sqrt[3]{xy})^{-2}$. |
| 3. \sqrt{xy} . | 6. $(\sqrt[5]{y})^4$. | 9. $5\sqrt{x^{-1}y^{-1}}$. |
| 4. $\sqrt{x^2y^5}$. | 7. $(\sqrt[5]{ab})^3$. | 10. $2\sqrt[3]{(a+b)^2}$. |

In the following, large numbers may be avoided by extracting the root first.

Find the value of

- | | | |
|----------------------------|--------------------------|------------------------------|
| 11. $8^{\frac{1}{3}}$. | 15. $64^{\frac{2}{3}}$. | 19. $64^{-\frac{1}{3}}$. |
| 12. $8^{\frac{2}{3}}$. | 16. $32^{\frac{2}{5}}$. | 20. $(-8)^{-\frac{1}{3}}$. |
| 13. $8^{-\frac{1}{3}}$. | 17. $25^{\frac{1}{2}}$. | 21. $(-32)^{-\frac{2}{5}}$. |
| 14. $(-8)^{\frac{1}{3}}$. | 18. $81^{\frac{1}{4}}$. | 22. $16^{-\frac{2}{3}}$. |

Simplify:

23. $\sqrt[3]{x^3} + x^{\frac{1}{3}} + 8^{\frac{2}{3}} + 3x^{\frac{2}{3}} - 5\sqrt[3]{x} - \sqrt[3]{27^2}$.
24. $4\sqrt[5]{x} + 5x^0 - 3x^{-\frac{1}{2}} + 2\sqrt[5]{x^{-1}} - 8^{\frac{2}{3}} - 2x^{\frac{1}{2}}$.
25. $\sqrt[3]{a^2} - \sqrt[3]{a^2b} + \sqrt[3]{ab^2} - b + a + 4\sqrt[3]{a^2b} - 4a^{\frac{1}{3}}b^{\frac{2}{3}} + \sqrt[3]{b^3}$.

Express with radical signs:

- | | | |
|-------------------------|---|--|
| 26. $a^{\frac{2}{3}}$. | 29. $a^{\frac{1}{3}}b^{\frac{2}{3}}$. | 32. $a^{\frac{1}{2}} + x^{\frac{1}{2}}$. |
| 27. $x^{\frac{5}{3}}$. | 30. $x^{\frac{2}{3}}y^{\frac{1}{3}}$. | 33. $m^{\frac{1}{2}} \div n^{\frac{3}{2}}$. |
| 28. $x^{\frac{2}{3}}$. | 31. $a^{\frac{1}{3}}b^{-\frac{2}{3}}$. | 34. $x^{\frac{2}{3}} \div y^{\frac{1}{3}}$. |

247. It now remains to complete the proof that the other laws of exponents are of general application for commensurable exponents by showing that they apply when negative and fractional exponents are employed with the meanings just obtained.

248. To prove that the law of exponents for division is general.

It is to be proved that $a^m \div a^n = a^{m-n}$ for all values of m and n .

Since dividing by a^n is equivalent to multiplying by its reciprocal $\frac{1}{a^n}$,

$$\S 106, \quad a^m \div a^n = a^m \times \frac{1}{a^n}$$

$$\S 244, \text{ Prin. 2,} \quad = a^m \times a^{-n}.$$

$$\text{Hence, for all values of } m \text{ and } n, \quad a^m \div a^n = a^{m-n}. \quad (\text{II})$$

249. To prove that the law of exponents for involution is general.

It is to be proved that $(a^m)^n = a^{mn}$ for all values of m and n .

CASE 1. — Let m represent any value and n a positive integer.

$$\text{Then,} \quad (a^m)^n = a^m \times a^m \times a^m \dots \text{to } n \text{ factors.}$$

$$\begin{aligned} \text{Law I,} \quad &= a^{m+m+m \dots \text{to } n \text{ terms}} \\ &= a^{mn}. \end{aligned}$$

CASE 2. — Let m represent any value, and let $n = \frac{p}{q}$, p and q being positive integers.

$$\text{Then, } \S 246, \text{ Prin. 2,} \quad (a^m)^{\frac{p}{q}} = \sqrt[q]{(a^m)^p}$$

$$\text{Case 1,} \quad = \sqrt[q]{a^{mp}}$$

$$\S 246, \text{ Prin. 1,} \quad = a^{\frac{mp}{q}}.$$

CASE 3. — Let m represent any value, and let $n = -r$, r being a positive integer or a positive fraction.

$$\text{Then,} \quad (a^m)^{-r} = \frac{1}{(a^m)^r}$$

$$\text{Case 1,} \quad = \frac{1}{a^{mr}}$$

$$\S 244, \text{ Prin. 2,} \quad = a^{-mr}.$$

$$\text{Hence, for all values of } m \text{ and } n, \quad (a^m)^n = a^{mn}. \quad (\text{III})$$

250. To prove that the law of exponents for evolution is general.

It is to be proved that $\sqrt[n]{a^m} = a^{m \div n}$ for all values of m and n .

Since $(a^m)^n = a^{mn}$ for all values of m and n , it is true when $\frac{1}{n}$ is substituted for n .

$$\text{Substituting } \frac{1}{n} \text{ for } n, \quad (a^m)^{\frac{1}{n}} = a^{m \times \frac{1}{n}},$$

$$\text{or, } \S 246, \quad \sqrt[n]{a^m} = a^{m \div n}. \quad (\text{IV})$$

251. To prove $(ab)^n = a^n b^n$ for all values of n .

CASE 1. — Let $n = \frac{p}{q}$, p and q being positive integers.

Then, § 249, Case 1, since q is a positive integer,

$$[(ab)^{\frac{p}{q}}]^q = (ab)^{\frac{p}{q} \times q} = (ab)^p$$

$$\S 240, 5, \qquad \qquad \qquad = a^p b^p. \qquad (1)$$

Also, § 249, Case 1,

$$(a^{\frac{p}{q}} b^{\frac{p}{q}})^q = a^{\frac{p}{q} \frac{p}{q} \times q} \times b^{\frac{p}{q} \frac{p}{q} \times q} \dots \text{to } q \text{ factors}$$

$$\S 83, \qquad \qquad \qquad = (a^{\frac{p}{q}} \times a^{\frac{p}{q}} \dots \text{to } q \text{ factors}) (b^{\frac{p}{q}} \times b^{\frac{p}{q}} \dots \text{to } q \text{ factors})$$

$$\qquad \qquad \qquad = a^p b^p. \qquad (2)$$

$$(1) \text{ and } (2), \text{ Ax. 1,} \qquad [(ab)^{\frac{p}{q}}]^q = (a^{\frac{p}{q}} b^{\frac{p}{q}})^q.$$

$$\text{Taking the } q\text{th root,} \qquad (ab)^{\frac{p}{q}} = a^{\frac{p}{q}} b^{\frac{p}{q}}.$$

CASE 2. — Let $n = -r$, r being a positive integer or a positive fraction.

$$\text{Then, § 244, Prin. 1,} \qquad (ab)^{-r} = \frac{1}{(ab)^r}$$

$$\text{Case 1,} \qquad \qquad \qquad = \frac{1}{a^r b^r}$$

$$\S 244, \text{ Prin. 2,} \qquad \qquad \qquad = a^{-r} b^{-r}.$$

$$\text{Hence, for all values of } n, \quad (ab)^n = a^n b^n. \qquad (V)$$

EXAMPLES

252. Multiply :

$$1. \ a^3 \text{ by } a^{-2}. \qquad \qquad 3. \ a^4 \text{ by } a^{-4}. \qquad \qquad 5. \ a^2 \text{ by } a^0.$$

$$2. \ a^2 \text{ by } a^{-1}. \qquad \qquad 4. \ a \text{ by } a^{-3}. \qquad \qquad 6. \ x^{\frac{1}{2}} \text{ by } x^{\frac{1}{3}}.$$

$$7. \ a^{\frac{1}{2}} b^{\frac{1}{3}} \text{ by } a^{\frac{1}{4}} b^{\frac{2}{3}}. \qquad \qquad 10. \ n^{-2} \text{ by } an^{\frac{5}{2}}.$$

$$8. \ m^{\frac{2}{3}} n \text{ by } m^{\frac{1}{3}} n^{-1}. \qquad \qquad 11. \ a^{m-n} \text{ by } a^{n-p}.$$

$$9. \ a^{\frac{1}{2}} b^{\frac{1}{3}} \text{ by } a^{-\frac{1}{4}} b^{\frac{2}{3}}. \qquad \qquad 12. \ a^{\frac{m+n}{2}} \text{ by } a^{\frac{m-n}{2}}.$$

$$13. \text{ Multiply } x^{\frac{1}{2}} y^{-\frac{1}{3}} + x^{\frac{2}{3}} + x^{\frac{2}{3}} y^{\frac{1}{3}} + x^{\frac{1}{2}} y^{\frac{2}{3}} + y^{\frac{2}{3}} \text{ by } x^{\frac{1}{2}} y^{\frac{1}{3}}.$$

$$14. \text{ Multiply } y^n + x^{-1} y^{n+1} + x^{-2} y^{n+2} + x^{-3} y^{n+3} \text{ by } x^n y^{-n}.$$

15. Expand $(a^{\frac{1}{2}}b^{-\frac{1}{2}} + 1 + a^{-\frac{1}{2}}b^{\frac{1}{2}})(a^{\frac{1}{2}}b^{-\frac{1}{2}} - 1 + a^{-\frac{1}{2}}b^{\frac{1}{2}})$

FIRST SOLUTION

$$\begin{array}{r}
 a^{\frac{1}{2}}b^{-\frac{1}{2}} + 1 + a^{-\frac{1}{2}}b^{\frac{1}{2}} \\
 a^{\frac{1}{2}}b^{-\frac{1}{2}} - 1 + a^{-\frac{1}{2}}b^{\frac{1}{2}} \\
 \hline
 a^{\frac{3}{2}}b^{-1} + a^{\frac{1}{2}}b^{-\frac{1}{2}} + a^0b^0 \\
 \quad - a^{\frac{1}{2}}b^{-\frac{1}{2}} - 1 - a^{-\frac{1}{2}}b^{\frac{1}{2}} \\
 \quad \quad + a^0b^0 + a^{-\frac{1}{2}}b^{\frac{1}{2}} + a^{-\frac{3}{2}}b \\
 \hline
 a^{\frac{3}{2}}b^{-1} \quad \quad + 1 \quad \quad + a^{-\frac{3}{2}}b
 \end{array}$$

SECOND SOLUTION

$$\begin{aligned}
 & (a^{\frac{1}{2}}b^{-\frac{1}{2}} + 1 + a^{-\frac{1}{2}}b^{\frac{1}{2}})(a^{\frac{1}{2}}b^{-\frac{1}{2}} - 1 + a^{-\frac{1}{2}}b^{\frac{1}{2}}) \\
 \S 97, \quad & = (a^{\frac{1}{2}}b^{-\frac{1}{2}} + a^{-\frac{1}{2}}b^{\frac{1}{2}})^2 - 1^2 \\
 & = a^{\frac{3}{2}}b^{-1} + 2a^0b^0 + a^{-\frac{3}{2}}b - 1 \\
 & = a^{\frac{3}{2}}b^{-1} + 2 + a^{-\frac{3}{2}}b - 1 \\
 & = a^{\frac{3}{2}}b^{-1} + 1 + a^{-\frac{3}{2}}b.
 \end{aligned}$$

Expand:

- | | |
|--|---|
| 16. $(a^{\frac{1}{2}} + b^{\frac{1}{2}})(a^{\frac{1}{2}} - b^{\frac{1}{2}})$. | 21. $(x^{\frac{2}{3}} - x^{\frac{1}{3}}y^{-\frac{1}{3}} + y^{-\frac{2}{3}})(x^{\frac{1}{3}} + y^{-\frac{1}{3}})$. |
| 17. $(x^{\frac{2}{3}} + y^{\frac{2}{3}})(x^{\frac{1}{3}} - y^{\frac{1}{3}})$. | 22. $(a^{\frac{2}{3}} - b^{-\frac{2}{3}} + a^{\frac{1}{3}}b^{-\frac{1}{3}} + 1)(a^{\frac{1}{3}} - b^{\frac{1}{3}})$. |
| 18. $(x^{-\frac{1}{2}} + 10)(x^{-\frac{1}{2}} - 1)$. | 23. $(1 - x + x^2)(x^{-3} + x^{-2} + x^{-1})$. |
| 19. $(x^{\frac{2}{3}} - 4)(x^{\frac{2}{3}} + 5)$. | 24. $(a^{-1} + b^{-\frac{1}{2}} + c^{\frac{1}{3}})(a^{-1} + b^{-\frac{1}{2}} + 2c^{\frac{1}{3}})$. |
| 20. $(x^{\frac{2}{3}} - y^{\frac{2}{3}})(x^{\frac{1}{3}} + y^{\frac{1}{3}})$. | 25. $(a^2b^2 - ab^3 + b^4)(a^{-2}b^{-2} + a^{-3}b^{-1} + a^{-4})$. |

Divide:

- | | | |
|---|---|--|
| 26. a^5 by a^6 . | 28. a^2 by a^{-2} . | 30. $x^{\frac{1}{2}}$ by $x^{\frac{1}{3}}$. |
| 27. a^3 by a^0 . | 29. $x^{\frac{2}{3}}$ by $x^{-\frac{1}{3}}$. | 31. $x^{n-\frac{1}{2}}$ by x^{n-2} . |
| 32. Divide $x^4 + x^2y^2 + y^4$ by x^2y^2 . | | |
| 33. Divide $a^{-4} + a^{-2}b + b^2$ by $a^{-2}b$. | | |
| 34. Divide $x^4 + 2ax^3 + 3a^2x^2 + a^3x - a^4$ by a^4x^4 . | | |

35. Divide $b^{-1} + 3a^{-\frac{1}{2}} - 10a^{-1}b$ by $a^{\frac{1}{2}}b^{-1} - 2$.

SOLUTION

$$\begin{array}{r}
 a^{\frac{1}{2}}b^{-1} - 2 \big) b^{-1} + 3a^{-\frac{1}{2}} - 10a^{-1}b \big(a^{-\frac{1}{2}} + 5a^{-1}b \\
 \underline{b^{-1} - 2a^{-\frac{1}{2}}} \\
 5a^{-\frac{1}{2}} - 10a^{-1}b \\
 \underline{5a^{-\frac{1}{2}} - 10a^{-1}b}
 \end{array}$$

Divide:

36. $a - b$ by $a^{\frac{1}{2}} + b^{\frac{1}{2}}$. 41. $a^{\frac{2}{3}} + 2a^{\frac{1}{3}}b^{-\frac{1}{3}} + b^{-1}$ by $a^{\frac{1}{3}} + b^{-\frac{1}{3}}$.
 37. $a - b$ by $a^{\frac{1}{3}} - b^{\frac{1}{3}}$. 42. $x^{\frac{4}{3}} - 2 + x^{-\frac{2}{3}}$ by $x^{\frac{2}{3}} - x^{-\frac{2}{3}}$.
 38. $a + b$ by $a^{\frac{1}{2}} + b^{\frac{1}{2}}$. 43. $3 - 4x^{-1} + x^{-2}$ by $x^{-1} - 3$.
 39. $a^2 + b^2$ by $a^{\frac{2}{3}} + b^{\frac{2}{3}}$. 44. $4x^2y^{-1} - 5y + x^{-2}y^3$ by $x^2 - y^2$.
 40. $x - 1$ by $x^{\frac{2}{3}} + x^{\frac{1}{3}} + 1$. 45. $a^2 - b^3$ by $a^{\frac{1}{3}} + b^{\frac{1}{3}}$.

Simplify the following:

46. $(a^{\frac{1}{2}})^2$. 49. $(-a^{\frac{1}{3}})^3$. 52. $(8^{-\frac{1}{3}})^4$.
 47. $(a^{-\frac{1}{3}})^6$. 50. $(-a^2)^4$. 53. $(16^{-\frac{1}{2}})^3$.
 48. $(a^{-4})^2$. 51. $(-a^{\frac{2}{3}})^{-1}$. 54. $(-\frac{1}{8}^{\frac{1}{2}})^{-\frac{3}{2}}$.

Expand by the binomial formula:

55. $(a^{\frac{1}{2}} - b^{\frac{1}{2}})^2$. 57. $(a^{-1} - b^{\frac{2}{3}})^3$. 59. $(a^{-\frac{1}{2}} + \frac{1}{2})^3$.
 56. $(a^{\frac{1}{2}} + b^{\frac{1}{2}})^3$. 58. $(x^{\frac{1}{2}} - y^{\frac{1}{2}})^4$. 60. $(1 - x^{\frac{2}{3}})^4$.

Simplify the following:

61. $\sqrt[4]{x^{\frac{1}{2}}}$. 64. $\sqrt[3]{a^{-\frac{1}{2}}b^{-3}}$. 67. $(\frac{1}{8}a^{-\frac{2}{3}}b^{\frac{1}{3}})^{-\frac{3}{2}}$.
 62. $\sqrt[4]{a^{-\frac{1}{2}}}$. 65. $\sqrt{x^{\frac{2}{3}}y^{-3}}$. 68. $(\frac{1}{8}m^{-1}n^{-\frac{1}{2}})^{\frac{1}{2}}$.
 63. $\sqrt[4]{a^{-\frac{1}{2}}}$. 66. $\sqrt[3]{a^2b^9}$. 69. $(4x^{2n}y^{-3z^2})^{\frac{5}{2}}$.

Extract the square root of

70. $x^2 + 2x^{\frac{3}{2}} + 3x + 4x^{\frac{1}{2}} + 3 + 2x^{-\frac{1}{2}} + x^{-1}$.
 71. $x^2 + y + 4x^{-2} - 2xy^{\frac{1}{2}} + 4xz^{-1} - 4y^{\frac{1}{2}}z^{-1}$.
 72. $a + 4b^{\frac{2}{3}} + 9c^{\frac{1}{3}} - 4a^{\frac{1}{2}}b^{\frac{1}{3}} + 6a^{\frac{1}{3}}c^{\frac{1}{2}} - 12b^{\frac{1}{3}}c^{\frac{1}{2}}$.

Extract the cube root of

73. $a^3 + 6a^{\frac{1}{2}} + 12a^{\frac{1}{3}} + 8$.

74. $a - 3a^{\frac{1}{2}}b^{\frac{1}{3}} + 3a^{\frac{1}{3}}b^{\frac{1}{2}} - b^2$.

75. $8x^{-1} - 12x^{-\frac{1}{2}}y + 6x^{-\frac{1}{3}}y^2 - y^3$.

76. $x^{\frac{1}{2}} - 6x + 15x^{\frac{1}{3}} - 20 + 15x^{-\frac{1}{2}} - 6x^{-1} + x^{-\frac{1}{3}}$.

77. Factor $4x^{-2} - 9y^{-2}$, and express the result with positive exponents.

SOLUTION

$$\begin{aligned} \S 128, \quad 4x^{-2} - 9y^{-2} &= (2x^{-1} + 3y^{-1})(2x^{-1} - 3y^{-1}) \\ &= \left(\frac{2}{x} + \frac{3}{y}\right)\left(\frac{2}{x} - \frac{3}{y}\right). \end{aligned}$$

Factor the following and express the results with positive exponents:

78. $a^{-3} - b^{-2}$.

83. $x^3 - x^{-3}$.

79. $9 - x^{-2}$.

84. $a^2 + 2 + a^{-2}$.

80. $16 - a^{-4}$.

85. $b^4 - 8 + 16b^{-4}$.

81. $27 - b^{-3}$.

86. $12 - x^{-1} - x^{-2}$.

82. $b^{-3} + y^{-3}$.

87. $2 - 3x^{-1} - 2x^{-2}$.

Solve the following equations:

88. $x^{\frac{1}{2}} = 2$.

96. $x^{-\frac{1}{2}} = 6$.

89. $x^{\frac{1}{3}} = 8$.

97. $x^{-\frac{1}{3}} = 16$.

90. $x^{\frac{2}{3}} = 4$.

98. $25x^{-\frac{2}{3}} = 1$.

91. $x^{\frac{1}{4}} = 16$.

99. $x^{\frac{1}{2}} = 243$.

92. $\frac{1}{3}x^{\frac{1}{2}} = 9$.

100. $x^{\frac{1}{3}} + 32 = 0$.

93. $x^{-\frac{1}{2}} = 5$.

101. $x^{\frac{1}{2}} + a^6 = 0$.

94. $\frac{1}{4}x^{\frac{1}{3}} = 25$.

102. $x^{\frac{1}{3}} - 64 = 0$.

95. $2x^{-\frac{1}{2}} = \frac{1}{\frac{1}{8}x}$.

103. $x^{-\frac{1}{2}} + 27 = 0$.

RADICALS

253. 1. What is indicated by \sqrt{x} ? by $x^{\frac{1}{2}}$? by $\sqrt[3]{a}$? by $a^{\frac{1}{3}}$?

2. Indicate in two ways the square root of 25; of 36; of 2; of 3.

3. Which of these indicated roots can be obtained exactly? Which cannot be obtained exactly?

254. An indicated root of an expression is called a **Radical**.

The root may be indicated by a *radical sign* or by a *fractional exponent*.

$\sqrt{5a}$, $(5a)^{\frac{1}{2}}$, $\sqrt[3]{ax^2}$, $(ax^2)^{\frac{1}{3}}$, $\sqrt{a^2 + 2ab + b^2}$, and $(a^2 + 2ab + b^2)^{\frac{1}{2}}$ are radicals.

In the discussion and treatment of radicals only principal roots will be considered.

255. The **Degree** of a radical is indicated by the index of the root or by the denominator of the fractional exponent.

$\sqrt{a+x}$ and $(b+x)^{\frac{1}{2}}$ are radicals of the second degree.

256. When the indicated root of a *rational* number cannot be exactly obtained, the expression is called a **Surd**.

$\sqrt{2}$ is a surd, since 2 is rational but has no rational square root.

$\sqrt{1+\sqrt{3}}$ is not a surd, because $1+\sqrt{3}$ is not rational.

Radicals may be either rational or irrational, but surds are always irrational.

Both $\sqrt{4}$ and $\sqrt{3}$ are radicals but only $\sqrt{3}$ is a surd.

257. An indicated even root of a negative number is called an **Imaginary Number**; as $\sqrt{-4}$, $\sqrt{-a}$.

All other numbers, whether rational or irrational, are called *Real Numbers*; as $\sqrt{25}$, $\sqrt{3}$, $a^{\frac{1}{2}}$, 4.

258. A surd may contain a *rational factor*, that is, a factor which is a perfect power of the same degree as the radical. The rational factor may be removed and written as the coefficient of the irrational factor.

In $\sqrt{8} = \sqrt{4 \times 2}$ and $\sqrt[3]{54} = \sqrt[3]{27 \times 2}$, the rational factors are $\sqrt{4}$ and $\sqrt[3]{27}$; that is, $\sqrt{8} = 2\sqrt{2}$ and $\sqrt[3]{54} = 3\sqrt[3]{2}$.

259. A surd that has a rational coefficient is called a **Mixed Surd**.

$2\sqrt{2}$, $a\sqrt[3]{x^2}$, and $(a-b)\sqrt{a+b}$ are mixed surds.

260. A surd that has no rational coefficient except unity is called an **Entire Surd**.

$\sqrt{5}$, $\sqrt[3]{11}$, and $\sqrt{a^2 + x^2}$ are entire surds.

261. A radical is in its **simplest form** when the expression under the radical sign is integral, contains no factor that is a power of the same degree as the radical, and is not itself a perfect power whose exponent is a factor of the index of the radical.

$\sqrt{7}$ is in its simplest form; but $\sqrt{\frac{7}{4}}$ is not in its simplest form, because $\frac{7}{4}$ is not integral in form; $\sqrt{8}$ is not in its simplest form, because the square root of 4, a factor of 8, may be extracted; $\sqrt[3]{25}$, or $25^{\frac{1}{3}}$, is not in its simplest form, because $25^{\frac{1}{3}} = (5^2)^{\frac{1}{3}} = 5^{\frac{2}{3}} = 5^{\frac{1}{3}} \cdot 5^{\frac{1}{3}}$, or $\sqrt[3]{5}$.

REDUCTION OF RADICALS

262. To reduce a radical to its simplest form when it has a rational factor.

EXAMPLES

1. Reduce $\sqrt{20a^6}$ to its simplest form.

PROCESS

$$\sqrt{20a^6} = \sqrt{4a^6 \times 5} = \sqrt{4a^6} \times \sqrt{5} = 2a^3\sqrt{5}$$

EXPLANATION. — Since the highest factor of $20a^6$ that is a perfect square is $4a^6$, $\sqrt{20a^6}$ is separated into two factors, a rational factor $\sqrt{4a^6}$, and an irrational factor $\sqrt{5}$. $\sqrt{20a^6} = \sqrt{4a^6} \times \sqrt{5}$, § 229, Prin. 3. Extracting the square root of $4a^6$ and prefixing the root to the irrational factor as a coefficient, the result is $2a^3\sqrt{5}$.

2. Reduce $\sqrt[3]{-864}$ to its simplest form.

PROCESS

$$\sqrt[3]{-864} = \sqrt[3]{-216 \times 4} = \sqrt[3]{-216} \times \sqrt[3]{4} = -6\sqrt[3]{4}$$

RULE.—Separate the radical into two factors one of which is its highest rational factor. Extract the required root of the rational factor, multiply the result by the coefficient, if any, of the given radical, and place the product as the coefficient of the irrational factor.

Simplify the following:

- | | | |
|--|---|--|
| 3. $\sqrt{12}$. | 15. $\sqrt{162}$. | 27. $\sqrt{243 a^2 x^{10}}$. |
| 4. $\sqrt{75}$. | 16. $\sqrt{18 a^2}$. | 28. $\sqrt[3]{128 a^6 b^4}$. |
| 5. $\sqrt[3]{16}$. | 17. $\sqrt{25 b}$. | 29. $\sqrt{405 a^2 y^2}$. |
| 6. $\sqrt{128}$. | 18. $\sqrt{98 c^2}$. | 30. $\sqrt{375 x^2 y^2}$. |
| 7. $\sqrt[3]{250}$. | 19. $\sqrt{50 a}$. | 31. $(245 a^4 y^{-4})^{\frac{1}{2}}$. |
| 8. $\sqrt[4]{32}$. | 20. $\sqrt[5]{640}$. | 32. $(135 x^4 y^6)^{\frac{1}{3}}$. |
| 9. $\sqrt{600}$. | 21. $\sqrt{84}$. | 33. $(a^3 + 5 a^2)^{\frac{1}{2}}$. |
| 10. $\sqrt{500}$. | 22. $\sqrt[3]{72}$. | 34. $(16x - 16)^{\frac{1}{2}}$. |
| 11. $\sqrt[5]{160}$. | 23. $\sqrt[3]{192}$. | 35. $\sqrt{18x - 9}$. |
| 12. $\sqrt[3]{3000}$. | 24. $\sqrt{800}$. | 36. $\sqrt[3]{x^6 - 2x^3}$. |
| 13. $\sqrt[3]{81}$. | 25. $\sqrt[3]{3645}$. | 37. $\sqrt{8 - 20b^2}$. |
| 14. $\sqrt[3]{189}$. | 26. $\sqrt{735}$. | 38. $5\sqrt{4a^2 + 4}$. |
| 39. $\sqrt{5x^2 - 10xy + 5y^2}$. | 41. $(3am^2 + 6am + 3a)^{\frac{1}{2}}$. | |
| 40. $\sqrt{4a^3 - 24a^2x + 36ax^2}$. | 42. $(x^4y - 3x^2y^2 + 3x^2y^3 - xy^4)^{\frac{1}{2}}$. | |
| 43. Reduce $\sqrt{\frac{a^2}{2y^3}}$ to its simplest form. | | |

PROCESS

$$\sqrt{\frac{a^2}{2y^3}} = \sqrt{\frac{a^2 \times 2y}{2y^3 \times 2y}} = \sqrt{\frac{a^2}{4y^4}} \times \sqrt{2y} = \frac{a}{2y^2} \sqrt{2y}$$

EXPLANATION. — Since a radical is not in its simplest form when the expression under the radical sign is fractional, the denominator is to be removed; and since the radical is of the second degree, the denominator must be made a perfect square. The smallest factor that will accomplish this is $2y$. Multiplying the terms of the fraction by this factor, the largest rational factor of the resulting radical is found to be $\sqrt{\frac{a^2}{4y^4}}$, which is equal to $\frac{a}{2y^2}$. Therefore, the irrational factor is $\sqrt{2y}$, and its coefficient is $\frac{a}{2y^2}$.

Simplify the following:

44. $\sqrt{\frac{1}{2}}$.

45. $\sqrt{\frac{1}{8}}$.

46. $\sqrt{\frac{1}{5}}$.

47. $\sqrt{\frac{1}{8}}$.

48. $\sqrt{\frac{2}{3}}$.

49. $\sqrt{\frac{3}{8}}$.

50. $\sqrt[3]{\frac{5}{12}}$.

51. $\sqrt[3]{\frac{4}{3}}$.

52. $\sqrt{\frac{2a^3}{b}}$.

53. $\sqrt{\frac{5x^4y^2}{2a^2}}$.

54. $\sqrt[4]{\frac{x}{y}}$.

55. $\sqrt[4]{\frac{1}{x^3}}$.

56. $\sqrt{\frac{2}{3y^3}}$.

57. $\sqrt{\frac{4a}{3x^2}}$.

58. $\sqrt{\frac{3x}{50a^3y}}$.

59. $\sqrt[3]{\frac{a}{3b^2}}$.

60. $(a+b)\sqrt{\frac{a+b}{a-b}}$.

62. $(1-x^2)\sqrt{\frac{1-x+x^2}{1+x+x^2}}$.

61. $\frac{2y}{x-2y}\sqrt{\frac{x-2y}{2y}}$.

63. $\frac{(a+b)^2}{a-b}\sqrt[3]{\frac{a+b}{(a-b)^2}}$.

263. To reduce a radical to its simplest form when the expression under the radical sign is a perfect power of a degree corresponding to some factor of the index of the root.

EXAMPLES

1. Reduce $\sqrt[6]{9a^2}$ to its simplest form.

PROCESS

$$\sqrt[6]{9a^2} = \sqrt[6]{(3a)^2} = (3a)^{\frac{2}{3}} = (3a)^{\frac{1}{3}} = \sqrt[3]{3a}$$

2. Reduce $\sqrt[9]{64a^6b^{15}}$ to its simplest form.

PROCESS

$$\sqrt[9]{64a^6b^{15}} = \sqrt[9]{2^6a^6b^6b^9} = b(2ab)^{\frac{2}{3}} = b(2ab)^{\frac{2}{3}} = b\sqrt[3]{4a^2b^2}$$

Simplify the following:

- | | | |
|----------------------|------------------------|---------------------------------|
| 3. $\sqrt[4]{36}$. | 7. $\sqrt[4]{1600}$. | 11. $\sqrt[4]{9a^2b^2c^3}$. |
| 4. $\sqrt[4]{25}$. | 8. $\sqrt[6]{27a^3}$. | 12. $\sqrt[4]{121a^6x^4}$. |
| 5. $\sqrt[4]{144}$. | 9. $\sqrt[6]{343}$. | 13. $\sqrt[6]{a^4b^2c^4d^8}$. |
| 6. $\sqrt[6]{81}$. | 10. $\sqrt[4]{289}$. | 14. $\sqrt[4]{(x^2-2xy+y^2)}$. |

264. To reduce a mixed surd to an entire surd.

EXAMPLES

1. Express $2a\sqrt{5b}$ as an entire surd.

PROCESS

$$2a\sqrt{5b} = \sqrt{4a^2}\sqrt{5b} = \sqrt{4a^2 \times 5b} = \sqrt{20a^2b}$$

RULE.—*Raise the coefficient to a power corresponding to the index of the given radical, and introduce the result under the radical sign as a factor.*

Express as entire surds:

- | | | | |
|---|---|--|---|
| 2. $2\sqrt{2}$. | 6. $3\sqrt[3]{3}$. | 10. $\frac{1}{2}\sqrt{2}$. | 14. $\frac{4}{5}\sqrt[4]{\frac{3}{8}}$. |
| 3. $3\sqrt{5}$. | 7. $4\sqrt{5}$. | 11. $\frac{3}{4}\sqrt{x^2}$. | 15. $\frac{3}{2}\sqrt{\frac{3}{7}a^2}$. |
| 4. $5\sqrt{2}$. | 8. $\frac{1}{2}\sqrt{8}$. | 12. $\frac{1}{2}\sqrt{bc}$. | 16. $\frac{2}{3}\sqrt[3]{1\frac{1}{8}}$. |
| 5. $3\sqrt[4]{2}$. | 9. $a^2\sqrt[3]{b}$. | 13. $\frac{3}{4}\sqrt[4]{\frac{7}{9}}$. | 17. $\frac{2}{3}\sqrt[4]{\frac{3}{8}}$. |
| 18. $\frac{x+y}{x-y}\sqrt{\frac{x-y}{x+y}}$. | 19. $\frac{a+4}{a-4}\sqrt{1-\frac{8}{a+4}}$. | 20. $\frac{1}{ab}(a-b)^{\frac{1}{2}}$. | |

265. To reduce radicals of different degrees to equivalent radicals of the same degree.

1. Express $a^{\frac{1}{2}}$ by an equivalent radical with an exponent in higher terms.

2. What is the degree of the radical $x^{\frac{1}{2}}$? Express $x^{\frac{1}{2}}$ as a radical of the 12th degree. Express $x^{\frac{1}{3}}$ as a radical of the 12th degree. Express $b^{\frac{1}{2}}$ and $b^{\frac{1}{3}}$ as radicals of the same degree.

EXAMPLES

1. Reduce $\sqrt[4]{3}$, $\sqrt{2}$, and $\sqrt[3]{4}$ to equivalent radicals of the same degree.

PROCESS

$$\sqrt[4]{3} = 3^{\frac{1}{4}} = 3^{\frac{3}{12}} = \sqrt[12]{3^3} = \sqrt[12]{27}$$

$$\sqrt{2} = 2^{\frac{1}{2}} = 2^{\frac{6}{12}} = \sqrt[12]{2^6} = \sqrt[12]{64}$$

$$\sqrt[3]{4} = 4^{\frac{1}{3}} = 4^{\frac{4}{12}} = \sqrt[12]{4^4} = \sqrt[12]{256}$$

RULE. — Express the given radicals with fractional exponents having a common denominator.

Raise each number to the power indicated by the numerator of its fractional exponent, and indicate the root expressed by the common denominator.

Reduce to equivalent radicals of the same degree:

2. $\sqrt{2}$ and $\sqrt[4]{3}$.

9. \sqrt{ab} , $\sqrt[3]{ab^2}$, and $\sqrt[4]{2}$.

3. $\sqrt{5}$ and $\sqrt[3]{6}$.

10. \sqrt{a} , $\sqrt[3]{b}$, $\sqrt[4]{x}$, and $\sqrt[6]{y}$.

4. $\sqrt[4]{7}$ and $\sqrt{10}$.

11. $\sqrt[3]{a+b}$ and $\sqrt{x+y}$.

5. $\sqrt[6]{10}$, $\sqrt{2}$, and $\sqrt[3]{5}$.

12. $\sqrt{\frac{2}{3}}$, $\sqrt[3]{\frac{1}{15}x}$, and $2\sqrt{5}$.

6. $\sqrt[6]{4}$, $\sqrt[4]{2}$, and $\sqrt{3}$.

13. $\sqrt[3]{x}$, \sqrt{xy} , and $\sqrt[3]{x^2y^2}$.

7. $\sqrt[10]{13}$, $\sqrt{5}$, and $\sqrt[5]{4}$.

14. $(a+b)\sqrt{a-b}$ and $\sqrt[3]{a-b}$.

8. $\sqrt{3}$, $\sqrt[3]{5}$, and $\sqrt[4]{\frac{1}{27}}$.

15. $\sqrt{a+b}$, $\sqrt[4]{a^2+b^2}$, and $\sqrt{a-b}$.

ADDITION AND SUBTRACTION OF RADICALS

266. Radical terms that, in their simplest forms, are of the same degree and have the same number under the radical sign are called **Similar Radicals**.

267. PRINCIPLE. — Only similar radicals can be united into one term by addition or subtraction.

EXAMPLES

1. Find the sum of $\sqrt{50}$, $2\sqrt[3]{8}$, and $6\sqrt{\frac{1}{2}}$.

PROCESS

$$\sqrt{50} = 5\sqrt{2}$$

$$2\sqrt[3]{8} = 2\sqrt{2}$$

$$6\sqrt{\frac{1}{2}} = 3\sqrt{2}$$

$$\text{Sum} = 10\sqrt{2}$$

EXPLANATION. — To ascertain whether or not the given expressions are similar radicals, each may be reduced to its simplest form. Since, in their simplest form, they are of the same degree and have the same number under the radical sign, they are similar, and their sum is the sum of the coefficients prefixed to the common radical factor.

Find the sum of

2. $\sqrt{50}$, $\sqrt{18}$, and $\sqrt{98}$.
3. $\sqrt{27}$, $\sqrt{12}$, and $\sqrt{75}$.
4. $\sqrt{20}$, $\sqrt{80}$, and $\sqrt{45}$.
5. $\sqrt{28}$, $\sqrt{63}$, and $\sqrt{700}$.
6. $\sqrt[3]{250}$, $\sqrt[3]{16}$, and $\sqrt[3]{54}$.
7. $\sqrt[3]{128}$, $\sqrt[3]{686}$, and $\sqrt[3]{\frac{1}{4}}$.
8. $\sqrt[3]{135}$, $\sqrt[3]{320}$, and $\sqrt[3]{625}$.
9. $\sqrt[3]{500}$, $\sqrt[3]{108}$, and $\sqrt[3]{-32}$.
10. $\sqrt{\frac{1}{2}}$, $\sqrt{12\frac{1}{2}}$, $\sqrt{\frac{1}{8}}$, and $\sqrt{1\frac{1}{8}}$.
11. $\sqrt{\frac{1}{8}}$, $\sqrt{75}$, $\frac{2}{3}\sqrt{3}$, and $\sqrt{12}$.
12. $\sqrt{\frac{2}{3}}$, $\frac{1}{3}\sqrt{3}$, $\frac{1}{4}\sqrt[4]{9}$, and $\sqrt{147}$.
13. $\sqrt[3]{40}$, $\sqrt{28}$, $\sqrt[6]{25}$, and $\sqrt{175}$.
14. $\sqrt[3]{375}$, $\sqrt{44}$, $\sqrt[3]{192}$, and $\sqrt{99}$.

Simplify :

15. $\sqrt{245} - \sqrt{405} + \sqrt{45}$.
16. $\sqrt{12} + 3\sqrt{75} - 2\sqrt{27}$.
17. $5\sqrt{72} + 3\sqrt{18} - \sqrt{50}$.
18. $\sqrt[3]{128} + \sqrt[3]{686} - \sqrt[3]{54}$.
19. $\sqrt{112} - \sqrt{343} + \sqrt{448}$.
20. $\sqrt[3]{135} - \sqrt[3]{625} + \sqrt[3]{320}$.
21. $\sqrt[3]{\frac{8}{9}} + \sqrt[3]{\frac{1}{5}} + \sqrt[3]{5\frac{2}{3}}$.
22. $\sqrt[3]{864} - \sqrt[3]{4000} + \sqrt[3]{32}$.
23. $\sqrt[3]{128x} + \sqrt[3]{375x} - \sqrt[3]{54x}$.
24. $\sqrt{\frac{a}{x^2}} + \sqrt{\frac{a}{y^2}} - \sqrt{\frac{a}{z^2}}$.
25. $\sqrt{\frac{ax^4}{by^2}} - \sqrt{\frac{16ax^2}{by^2}} + \sqrt{\frac{4ax^2}{by^2}}$.
26. $\sqrt{\frac{a}{bc}} + \sqrt{\frac{b}{ac}} + \sqrt{\frac{c}{ab}}$.
27. $\sqrt{(a+b)^2c} - \sqrt{(a-b)^2c}$.
28. $6\sqrt[3]{\frac{49}{27}} + 4\sqrt[3]{1\frac{0}{8}} - 8\sqrt[3]{\frac{875}{8}}$.
29. $\sqrt[5]{-96x^4} + 2\sqrt[5]{3x^4} - \sqrt[5]{5x} + \sqrt[5]{40x^4}$.
30. $\sqrt[3]{abx} - \sqrt[6]{a^2b^2x^2} + \sqrt[9]{8a^3b^3x^3}$.
31. $\sqrt{3x^3 + 30x^2 + 75x} - \sqrt{3x^3 - 6x^2 + 3x}$.
32. $\sqrt{5a^5 + 30a^4 + 45a^3} - \sqrt{5a^5 - 40a^4 + 80a^3}$.
33. $\sqrt{50} + \sqrt[6]{9} - 4\sqrt{\frac{1}{3}} + \sqrt[3]{-24} + \sqrt[9]{27} - \sqrt[4]{64}$.
34. $\sqrt{\frac{2}{3}} + 6\sqrt{\frac{1}{4}} - \frac{1}{5}\sqrt{18} + \sqrt[4]{36} - \sqrt[3]{\frac{1}{8}} + \sqrt[6]{125} - 2\sqrt{\frac{2}{3}}$.

MULTIPLICATION OF RADICALS

268. 1. What is the exponent of a in $a^2 \times a^3$? in $a^{\frac{1}{2}} \times a^{\frac{1}{3}}$? in $a^{\frac{2}{3}} \times a^{\frac{1}{2}}$? in $a^{\frac{1}{2}} \times a^{\frac{1}{3}}$? in $a^{\frac{1}{3}} \times a^{\frac{1}{2}}$? in $\sqrt{a} \times \sqrt[3]{a}$?

2. When the fractional exponents indicate different roots, what must be done before the radicals can be multiplied together?

EXAMPLES

PROCESSES.—1. $\sqrt{7} \times \sqrt{5} = \sqrt{35}$

$$2. 5\sqrt{3} \times 2\sqrt{15} = 10\sqrt{45} = 10 \times 3\sqrt{5} = 30\sqrt{5}$$

$$3. 2\sqrt{3} \times 3\sqrt[3]{2} = 2\sqrt[6]{27} \times 3\sqrt[6]{4} = 6\sqrt[6]{108}$$

RULE.—*If necessary, reduce the radicals to the same degree.*

Multiply the coefficients together for the coefficient of the product and the factors under the radical sign for the radical factor of the product, and simplify the result.

Multiply :

- | | |
|------------------------------------|--|
| 4. $\sqrt{2}$ by $\sqrt{8}$. | 13. $2\sqrt{6}$ by $\sqrt{18}$. |
| 5. $\sqrt{2}$ by $\sqrt{6}$. | 14. $2\sqrt[3]{3}$ by $3\sqrt[3]{45}$. |
| 6. $\sqrt{3}$ by $\sqrt{15}$. | 15. $2\sqrt[4]{6}$ by $3\sqrt{6}$. |
| 7. $\sqrt{3}$ by $\sqrt{48}$. | 16. $3\sqrt{3}$ by $2\sqrt[3]{5}$. |
| 8. $2\sqrt{5}$ by $3\sqrt{10}$. | 17. $\sqrt[4]{5}$ by $\sqrt[5]{10}$. |
| 9. $3\sqrt{20}$ by $2\sqrt{2}$. | 18. $2\sqrt[3]{250}$ by $\sqrt{2}$. |
| 10. $\sqrt{2}$ by $3\sqrt[3]{3}$. | 19. $2\sqrt[3]{24}$ by $\sqrt[3]{18}$. |
| 11. $\sqrt[3]{2}$ by $2\sqrt{5}$. | 20. $\sqrt{28}$ by $3\sqrt{7}$. |
| 12. $\sqrt[3]{3}$ by $3\sqrt{3}$. | 21. $2\sqrt[5]{2}$ by $\sqrt[10]{512}$. |

Find the value of

$$22. \sqrt{ab} \times \sqrt{bc} \times \sqrt{cd} \times \sqrt{da}.$$

$$23. \sqrt{x^2y^2} \times \sqrt{12x} \times \sqrt{75xy^2}.$$

$$24. \sqrt{2ab} \times \sqrt[3]{abc} \times \sqrt[4]{4a^2b^2}.$$

25. $\sqrt{mn} \times \sqrt[4]{m^2n} \times \sqrt[8]{mn^4}.$

26. $\sqrt{2axy} \times \sqrt[3]{xy} \times \sqrt[4]{a^2xy}.$

27. $\sqrt{x^{-1}y} \times \sqrt[3]{x^{-2}y^2} \times \sqrt{x^{-3}y^3}.$

28. $\sqrt{a-b} \times \sqrt[4]{a^2b^2} \times \sqrt[4]{(a-b)^{-2}}.$

29. $\sqrt{\frac{2}{3}} \times \sqrt{\frac{4}{5}} \times \sqrt{\frac{3}{4}}.$

32. $\sqrt[3]{\frac{1}{8}} \times \sqrt[6]{\frac{3}{2}} \times \sqrt{\frac{3}{2}}.$

30. $\sqrt{\frac{1}{2}} \times \sqrt{\frac{4}{5}} \times \sqrt{\frac{7}{2}}.$

33. $\sqrt[6]{\frac{3}{2}} \times \sqrt[3]{\frac{4}{3}} \times \sqrt{\frac{3}{2}}.$

31. $\sqrt[3]{\frac{2}{3}} \times \sqrt[3]{\frac{3}{4}} \times \sqrt{\frac{1}{2}}.$

34. $\sqrt[4]{\frac{2}{3}} \times \sqrt{\frac{3}{4}} \times \sqrt[4]{\frac{8}{3}}.$

35. Multiply $2\sqrt{2} + 3\sqrt{3}$ by $5\sqrt{2} - 2\sqrt{3}.$

SOLUTION

$$2\sqrt{2} + 3\sqrt{3}$$

$$5\sqrt{2} - 2\sqrt{3}$$

$$\hline 20 + 16\sqrt{6}$$

$$- 4\sqrt{6} - 18$$

$$\hline 20 + 11\sqrt{6} - 18$$

$$= 2 + 11\sqrt{6}.$$

Multiply:

36. $\sqrt{5} + \sqrt{3}$ by $\sqrt{5} - \sqrt{3}.$

37. $\sqrt{7} + \sqrt{2}$ by $\sqrt{7} - \sqrt{2}.$

38. $\sqrt{6} - \sqrt{5}$ by $\sqrt{6} - \sqrt{5}.$

39. $5 - \sqrt{5}$ by $1 + \sqrt{5}.$

40. $4\sqrt{7} + 1$ by $4\sqrt{7} - 1.$

41. $2\sqrt{2} + \sqrt{3}$ by $4\sqrt{2} + \sqrt{3}.$

42. $2\sqrt{3} + 3\sqrt{5}$ by $3\sqrt{3} + 2\sqrt{5}.$

43. $3a + \sqrt{5}$ by $2a - \sqrt{5}.$

44. $2\sqrt{6} - 3\sqrt{5}$ by $4\sqrt{3} - \sqrt{10}.$

45. $a^2 - ab\sqrt{2} + b^2$ by $a^2 + ab\sqrt{2} + b^2.$

46. $x - \sqrt{xyz} + yz$ by $\sqrt{x} + \sqrt{yz}.$

47. $x\sqrt{x} - x\sqrt{y} + y\sqrt{x} - y\sqrt{y}$ by $\sqrt{x} + \sqrt{y}.$

DIVISION OF RADICALS

269. 1. What is the exponent of a in $a^5 + a^2$? in $a^{\frac{1}{2}} + a^{\frac{1}{3}}$? in $a^{\frac{1}{2}} \div a^{\frac{1}{3}}$? in $a^{\frac{1}{2}} + a^{\frac{1}{3}}$? in $\sqrt{a} \div \sqrt[3]{a}$?

2. When the fractional exponents indicate different roots, what must be done before one radical can be divided by another?

EXAMPLES

PROCESSES. — 1. $\sqrt{60} \div \sqrt{12} = \sqrt{5}$

$$2. \sqrt[3]{2} \div \sqrt{2} = \sqrt[6]{4} \div \sqrt[6]{8} = \sqrt[6]{\frac{4}{8}} = \sqrt[6]{\frac{1}{2}} = \frac{1}{\sqrt[6]{2}}$$

$$3. \sqrt[3]{x^2} \div \sqrt[4]{y} = \frac{(x^2)^{\frac{1}{3}}}{(y)^{\frac{1}{4}}} = \frac{(x^2)^{\frac{1}{12}}}{(y^3)^{\frac{1}{12}}} = \left(\frac{x^2}{y^3}\right)^{\frac{1}{12}} = \sqrt[12]{\frac{x^2}{y^3}} = \frac{1}{y^{\frac{1}{6}}} \sqrt[12]{x^2 y^5}$$

RULE. — *If necessary, reduce the radicals to the same degree.*

To the quotient of the coefficients annex the quotient of the radical factors under the common radical sign, and reduce the result to its simplest form.

Find the quotient of

- | | |
|--|---|
| 4. $\sqrt{50} \div \sqrt{8}.$ | 12. $2\sqrt[3]{12} \div \sqrt{8}.$ |
| 5. $\sqrt{72} \div 2\sqrt{6}.$ | 13. $\sqrt[3]{ax} \div \sqrt{xy}.$ |
| 6. $4\sqrt{5} \div \sqrt{40}.$ | 14. $\sqrt{2ab^3} \div \sqrt[4]{a^4b^4}.$ |
| 7. $6\sqrt{7} \div \sqrt{126}.$ | 15. $\sqrt[3]{a^2x^2} \div \sqrt{2ax}.$ |
| 8. $\sqrt[3]{4} \div \sqrt{2}.$ | 16. $\sqrt[3]{9a^2b^3} \div \sqrt{3ab}.$ |
| 9. $7\sqrt[3]{135} \div \sqrt[3]{9}.$ | 17. $\sqrt[4]{4x^2y^3} \div \sqrt[3]{2xy}.$ |
| 10. $7\sqrt{75} \div 5\sqrt{28}.$ | 18. $\sqrt{a-b} \div \sqrt{a+b}.$ |
| 11. $\sqrt[3]{16} \div \sqrt[6]{32}.$ | 19. $3\sqrt[3]{\frac{2}{3}} \div \sqrt{\frac{2}{3}}.$ |
| 20. Divide $\sqrt{15} - \sqrt{3}$ by $\sqrt{3}.$ | |
| 21. Divide $\sqrt{6} - 2\sqrt{3} + 4$ by $\sqrt{2}.$ | |
| 22. Divide $\sqrt{2} + 2 + \frac{1}{8}\sqrt{42}$ by $\frac{1}{8}\sqrt{6}.$ | |
| 23. Divide $5\sqrt{2} + 5\sqrt{3}$ by $\sqrt{10} + \sqrt{15}.$ | |
| 24. Divide $5 + 5\sqrt{30} + 36$ by $\sqrt{5} + 2\sqrt{6}.$ | |

INVOLUTION AND EVOLUTION OF RADICALS

270. In finding powers and roots of radicals it is frequently convenient to use fractional exponents.

EXAMPLES

1. What is the cube of $2\sqrt{ax^3}$?

SOLUTION. $(2\sqrt{ax^3})^3 = 2^3(ax^3)^{\frac{3}{2}} = 8a^{\frac{3}{2}}x^{\frac{9}{2}} = 8\sqrt{a^3x^9} = 8ax^4\sqrt{ax}$.

2. What is the square of $3\sqrt[6]{x^5}$?

SOLUTION. $(3\sqrt[6]{x^5})^2 = 9(x^5)^{\frac{2}{3}} = 9x^{\frac{10}{3}} = 9\sqrt[3]{x^5} = 9x\sqrt[3]{x^2}$.

3. What is the cube of $\sqrt{2} + 1$?

SOLUTION

$$\begin{aligned}(\sqrt{2} + 1)^3 &= (\sqrt{2})^3 + 3(\sqrt{2})^2 \cdot 1 + 3\sqrt{2} \cdot 1^2 + 1^3 \\&= 2\sqrt{2} + 6 + 3\sqrt{2} + 1 \\&= 7 + 5\sqrt{2}.\end{aligned}$$

In such cases expand by the binomial formula.

Square:	Cube:	Involve as indicated:
4. $3\sqrt{ab}$.	9. $2\sqrt{5}$.	14. $(-2\sqrt{2ab})^4$.
5. $2\sqrt[3]{3x}$.	10. $3\sqrt{2}$.	15. $(-\sqrt{2}\sqrt[6]{x})^3$.
6. $x\sqrt[3]{2x^3}$.	11. $2\sqrt[3]{a^2}$.	16. $(-\sqrt{2}\sqrt[3]{ax^3})^4$.
7. $n^2\sqrt[4]{4b}$.	12. $\sqrt[4]{a^2b^3}$.	17. $(-2\sqrt{x}\sqrt[3]{y})^5$.
8. $a\sqrt[4]{a^3b}$.	13. $\sqrt[6]{4n^3}$.	18. $(-3a^{\frac{n}{2}}x^{\frac{n}{3}})^6$.

Expand:

- | | | |
|--------------------------|----------------------------------|---------------------------------|
| 19. $(2 + \sqrt{6})^3$. | 22. $(2 - \sqrt{3})^3$. | 25. $(\sqrt{x} \pm 1)^2$. |
| 20. $(2 + \sqrt{2})^2$. | 23. $(\sqrt{7} - \sqrt{6})^2$. | 26. $(\sqrt{a} - \sqrt{b})^3$. |
| 21. $(2 + \sqrt{5})^3$. | 24. $(2\sqrt{2} - \sqrt{3})^2$. | 27. $\sqrt{x} \pm 1)^3$. |

28. What is the cube root of $-27\sqrt{ax}$?

SOLUTION. $\sqrt[3]{-27\sqrt{ax}} = (-27)^{\frac{1}{3}}(ax)^{\frac{1}{6}} = -3\sqrt[6]{ax}$.

29. What is the fourth root of $\sqrt{2x}$?

SOLUTION. $\sqrt[4]{\sqrt{2x}} = [(2x)^{\frac{1}{2}}]^{\frac{1}{4}} = (2x)^{\frac{1}{8}} = \sqrt[8]{2x}$.

Find the square root of

Find the cube root of

30. $\sqrt{2}$.

33. $\sqrt[3]{x^a}$.

36. $\sqrt{2x}$.

39. $-27\sqrt{x^5}$.

31. $\sqrt[3]{5}$.

34. $\sqrt[5]{x^{12}}$.

37. $\sqrt{7a^3}$.

40. $-64\sqrt[5]{a^3y^3}$.

32. $\sqrt{x^3}$.

35. $\sqrt[n]{a^nx^2}$.

38. $\sqrt[4]{8m^3x^5}$.

41. $-\sqrt{a^nb^n}$.

Simplify the following indicated roots:

42. $\sqrt{\sqrt[3]{4a^2x^4}}$.

44. $(\sqrt{8a^3x^5})^{\frac{1}{2}}$.

46. $\sqrt{\left(\frac{x^nb^2}{a^{-2}y^m}\right)^{\frac{2}{n}}}$.

43. $\sqrt[3]{\sqrt{a^{12}x^4}}$.

45. $(\sqrt{x^my^m})^{\frac{1}{mn}}$.

RATIONALIZATION

271. How may the fraction $\frac{1}{\sqrt{2}}$ be reduced to the form $\frac{\sqrt{2}}{2}$?

$$\frac{3}{\sqrt{2}} \text{ to } \frac{3\sqrt{2}}{2} \quad \frac{2}{\sqrt{5}} \text{ to } \frac{2\sqrt{5}}{5} \quad \frac{a}{\sqrt{x}} \text{ to } \frac{a\sqrt{x}}{x}$$

272. The process of rendering a surd expression rational is called **Rationalization**.

The factor by which a surd expression is multiplied to render it rational is called the **Rationalizing Factor**.

The denominator of $\frac{2}{\sqrt{3}}$ is rendered rational by multiplying it by $\sqrt{3}$.

Thus, $\frac{2}{\sqrt{3}} = \frac{2\sqrt{3}}{3}$. $\sqrt{3}$ is a rationalizing factor for the denominator.

273. A binomial, one or both of whose terms are surds, is called a **Binomial Surd**.

$2 + \sqrt{3}$, $\sqrt{x} + \sqrt{y}$, and $a^{\frac{1}{2}} - b^{\frac{1}{2}}$ are binomial surds.

274. Two binomial surds of the second degree whose product is rational are called **Conjugate Surds**.

$\sqrt{a} + \sqrt{b}$ and $\sqrt{a} - \sqrt{b}$ are conjugate surds; also $a + \sqrt{b}$ and $a - \sqrt{b}$.

Conjugate surds differ only in the sign of one of the terms.

275. PRINCIPLE. — *A binomial surd of the second degree may be rationalized by multiplying it by its conjugate.*

Thus, the product of $a + \sqrt{b}$ and $a - \sqrt{b}$ is $a^2 - b$.

EXAMPLES

1. Find the simplest rationalizing factor for $\sqrt{3x}$.

PROCESS

$$\sqrt{3x} \times \sqrt{3x} = \sqrt{9x^2} = 3x$$

$\therefore \sqrt{3x}$ is the simplest rationalizing factor.

EXPLANATION. — Since the number under the radical sign is to be changed so that the indicated root may be extracted, $\sqrt{3x}$ must be multiplied by some factor that will render the number under the radical sign a perfect square. Since the simplest factor by which $3x$ can be multiplied so that the product shall be a perfect square is $3x$, the simplest rationalizing factor for $\sqrt{3x}$ is $\sqrt{3x}$.

2. Find the simplest rationalizing factor for $\sqrt[3]{4a}$.

SOLUTION.
$$\sqrt[3]{4a} \times \sqrt[3]{2a^2} = \sqrt[3]{8a^3} = 2a.$$

$\therefore \sqrt[3]{2a^2}$ is the simplest rationalizing factor.

Find the simplest rationalizing factor for

- | | | | |
|------------------|------------------|------------------------|---------------------------|
| 3. $\sqrt{3}$. | 6. $\sqrt{4a}$. | 9. $\sqrt[3]{4}$. | 12. $\sqrt{x+1}$. |
| 4. $\sqrt{6}$. | 7. $\sqrt{9x}$. | 10. $\sqrt[3]{a^2}$. | 13. $\sqrt[3]{a-b}$. |
| 5. \sqrt{ax} . | 8. $\sqrt{8}$. | 11. $\sqrt[3]{mn^2}$. | 14. $\sqrt[3]{(a+b)^2}$. |

15. Find the simplest rationalizing factor for $\sqrt{5} + \sqrt{3}$.

PROCESS

$$(\sqrt{5} + \sqrt{3})(\sqrt{5} - \sqrt{3}) = (\sqrt{5})^2 - (\sqrt{3})^2 = 5 - 3 = 2$$

$\therefore \sqrt{5} - \sqrt{3}$ is the simplest rationalizing factor.

EXPLANATION. — Since $(\sqrt{5})^2 = 5$ and $(\sqrt{3})^2 = 3$, $\sqrt{5} + \sqrt{3}$ may be rationalized by multiplying it by some factor that will give the square of each term, but no other terms. Since the product of the sum and difference of two numbers is equal to the difference of their squares, the simplest rationalizing factor for $\sqrt{5} + \sqrt{3}$ is its conjugate, $\sqrt{5} - \sqrt{3}$ (Prin.).

Find the simplest rationalizing factor for

16. $\sqrt{3} + \sqrt{2}$.

18. $2 + \sqrt{3}$.

20. $a - 2\sqrt{b}$.

17. $\sqrt{3} - \sqrt{2}$.

19. $\sqrt{a} + \sqrt{x}$.

21. $3\sqrt{x} + 2y$.

22. Rationalize the denominator of $\frac{3}{\sqrt{12}}$.

SOLUTION. $\frac{3}{\sqrt{12}} = \frac{3 \times \sqrt{3}}{\sqrt{12} \times \sqrt{3}} = \frac{3\sqrt{3}}{\sqrt{36}} = \frac{3}{2}\sqrt{3} = \frac{1}{2}3\sqrt{3}$.

Rationalize the denominators:

23. $\frac{1}{\sqrt{3}}$.

27. $\frac{2}{\sqrt[3]{4}}$.

31. $\frac{\sqrt{6}}{\sqrt[3]{12}}$.

24. $\frac{2}{\sqrt{5}}$.

28. $\frac{2\sqrt{a}}{\sqrt{by}}$.

32. $\frac{\sqrt{a}}{\sqrt[3]{ax^3}}$.

25. $\sqrt{\frac{2}{3}}$.

29. $\frac{1}{\sqrt{x^j}}$.

33. $\frac{\sqrt{a+b}}{\sqrt{a-b}}$.

26. $\frac{\sqrt{a}}{\sqrt{b}}$.

30. $\frac{ax}{\sqrt{2ax}}$.

34. $\frac{\sqrt{x-2}}{\sqrt{x+2}}$.

35. Rationalize the denominator of $\frac{\sqrt{7} - \sqrt{3}}{\sqrt{7} + \sqrt{3}}$.

SOLUTION

$$\frac{\sqrt{7} - \sqrt{3}}{\sqrt{7} + \sqrt{3}} = \frac{(\sqrt{7} - \sqrt{3})(\sqrt{7} - \sqrt{3})}{(\sqrt{7} + \sqrt{3})(\sqrt{7} - \sqrt{3})} = \frac{7 - 2\sqrt{21} + 3}{7 - 3} = \frac{10 - 2\sqrt{21}}{4} = \frac{5 - \sqrt{21}}{2}.$$

Rationalize the denominators:

36. $\frac{\sqrt{3} + \sqrt{2}}{\sqrt{3} - \sqrt{2}}$.

38. $\frac{\sqrt{2} - 1}{\sqrt{2} + \sqrt{3}}$.

40. $\frac{1 - \sqrt{7}}{\sqrt{8} - \sqrt{7}}$.

37. $\frac{5}{\sqrt{5} - \sqrt{3}}$.

39. $\frac{5 - 3\sqrt{2}}{2 - \sqrt{2}}$.

41. $\frac{\sqrt{x} + \sqrt{y}}{\sqrt{x} - \sqrt{y}}$.

42. $\frac{4\sqrt{2} + 6\sqrt{3}}{3\sqrt{3} - 2\sqrt{2}}$.

44. $\frac{x - \sqrt{x^2 - 1}}{x + \sqrt{x^2 - 1}}$.

43. $\frac{\sqrt{x+1} - 2}{\sqrt{x+1} + 2}$.

45. $\frac{a\sqrt{a} - \sqrt{x+1}}{\sqrt{a^3} + \sqrt{x+1}}$.

$$46. \frac{\sqrt{x+y} - \sqrt{x-y}}{\sqrt{x+y} + \sqrt{x-y}}.$$

$$47. \frac{\sqrt{a^2+a+1} - 1}{\sqrt{a^2+a+1} + 1}.$$

$$48. \text{ Find the approximate value of } \frac{5}{\sqrt{3}}.$$

SOLUTION

$$\frac{5}{\sqrt{3}} = \frac{5\sqrt{3}}{3} = \frac{5 \times 1.73205}{3} = 2.88675.$$

In solving the following examples, the work will be lessened by observing that $\sqrt{2} = 1.41421$, $\sqrt{3} = 1.73205$, and $\sqrt{5} = 2.23607$.

Find the approximate value of

$$49. \frac{5}{\sqrt{2}}.$$

$$52. \frac{10}{\sqrt{45}}.$$

$$55. \frac{8 - \sqrt{3}}{2 - \sqrt{3}}.$$

$$50. \frac{2}{\sqrt{5}}.$$

$$53. \frac{15}{\sqrt{50}}.$$

$$56. \frac{1 + \sqrt{2}}{2 - \sqrt{2}}.$$

$$51. \frac{6}{\sqrt{8}}.$$

$$54. \frac{1}{\sqrt{125}}.$$

$$57. \frac{5 + 2\sqrt{5}}{5 - 2\sqrt{5}}.$$

$$58. \text{ Simplify } \frac{\sqrt{2} - \sqrt{3} - \sqrt{5}}{\sqrt{2} + \sqrt{3} + \sqrt{5}}.$$

SOLUTION

$$\text{Since } (\sqrt{2} + \sqrt{3}) - \sqrt{5} = (\sqrt{2} - \sqrt{5}) + \sqrt{3},$$

$$\begin{aligned} \frac{\sqrt{2} - \sqrt{3} - \sqrt{5}}{\sqrt{2} + \sqrt{3} + \sqrt{5}} &= \frac{(\sqrt{2} - \sqrt{5}) - \sqrt{3}}{(\sqrt{2} + \sqrt{3}) + \sqrt{5}} \times \frac{(\sqrt{2} - \sqrt{5}) + \sqrt{3}}{(\sqrt{2} + \sqrt{3}) - \sqrt{5}} \\ &= \frac{2 - 2\sqrt{10} + 5 - 3}{2 + 2\sqrt{6} + 3 - 5} = \frac{4 - 2\sqrt{10}}{2\sqrt{6}} \\ &= \frac{2 - \sqrt{10}}{\sqrt{6}} = \frac{2\sqrt{6} - 2\sqrt{15}}{6} = \frac{\sqrt{6} - \sqrt{15}}{3}. \end{aligned}$$

Rationalize the denominators:

$$59. \frac{\sqrt{2} - \sqrt{5} - \sqrt{7}}{\sqrt{2} + \sqrt{5} + \sqrt{7}}.$$

$$61. \frac{1}{\sqrt{2} + \sqrt{3} + \sqrt{5}}.$$

$$60. \frac{\sqrt{3} + \sqrt{2}}{\sqrt{3} + \sqrt{2} - \sqrt{6}}.$$

$$62. \frac{2\sqrt{2} - 3\sqrt{3} + 4\sqrt{5}}{\sqrt{2} + \sqrt{3} - \sqrt{5}}.$$

276. To find the square root of a binomial surd.

$$(\sqrt{2} + \sqrt{6})^2 = 2 + 2\sqrt{12} + 6 = 8 + 2\sqrt{12};$$

$$(\sqrt{4} + \sqrt{3})^2 = 4 + 2\sqrt{12} + 3 = 7 + 2\sqrt{12};$$

or

$$(2 + \sqrt{3})^2 = 4 + 4\sqrt{3} + 3 = 7 + 4\sqrt{3}.$$

1. Since $(\sqrt{2} + \sqrt{6})^2 = 8 + 2\sqrt{12}$, what is the square root of $8 + 2\sqrt{12}$?

2. How does the product of the terms of the square root of $8 + 2\sqrt{12}$ compare with the irrational term $2\sqrt{12}$?

3. How does the sum of the squares of the terms of the square root compare with the rational term 8?

4. How, then, may the square root of $8 + 2\sqrt{12}$ be found from the terms of $8 + 2\sqrt{12}$?

5. How may the terms of the square root of $7 + 2\sqrt{12}$, or the equivalent expression $7 + 4\sqrt{3}$, be found? After the irrational term is divided by 2, what two factors of the result are selected for the terms of the root?

277. A surd of the second degree is called a **Quadratic Surd**.

\sqrt{x} , $4\sqrt{x}$, $\sqrt{x} + \sqrt{y}$, and $3 + 2\sqrt{5}$ are quadratic surds.

278. PRINCIPLE. — *The terms of the square root of a quadratic binomial surd that is a perfect square may be obtained by dividing the irrational term by 2 and then separating the quotient into two factors, the sum of whose squares is the rational term.*

EXAMPLES

1. Find the square root of $14 + 8\sqrt{3}$.

SOLUTION

Since, if $14 + 8\sqrt{3}$ is the square of a binomial quadratic surd, the irrational term $8\sqrt{3}$ is *twice* the product of the terms of the root (Prin.), $4\sqrt{3}$, or $\sqrt{48}$, is the product of the terms of the binomial surd. Since the two factors of $\sqrt{48}$, the sum of whose squares is 14, are $\sqrt{6}$ and $\sqrt{8}$, the required square root is equal to $\sqrt{6} + \sqrt{8}$.

$$\therefore \sqrt{14 + 8\sqrt{3}} = \sqrt{6} + \sqrt{8}.$$

2. Find the square root of $11 - 6\sqrt{2}$.

SOLUTION

$$\begin{aligned}\sqrt{11 - 6\sqrt{2}} &= \sqrt{11 - 2\sqrt{18}} \\ &= 3 - \sqrt{2}.\end{aligned}$$

Find the square root of each of the following:

- | | |
|------------------------|--------------------------------|
| 3. $12 + 2\sqrt{35}$. | 11. $12 + 4\sqrt{5}$. |
| 4. $16 - 2\sqrt{60}$. | 12. $11 + 4\sqrt{7}$. |
| 5. $15 + 2\sqrt{26}$. | 13. $12 - 6\sqrt{3}$. |
| 6. $16 - 2\sqrt{55}$. | 14. $17 + 12\sqrt{2}$. |
| 7. $11 + 2\sqrt{30}$. | 15. $15 - 6\sqrt{6}$. |
| 8. $7 - 2\sqrt{10}$. | 16. $18 + 6\sqrt{5}$. |
| 9. $3 - 2\sqrt{2}$. | 17. $a^2 + b + 2a\sqrt{b}$. |
| 10. $6 + 2\sqrt{5}$. | 18. $2a - 2\sqrt{a^2 - b^2}$. |

PROPERTIES OF QUADRATIC SURDS

279. *The square root of a rational number cannot be partly rational and partly a quadratic surd.*

For, if possible, let

$$\sqrt{y} = \sqrt{b} \pm m.$$

By squaring,

$$y = b \pm 2m\sqrt{b} + m^2,$$

and

$$\sqrt{b} = \pm \frac{y - m^2 - b}{2m};$$

that is, a surd is equal to a rational number, which is impossible. Therefore, \sqrt{y} cannot be equal to $\sqrt{b} \pm m$.

280. *In any equation containing rational numbers and quadratic surds, as $a + \sqrt{b} = x + \sqrt{y}$, the rational parts are equal, and also the irrational parts.*

Let

$$a + \sqrt{b} = x + \sqrt{y}. \quad (1)$$

Since a and x are both rational, if possible, let

$$a = x \pm m. \quad (2)$$

Then,

$$x \pm m + \sqrt{b} = x + \sqrt{y}, \quad (3)$$

and

$$\sqrt{y} = \sqrt{b} \pm m. \quad (4)$$

Since, § 279, equation (4) is impossible, $a = x \pm m$ is impossible; that is, a is neither greater nor less than x .

Therefore, $a = x$, and, Eq. (1), $\sqrt{b} = \sqrt{y}$.

Hence, if $a + \sqrt{b} = x + \sqrt{y}$, $a = x$, and $\sqrt{b} = \sqrt{y}$.

281. If $\sqrt{a + \sqrt{b}} = \sqrt{x} + \sqrt{y}$, then $\sqrt{a - \sqrt{b}} = \sqrt{x} - \sqrt{y}$, when a, b, x , and y are rational and $a > \sqrt{b}$.

For, squaring, $a + \sqrt{b} = x + 2\sqrt{xy} + y$.

Therefore, § 280, $a = x + y$, and $\sqrt{b} = 2\sqrt{xy}$.

Hence, $a - \sqrt{b} = x + y - 2\sqrt{xy}$.

Whence, $\sqrt{a - \sqrt{b}} = \sqrt{x} - \sqrt{y}$.

EXAMPLES

1. Find the square root of $21 + 6\sqrt{10}$.

SOLUTION

Let $\sqrt{x} + \sqrt{y} = \sqrt{21 + 6\sqrt{10}}$. (1)

Then, § 281, $\sqrt{x} - \sqrt{y} = \sqrt{21 - 6\sqrt{10}}$. (2)

Multiplying, $x - y = \sqrt{441 - 360} = \sqrt{81}$,
or $x - y = 9$. (3)

Squaring (1), $x + 2\sqrt{xy} + y = 21 + 6\sqrt{10}$.

Therefore, § 280 $x + y = 21$. (4)

Solving (4) and (3), $x = 15, y = 6$.

$$\therefore \sqrt{x} = \sqrt{15}, \sqrt{y} = \sqrt{6}.$$

Hence, the square root of $21 + 6\sqrt{10}$ is $\sqrt{15} + \sqrt{6}$.

Find the square root of

- | | | |
|------------------------|--------------------------|---------------------------------|
| 2. $25 + 10\sqrt{6}$. | 8. $16 + 6\sqrt{7}$. | 14. $2 + \sqrt{3}$. |
| 3. $19 + 6\sqrt{2}$. | 9. $21 - 8\sqrt{5}$. | 15. $6 + \sqrt{35}$. |
| 4. $45 + 30\sqrt{2}$. | 10. $47 - 12\sqrt{11}$. | 16. $1 + \frac{2}{3}\sqrt{2}$. |
| 5. $35 - 14\sqrt{6}$. | 11. $56 + 32\sqrt{3}$. | 17. $2 + \frac{4}{5}\sqrt{6}$. |
| 6. $11 + 6\sqrt{2}$. | 12. $35 - 12\sqrt{6}$. | 18. $30 + 20\sqrt{2}$. |
| 7. $24 - 8\sqrt{5}$. | 13. $56 - 12\sqrt{3}$. | 19. $18 - 6\sqrt{5}$. |

RADICAL EQUATIONS

282. An equation involving an irrational root of an unknown number is called an **Irrational**, or **Radical Equation**.

$x^{\frac{1}{2}} = 3$, $\sqrt{x+1} = \sqrt{x-4} + 1$, and $\sqrt[3]{x-1} = 2$ are radical equations.

283. An equation containing quadratic surds involving x may be rationalized with respect to x by writing all the terms in the first member and multiplying both members by the proper rationalizing factor.

To rationalize $\sqrt{x-3} = 2.$ (1)

Transposing, $\sqrt{x-3} - 2 = 0.$ (2)

Multiplying by the conjugate surd,
 $(\sqrt{x-3} - 2)(\sqrt{x-3} + 2) = 0,$ (3)

or $x - 3 - 4 = 0.$ (4)

The rationalization of such equations, however, is more conveniently accomplished by the process of squaring.

Thus, by squaring (1), (4) may be obtained in the form

$$x - 3 = 4. \quad (5)$$

It is evident, then, that squaring an equation is equivalent to multiplying both members by the same unknown expression, an operation likely to introduce roots (§ 196). The roots introduced, if any, when an equation is rationalized are those of the equation or equations formed by placing each rationalizing factor equal to zero.

Thus, in squaring the equation $\sqrt{x-3} = 2$, the root of $\sqrt{x-3} = -2$ is introduced. But if $\sqrt{4}$ is taken to mean either $+2$ or -2 , this equation has the same root as the given equation and no root has been introduced.

Repeated squaring is often necessary to free an equation of quadratic surds involving x . This corresponds to repeated rationalization with respect to particular surds.

Thus, $\sqrt{x} - 5 = -\sqrt{x+5}.$
 Squaring, $x - 10\sqrt{x} + 25 = +(x+5) = x+5.$
 Simplifying, $\sqrt{x} = 2.$
 Squaring, $x = 4.$

Or, transposing in the given equation,

$$\sqrt{x} - 5 + \sqrt{x+5} = 0.$$

Rationalizing with respect to $\sqrt{x+5}$,

$$(\sqrt{x}-5+\sqrt{x+5})(\sqrt{x}-5-\sqrt{x+5})=0.$$

Expanding, $x-10\sqrt{x}+25-(x+5)=0.$

Simplifying, $\sqrt{x}-2=0.$

Rationalizing, $(\sqrt{x}-2)(\sqrt{x}+2)=0,$

or $x-4=0.$

In squaring the first time, the factor $\sqrt{x}-5-\sqrt{x+5}$ is introduced; and in squaring the second time the factor $\sqrt{x}+2$, which is $\frac{1}{10}$ of the product $(\sqrt{x+5}+\sqrt{x+5})(\sqrt{x+5}-\sqrt{x+5})$, or $10\sqrt{x}+20$, is introduced.

284. It follows from the preceding discussion that:

If a radical equation is rationalized by multiplying by a rationalizing factor or by squaring, the resulting equation has all the roots of the given equation.

Whether the given radical equation has all the roots of the rational equation depends upon the method agreed upon of *verifying* radical equations.

As illustrated above, each of the equations

$$\sqrt{x}-5+\sqrt{x+5}=0, \quad (1)$$

$$\sqrt{x}-5-\sqrt{x+5}=0, \quad (2)$$

$$\sqrt{x}+5+\sqrt{x+5}=0, \quad (3)$$

and $\sqrt{x}+5-\sqrt{x+5}=0 \quad (4)$

is rationalized by finding the product of them all, which is $x-4=0$. Hence, the equation $x-4=0$ has the roots of the four equations; that is, each equation has the root $x=4$, or has no root.

If $x=4$, $\sqrt{x}=\sqrt{4}$ and $\sqrt{x+5}=\sqrt{9}$. If $\sqrt{4}$ is either $+2$ or -2 and $\sqrt{9}$ is $+3$ or -3 , the equations are verified as follows:

(1) becomes $2-5+3=0,$

(2) becomes $2-5-(-3)=0,$

(3) becomes $-2+5+(-3)=0,$

(4) becomes $-2+5-3=0.$

To prevent confusion in making numerical substitutions, it is customary to regard only the positive or principal square root in expressions like $\sqrt{4}$, $\sqrt{9}$, $\sqrt{5}$, etc. (See § 225.)

For example, by common agreement $+\sqrt{9}$ means $+(+3)$, or $+3$, $-\sqrt{9}$ means $-(+3)$, or -3 ; $+\sqrt{5}$ means the positive square root of 5; etc. With this understanding equations (2), (3), and (4) cannot be verified for $x=4$, and since they have no other root, they may in this sense be regarded as *impossible* equations.

When the equations given in this section have been freed from the radical signs, the resulting equations will be found to be simple equations. Other varieties of radical equations are treated subsequently.

EXAMPLES

1. Given $\sqrt{2x} + 4 = 10$, to find the value of x .

SOLUTION

$$\sqrt{2x} + 4 = 10.$$

Transposing,

$$\sqrt{2x} = 6.$$

Squaring,

$$2x = 36.$$

$$\therefore x = 18.$$

2. Given $\sqrt{x-7} + \sqrt{x} = 7$, to find the value of x .

SOLUTION

$$\sqrt{x-7} + \sqrt{x} = 7.$$

Transposing,

$$\sqrt{x-7} = 7 - \sqrt{x}.$$

Squaring,

$$x - 7 = 49 - 14\sqrt{x} + x.$$

Transposing and combining, $14\sqrt{x} = 56.$

Dividing by 14,

$$\sqrt{x} = 4.$$

Squaring,

$$x = 16.$$

3. Given $\sqrt{14 + \sqrt{1 + \sqrt{x + 8}}} = 4$, to find the value of x .

SOLUTION

$$\sqrt{14 + \sqrt{1 + \sqrt{x + 8}}} = 4.$$

Squaring,

$$14 + \sqrt{1 + \sqrt{x + 8}} = 16.$$

Transposing, etc.,

$$\sqrt{1 + \sqrt{x + 8}} = 16 - 14 = 2.$$

Squaring,

$$1 + \sqrt{x + 8} = 4.$$

Transposing, etc.,

$$\sqrt{x + 8} = 4 - 1 = 3.$$

Squaring,

$$x + 8 = 9.$$

$$\therefore x = 9 - 8 = 1.$$

VERIFICATION. $\sqrt{14 + \sqrt{1 + \sqrt{1 + 8}}} = \sqrt{14 + \sqrt{1 + 3}}$
 $= \sqrt{14 + 2} = 4.$

4. Given $\frac{2\sqrt{ax}-b}{2\sqrt{ax}+b} = \frac{3\sqrt{ax}-2b}{3\sqrt{ax}+3b}$, to find the value of x .

SOLUTION

$$\frac{2\sqrt{ax}-b}{2\sqrt{ax}+b} = \frac{3\sqrt{ax}-2b}{3\sqrt{ax}+3b}$$

Reducing to mixed numbers, $1 - \frac{2b}{2\sqrt{ax}+b} = 1 - \frac{5b}{3\sqrt{ax}+3b}$.

Canceling, $-\frac{2b}{2\sqrt{ax}+b} = -\frac{5b}{3\sqrt{ax}+3b}$.

Dividing by $-b$, $\frac{2}{2\sqrt{ax}+b} = \frac{5}{3\sqrt{ax}+3b}$.

Clearing of fractions, etc., $6\sqrt{ax} - 10\sqrt{ax} = 5b - 6b$.

$$\therefore \sqrt{ax} = \frac{b}{4}$$

Squaring, etc.,

$$x = \frac{b^2}{16a}$$

SUGGESTIONS.—1. When the equation is free from fractions, transpose so that the radical term, if there is but one, or the more complex radical term, if there is more than one, may constitute one member of the equation; then raise each member to a power of the same degree as that radical. Simplify the result. If the equation is not freed from radicals by the first involution, proceed again as at first.

2. It is sometimes convenient to rationalize denominators before clearing of fractions or involving.

Solve the following equations:

- | | |
|----------------------------|--------------------------------------|
| 5. $\sqrt{x+11} = 4$. | 12. $1 + 2\sqrt{x} = 7 - \sqrt{x}$. |
| 6. $\sqrt{x+5} = 3$. | 13. $\sqrt{x+16} - \sqrt{x} = 2$. |
| 7. $\sqrt{x-a^2} = b$. | 14. $\sqrt{2x} - \sqrt{2x-15} = 1$. |
| 8. $\sqrt[3]{x-1} = 2$. | 15. $\sqrt{x^2+x+1} = 2-x$. |
| 9. $\sqrt[3]{x-a^3} = a$. | 16. $3\sqrt{x^2-9} = 3x-3$. |
| 10. $\sqrt[3]{x+b} = a$. | 17. $\sqrt{x+2} = \sqrt{x+32}$. |
| 11. $1 + \sqrt{x} = 5$. | 18. $5 - \sqrt{x+5} = \sqrt{x}$. |

19. $2\sqrt{x} - x = x - 8\sqrt{x}$. 25. $\sqrt{3x-5} + \sqrt{3x+7} = 6$.
20. $\sqrt{4x^2+6x-10} = 2x+4$. 26. $\sqrt{16x+3} + \sqrt{16x+8} = 5$.
21. $\sqrt{x^2-5x+7} + 2 = x$. 27. $\sqrt{9x+8} + \sqrt{9x-4} = 0$.
22. $4 - \sqrt{4-8x+9x^2} = 3x$. 28. $\sqrt{1+x\sqrt{x^2+12}} = 1+x$.
23. $\sqrt{2(1-x)(3-2x)} - 1 = 2x$. 29. $\sqrt{7+3\sqrt{5x-16}} - 4 = 0$.
24. $\sqrt{2x-1} + \sqrt{2x+4} = 5$. 30. $2x + \sqrt{4x^2 - \sqrt{16x^2-7}} = 1$.
31. $\sqrt{7 + \sqrt{1 + \sqrt{4 + \sqrt{1 + 2\sqrt{x}}}}} = 3$.
32. $\frac{5}{\sqrt{3x+2}} = \sqrt{3x+2} + \sqrt{3x-1}$.
33. $\frac{\sqrt{2x}+9}{\sqrt{2x}-7} = \frac{\sqrt{2x}+20}{\sqrt{2x}-12}$. 38. $\frac{\sqrt{2x}+6}{\sqrt{2x}+4} = \frac{\sqrt{2x}+2}{\sqrt{2x}+1}$.
34. $\frac{\sqrt{x}+18}{\sqrt{x}+2} = \frac{32}{\sqrt{x}+6} + 1$. 39. $\frac{\sqrt{11x} + \sqrt{2x+3}}{\sqrt{11x} - \sqrt{2x+3}} = \frac{8}{3}$.
35. $\frac{\sqrt{x-1}}{\sqrt{x+5}} = \frac{\sqrt{x-3}}{\sqrt{x-1}}$. 40. $\frac{2\sqrt{2x}+4}{2\sqrt{2x}-4} = \frac{3\sqrt{x+1}+9}{3\sqrt{x+1}-9}$.
36. $\frac{\sqrt{x}-6}{\sqrt{x}-1} = \frac{\sqrt{x}-8}{\sqrt{x}-5}$. 41. $\frac{\sqrt{\sqrt{5x}-9}}{\sqrt{\sqrt{5x}+11}} = \frac{\sqrt{\sqrt{5x}-21}}{\sqrt{\sqrt{5x}-16}}$.
37. $\frac{\sqrt{x-3}}{\sqrt{x+1}} = \frac{\sqrt{x-4}}{\sqrt{x-2}}$. 42. $\frac{\sqrt{4x+3} + 2\sqrt{x-1}}{\sqrt{4x+3} - 2\sqrt{x-1}} = 5$.
43. $\frac{\sqrt{x+1} - \sqrt{x-1}}{\sqrt{x+1} + \sqrt{x-1}} = \frac{1}{2}$.
44. $\frac{x-3}{\sqrt{x}-\sqrt{3}} = \frac{\sqrt{x}+\sqrt{3}}{2} + 2\sqrt{3}$.
45. $\frac{\sqrt{19x} + \sqrt{2x+11}}{\sqrt{19x} - \sqrt{2x+11}} = 2\frac{1}{8}$.
46. $2\sqrt{x} - \sqrt{4x-22} - \sqrt{2} = 0$.

$$47. 1 + \sqrt{(3-5x)^2 + 16} = 2(3-x).$$

$$48. \sqrt{x} + \sqrt{x - \sqrt{a^2 - x}} = \sqrt{a}.$$

$$49. \sqrt{x} + \sqrt{x - (a-b)^2} = a + b.$$

$$50. \sqrt{mn-x} - \sqrt{x}\sqrt{mn-1} = \sqrt{mn}\sqrt{1-x}.$$

$$51. a\sqrt{x} - b\sqrt{x} = a^2 + b^2 - 2ab.$$

$$52. \sqrt{5ax - 9a^3} + a = \sqrt{5ax}.$$

$$53. \sqrt{x+3a} = \frac{6a}{\sqrt{x+3a}} - \sqrt{x}.$$

$$54. \sqrt{2x} - \sqrt{2x-7} = \frac{3}{\sqrt{2x-7}}.$$

$$55. \sqrt{2x + \sqrt{10x+1}} = \sqrt{2x} + 1.$$

$$56. \frac{\sqrt{x+a} + \sqrt{x-a}}{\sqrt{x+a} - \sqrt{x-a}} = 2 + \frac{\sqrt{x^2 - a^2}}{a}.$$

$$57. \sqrt{x} + \sqrt{2x} + \sqrt{3x} = \sqrt{a}.$$

SOLUTION

$$\sqrt{x} + \sqrt{2x} + \sqrt{3x} = \sqrt{a}.$$

Factoring, $\sqrt{x}(1 + \sqrt{2} + \sqrt{3}) = \sqrt{a}.$

Multiplying by $1 + \sqrt{2} - \sqrt{3},$

$$\sqrt{x}(1 + 2\sqrt{2} + 2 - 3) = \sqrt{a}(1 + \sqrt{2} - \sqrt{3}).$$

$$\sqrt{x} \cdot 2\sqrt{2} = \sqrt{a}(1 + \sqrt{2} - \sqrt{3}).$$

Squaring, $8x = a(1 + \sqrt{2} - \sqrt{3})^2.$

$\therefore x = \frac{a}{8}(1 + \sqrt{2} - \sqrt{3})^2.$

$$58. \sqrt{2x} + \sqrt{3x} + \sqrt{5x} = \sqrt{m}.$$

$$59. \sqrt{2x} + \sqrt{3x} - \sqrt{5x} = \sqrt{c}.$$

$$60. \sqrt{x-a} + \sqrt{2(x-a)} = \sqrt{3x+a\sqrt{2}}.$$

$$61. \sqrt{x-1} + \sqrt{2x-2} = \sqrt{3x-3} + \sqrt{2}.$$

$$62. \sqrt{2x-3} + \sqrt{4x-6} = \sqrt{2x} + \sqrt{x}.$$

REVIEW

Reduce the following to their simplest forms :

$$1. \frac{6x^2 - 7x^2 - 5x}{9x^2 - 25x}.$$

$$7. \frac{x-y}{x+y} - \frac{y+x}{y-x} - \frac{4x^2y^2}{x^4-y^4}.$$

$$2. \frac{8x^2 + 18x - 5}{12x^2 + 5x - 2}.$$

$$8. \frac{\sqrt{2}-\sqrt{3}}{\sqrt{2}+\sqrt{3}} - \frac{\sqrt{2}+\sqrt{3}}{\sqrt{2}-\sqrt{3}}.$$

$$3. \frac{a^2x^2 - a\sqrt{x} + x}{\sqrt{x}}.$$

$$9. \frac{x+\sqrt{xy}+y}{\sqrt{x}+\sqrt{y}} + \frac{x\sqrt{x}+y\sqrt{y}}{x+y}.$$

$$4. \frac{a^2 - 2a\sqrt{b} + b}{a - \sqrt{b}}.$$

$$10. \frac{1}{\sqrt{a}+\sqrt{b}} - 1 + \frac{1}{\sqrt{a}-\sqrt{b}}.$$

$$5. \frac{\sqrt{2}-\sqrt{3}-\sqrt{5}}{\sqrt{2}+\sqrt{3}+\sqrt{5}}.$$

$$11. \frac{\sqrt{1+x^2}-\sqrt{1-x^2}}{\sqrt{1+x^2}+\sqrt{1-x^2}}.$$

$$6. \frac{2-\sqrt{5}}{2+\sqrt{5}} + \frac{2\sqrt{3}}{\sqrt{243}}.$$

$$12. \frac{1}{1-\sqrt{2}x} + \frac{1}{1+\sqrt{2}x} + \frac{1}{1-2x}.$$

$$13. \frac{x+\sqrt{x^2-a^2}}{x-\sqrt{x^2-a^2}} - \frac{x-\sqrt{x^2-a^2}}{x+\sqrt{x^2-a^2}}.$$

$$14. \frac{\sqrt{a+1}+\sqrt{a-1}}{\sqrt{a+1}-\sqrt{a-1}} - \frac{\sqrt{a+1}-\sqrt{a-1}}{\sqrt{a+1}+\sqrt{a-1}}.$$

$$15. \frac{a^2-2ax-3x^2}{3a^2+5ax+2x^2} \times \frac{6a^2+7ax+2x^2}{a^2-4ax+3x^2}.$$

$$16. \frac{a^2-b}{a^2-2a\sqrt{b}+b} \times \frac{a^2-4a\sqrt{b}+4b}{a^2+2a\sqrt{b}+b}.$$

$$17. \left(1 - \frac{a}{b}\right) \div \left(1 + \sqrt{\frac{a}{b}}\right).$$

$$19. 1 + \left(\sqrt{\frac{a}{b}} - \sqrt{\frac{b}{a}}\right) + \frac{\sqrt{ab}}{a+b}.$$

$$18. \frac{\frac{1+a+a^2}{1+\sqrt{a}+a}}{1-\sqrt{a}+a}.$$

$$20. \frac{\left(\frac{a}{\sqrt{x}} + \frac{\sqrt{x}}{a}\right)\left(\frac{a}{\sqrt{x}} - \frac{\sqrt{x}}{a}\right)}{\left(\frac{a}{\sqrt{x}} - \frac{\sqrt{x}}{a}\right)\left(\frac{a}{\sqrt{x}} - \frac{\sqrt{x}}{a}\right)}.$$

Expand:

- | | | |
|--|---|---------------------------------------|
| 21. $(a^3 - b^2)^3$. | 25. $(a^{-2} + a^{-1})^2$. | 31. $(a - \sqrt{b})^4$. |
| 22. $(2a - 3b)^4$. | 26. $(a^{-1} + b)^4$. | 32. $(\sqrt{x} + \sqrt{y})^6$. |
| 23. $\left(\frac{a}{3} - \frac{b}{2}\right)^3$. | 27. $(a^{\frac{1}{2}} - b^{\frac{1}{2}})^6$. | 33. $(\sqrt{2} - \sqrt{3})^4$. |
| 24. $\left(ax + \frac{1}{a}\right)^5$. | 28. $(a^{\frac{1}{2}} - b^{-\frac{1}{2}})^4$. | 34. $(\sqrt{5} - 2)^6$. |
| | 29. $(a^{-\frac{1}{2}} - b^{-\frac{1}{2}})^6$. | 35. $(\sqrt[3]{4} - \sqrt[3]{2})^3$. |
| | 30. $(a^{\frac{1}{2}} + b^{\frac{1}{2}})^6$. | 36. $(\sqrt{2} - \sqrt[3]{2})^6$. |

Extract the square root of

37. $\frac{9x^4}{4} + 3x^3 - x^2 - \frac{4x}{3} + \frac{4}{9}$.
38. $\frac{x^2}{4} + 4y^2 + \frac{z^2}{16} - 2xy + \frac{xz}{4} - yz$.
39. $a^2 + 12a^{\frac{1}{2}}b^{\frac{1}{2}} + 54ab + 108a^{\frac{1}{2}}b^{\frac{3}{2}} + 81b^2$.
40. $1 + 2\sqrt{x} - x - 2x\sqrt{x} + x^2$.
41. $a + 4b + 9c - 4\sqrt{ab} + 6\sqrt{ac} - 12\sqrt{bc}$.
42. $x^2 - 4x\sqrt{xy} + 6xy - 4y\sqrt{xy} + y^2$.
43. $x^2 - 12x^{\frac{1}{2}}y^{\frac{1}{2}} + 60x^{\frac{1}{2}}y - 160xy^{\frac{1}{2}} + 240x^{\frac{1}{2}}y^2$
 $- 192x^{\frac{1}{2}}y^{\frac{3}{2}} + 64y^3$.

Find the square root of

- | | |
|------------------------|--------------------------|
| 44. 81234169. | 48. $56 + 14\sqrt{15}$. |
| 45. 64064016. | 49. $47 - 12\sqrt{15}$. |
| 46. .00022801. | 50. $62 + 20\sqrt{6}$. |
| 47. .1 to four places. | 51. $51 - 36\sqrt{2}$. |

Extract the cube root of

52. $x^3 - 9x + 27x^{-1} - 27x^{-3}$.
53. $27x^3 + 27x^2 - 5 + \frac{1}{3x^3} - \frac{1}{27x^3}$.
54. $x^3 + 3x^2\sqrt{x} - 5x\sqrt{x} + 3\sqrt{x} - 1$.

55. Find the cube root of $2\sqrt{2} - 6\sqrt[3]{2} + 3\sqrt{2}\sqrt[3]{4} - 2$.
56. Find the cube root of $(a + 12b + 3c)\sqrt{a}$
 $-(6a + 8b + 6c)\sqrt{b} - (3a + 12b + c)\sqrt{c} + 12\sqrt{abc}$.
57. Extract the cube root of 510,082,399.
58. Extract the cube root of 1,042,590,744.
59. Extract the cube root of 2 to three decimal places.
60. Find the first four terms of $\sqrt{1+x-x^2}$.
61. Find the first three terms of $\sqrt[3]{1+x^3}$.
62. Find the fourth root of
 $a^3 - 4a^4\sqrt{ab^{-1}} + 6a^3b^{-1} - 4ab^{-1}\sqrt{ab^{-1}} + b^{-2}$.
63. Find the sixth root of
 $8 - 48\sqrt{a} + 120a - 160a\sqrt{a} + 120a^2 - 48a^2\sqrt{a} + 8a^3$.

If $a^m \times a^n = a^{m+n}$ for all values of m and n , show that

64. $a^{-2} = \frac{1}{a^2}$. 67. $(ab)^0 = 1$.
65. $a^{\frac{1}{2}} = \sqrt{a} = (\sqrt{a})^2$. 68. $(abc)^3 = a^3b^3c^3$.
66. $2a^{-\frac{1}{2}} = \frac{2\sqrt[3]{a^2}}{a}$. 69. $\left(\frac{a}{b}\right)^3 = \frac{a^3}{b^3}$.

Find the values of the following:

70. $16^{\frac{1}{2}}$. 73. $(a^4x^4)^{\frac{1}{2}}$. 76. $\left(\frac{32}{248}\right)^{-\frac{1}{2}}$.
71. $27^{\frac{1}{3}}$. 74. $(b^2y^4)^{-\frac{3}{2}}$. 77. $\left(\frac{36}{144}\right)^{-\frac{1}{2}}$.
72. $8^{-\frac{1}{2}}$. 75. $(a^nb^n)^{-\frac{1}{n}}$. 78. $\left(-\frac{8}{27}\right)^{-\frac{1}{3}}$.
79. For what values of n is $(a-b)^n = (b-a)^n$?

Simplify and express with positive exponents:

80. $(36a^{-3} + 25a^{-2})^{-\frac{1}{2}}$. 83. $(\sqrt{a^3x^{-3}} + \sqrt[3]{a^2x^{-2}})^{\frac{2}{3}}$.
81. $(8a^2x^3 \times 64a^{-4}x^{-5})^{-\frac{1}{2}}$. 84. $(\sqrt{a^{-1}b^4} + \sqrt{a^2b})^{-\frac{1}{2}}$.
82. $(a^{\frac{1}{2}}b^{\frac{1}{4}})^{\frac{1}{2}} + (a^{\frac{1}{2}}b^{\frac{1}{4}})^2$. 85. $(\sqrt{a} \div \sqrt[3]{a}) + \sqrt[4]{a}$.

$$86. \frac{a-b}{a^{\frac{1}{3}}-b^{\frac{1}{3}}} - \frac{a+b}{a^{\frac{1}{3}}+b^{\frac{1}{3}}} + \frac{2a^{\frac{1}{3}}b}{a^{\frac{2}{3}}-b^{\frac{2}{3}}}.$$

$$87. \frac{1+a^{-1}b}{1-a^{-1}b} + \left(\frac{1+ab^{-1}+a^2b^{-2}}{1-ab^{-1}+a^2b^{-2}} \times \frac{1+a^{-2}b^3}{1-a^{-2}b^3} \right).$$

Solve the following equations:

$$88. \frac{x+1}{x-1} - \frac{x-1}{x+1} = \frac{3-5x}{1-x^2}.$$

$$89. \frac{7-2x}{10} - \frac{2x-1}{5} + \frac{1}{2}x = \frac{5x-6}{2x} - \frac{17+3x}{30}.$$

$$90. \frac{4x-17}{9} - \frac{3\frac{2}{3}-22x}{33} = x - \frac{6}{x} \left(1 - \frac{x^2}{54} \right).$$

$$91. \begin{cases} \frac{3x-5y}{3} - \frac{2x-8y-9}{12} = \frac{y}{2} + \frac{7}{12}, \\ \frac{7}{2} \left(\frac{x}{7} + \frac{y}{4} + 1\frac{1}{8} \right) - \frac{10}{3} \left(4x - \frac{y}{8} - 24 \right) = 0. \end{cases}$$

$$92. \begin{cases} 3x+1=2y, \\ (x+5)(y+7)=(x+1)(y-9)+112. \end{cases}$$

$$93. \begin{cases} x-y=3, \\ (x+1)(x+2)-(x-2)(x+1)=11y+2. \end{cases}$$

Simplify and express with positive exponents:

$$94. \frac{\left(\frac{a}{27} + \frac{a^{-2}}{8} \right)^{-\frac{2}{3}} - x^2}{\frac{3a^{-1}+2x}{2}}.$$

$$97. \left[\frac{x^{-\frac{1}{2}}y^{-\frac{2}{3}}}{x^{-\frac{1}{3}}y^{-1}} + \frac{x^{-2}y^2}{(xy)^{-3}} \right]^{-3}.$$

$$98. \{ (a^{\frac{1}{2}}b^{\frac{5}{3}})^{\frac{1}{2}} + (a^{-\frac{1}{2}}b)^{-2} \}^{\frac{1}{3}}.$$

$$99. \frac{a+b}{a^{\frac{1}{3}}-b^{\frac{1}{3}}} - \frac{a-b}{a^{\frac{1}{3}}+b^{\frac{1}{3}}}.$$

$$96. \left[\frac{\left(\frac{a^3b}{x^2y} \right)^{\frac{1}{2}}}{\left(\frac{a^3b^2}{xy^2} \right)^{\frac{1}{3}}} \right]^6 - \frac{(ax^{-1})^3}{b}.$$

$$100. \frac{\left[-\frac{a^{-1}+b^{-1}}{a^{-1}-b^{-1}} \times (a^2-b^2) \right]^{\frac{1}{2}}}{\frac{b+a}{ab}}.$$

QUADRATIC EQUATIONS

285. 1. What is the value of x in the equation $3x = 24$? What kind of an equation is it?

2. What powers of x are found in the equation $x^2 + 2x = 3$? Which is the higher power?

3. What is the value of x in the equation $x^2 = 9$? How many values has x ? How do they compare numerically?

4. Factor $x^2 - 5x + 6 = 0$, and so find the values of x . How many values has x ?

286. An integral equation that contains the *square* of the unknown number, but no higher power, is called a **Quadratic Equation**, or an equation of the *second degree*.

It is evident, therefore, that quadratic equations may be of two kinds—those which contain only the second power of the unknown number, and those which contain both the second and first powers.

$x^2 = 16$ and $ax^2 + bx = c$ are quadratic equations.

PURE QUADRATICS

287. An equation that contains only the second power of the unknown number is called a **Pure Quadratic**.

$ax^2 = b$ and $ax^2 - cx^2 = bc$ are pure quadratics.

Pure Quadratics are also called *Incomplete Quadratics*, because they lack the first power of the unknown number.

288. Since pure quadratics contain only the second power of the unknown number, they may be reduced to the general form $ax^2 = b$, in which a represents the coefficient of x^2 , and b the sum of the terms that do not involve x^2 .

289. PRINCIPLES. — 1. *If the square root of each member of a quadratic equation is extracted and the second member of the resulting equation is given the sign \pm , the resulting equation is equivalent to the given equation.*

2. *Every pure quadratic equation has two roots numerically equal, but having opposite signs.*

For the equation $A^2 = B^2$, or $A^2 - B^2 = 0$, or $(A - B)(A + B) = 0$, is equivalent to the two equations

$$A - B = 0 \text{ and } A + B = 0,$$

or

$$A = +B \text{ and } A = -B;$$

that is, to the two equations $A = \pm B$.

EXAMPLES

1. Given $10x^2 = 99 - x^2$, to find the value of x .

SOLUTION

$$10x^2 = 99 - x^2.$$

Transposing, etc.,

$$11x^2 = 99.$$

Dividing by 11,

$$x^2 = 9.$$

Extracting the square root of each member, Ax. 7,

$$x = \pm 3.$$

Strictly speaking, the last equation should be $\pm x = \pm 3$, which stands for the equations $+x = +3$, $+x = -3$, $-x = -3$, and $-x = +3$. But since the last two equations may be derived from the first two, the first two express *all* the values of x . For convenience, the two expressions, $x = +3$ and $x = -3$, are written $x = \pm 3$.

Consequently, in extracting the square roots of the members of an equation, it will be sufficient to write the ambiguous sign before the root of *one* member.

2. Find the roots of the equation $3x^2 = -15$.

SOLUTION

$$3x^2 = -15.$$

Dividing by 3,

$$x^2 = -5.$$

Extracting the square root,

$$x = \pm \sqrt{-5}.$$

Solve the following equations:

3. $7x^2 - 25 = 5x^2 + 73$.
4. $(x+4)^2 = 8x + 25$.
5. $(a-x)^2 = (3x+a)(x-a)$.
6. $ax^2 = (a-b)(a^2-b^2) - bx^2$.
7. $a^2x^2 + 2ax^2 = (a^2-1)^2 - x^2$.
8. $(x+2)^2 - 4(x+2) = 4$.
9. $\frac{x-8}{6} = \frac{6}{x+8}$.
10. $\frac{1}{1-x} + \frac{1}{1+x} = 2\frac{2}{3}$.
11. $\frac{x}{12} + \frac{x^2-15}{5x} = \frac{x}{5}$.
12. $\frac{x+3}{x-3} + \frac{x-3}{x+3} = 4$.
13. $\frac{x-2}{x+1} + \frac{x+2}{x-1} = -1$.
14. $\frac{x}{a+b} - \frac{a-b}{x} = 0$.
15. $\frac{x-3}{x-2} + \frac{x+3}{x+2} = 1\frac{1}{3}$.
16. $\frac{a}{x} + \frac{x}{a} = \frac{ab}{x}$.
17. $\sqrt{x^2+8} - \frac{6}{\sqrt{x^2+8}} = x$.
18. $x + \sqrt{x^2+m^2} = \frac{2m^2}{\sqrt{x^2+m^2}}$.
19. $\frac{x+a}{x+b} + \frac{x-a}{x-b} = \frac{a^2+b^2}{x^2-b^2}$.
20. $x + \sqrt{x^2+3} = \frac{6}{\sqrt{x^2+3}}$.
21. $\frac{24}{\sqrt{x^3+12}} - \sqrt{x^2+12} = x$.
22. $\frac{x+a}{x-a} + \frac{x-a}{x+a} = \frac{2a}{1-a}$.
23. $\frac{x+7}{x^2-7x} - \frac{x-7}{x^2+7x} = \frac{7}{x^2-73}$.
24. $x + \sqrt{x^2-a^2} = \frac{a^2}{\sqrt{x^2-a^2}}$.
25. $\sqrt{25-6x} + \sqrt{25+6x} = 8$.
26. $\frac{2x + \sqrt{4x^2-1}}{2x - \sqrt{4x^2-1}} = 4$.
27. $\frac{\sqrt{1+x}}{1+\sqrt{1+x}} + \frac{\sqrt{1-x}}{1+\sqrt{1-x}} = 0$.
28. $\frac{\sqrt{x^2+1} - \sqrt{x^2-1}}{\sqrt{x^2+1} + \sqrt{x^2-1}} = \frac{1}{2}$.
29. $\frac{1}{1+\sqrt{1-x}} - \frac{1}{\sqrt{1+x}+1} = \frac{1}{x}$.

$$30. \frac{2}{x + \sqrt{2 - x^2}} + \frac{2}{x - \sqrt{2 - x^2}} = x.$$

$$31. \frac{\sqrt{x + 2a} - \sqrt{x - 2a}}{\sqrt{x - 2a} + \sqrt{x + 2a}} = \frac{x}{2a}.$$

$$32. \sqrt{\frac{x-a}{x+a}} + \sqrt{\frac{x+a}{x-a}} = a^2.$$

PROBLEMS

1. If 25 is added to the square of a certain number, the sum is equal to the square of 13. What is the number?

2. What number is that whose square is equal to the difference of the squares of 25 and 20?

3. If a certain number is increased by 5 and also decreased by 5, the product of these results will be 75. What is the number?

4. Two numbers are as 3 to 4, and the sum of their squares is equal to the square of 15. What are the numbers?

5. Two numbers are as 4 to 3, and the sum of their squares is 400. What are the numbers?

6. A gentleman has two square rooms whose sides are as 2 to 3. He finds that it takes 9 square yards more than twice as much carpeting for the larger room as for the smaller. What is the length of a side of each room?

7. A man who owns a field 80 rods square sells one fourth of it. If the part he sells is also a square, how long is each of its sides?

8. A man had a rectangular field whose width was $\frac{2}{3}$ of its length. He built a fence across it so that one of the two parts thus formed was a square. If the square field contained 10 acres, what were the dimensions of the original field?

9. How many rods of fence will inclose a square garden whose area is $2\frac{1}{2}$ acres?

10. The sum of two numbers is 10, and their product is 21. What are the numbers?

SUGGESTION. — Represent the numbers by $5 + x$ and $5 - x$.

11. The sum of two numbers is 16, and their product is 55. What are the numbers?

12. The sum of two numbers is 26, and their product is 69. What are the numbers?

13. The sum of two numbers is 5, and their product is -14 . What are the numbers?

14. Factor $a^2 + 17a + 60$ by the method suggested in the preceding problems.

SUGGESTION. $+60$ is the product of the arithmetical terms, and $+17$ is their algebraic sum.

15. Separate $a^2 + 2a - 2$ into two factors.

16. Separate $x^2 - 2x - 1$ into two factors.

17. Divide 24 into two parts whose product is 143.

18. The length of a ten-acre field was 4 times its width. What were its dimensions?

19. The sum of the squares of two numbers is 394, and the difference of their squares is 56. What are the numbers?

20. A man has two square fields that together contain $51\frac{1}{4}$ acres. If the side of one is as much longer than 50 rods as that of the other is shorter than 50 rods, what are the dimensions of each field?

AFFECTED QUADRATICS

290. A quadratic equation that contains both the second and the first powers of one unknown number is called an **Affected Quadratic**.

$x^2 + 3x = 10$, $4x^2 - x + 1 = 0$, and $ax^2 + bx + c = 0$ are affected quadratics.

Affected Quadratics are also called *Complete Quadratics*.

291. Since affected quadratic equations contain both the second and first powers of the unknown number, they may always be reduced to the general form of $ax^2 + bx + c = 0$, in which a , b , and c may represent any numbers whatever.

The term c is sometimes called the *absolute term*.

292. To solve affected quadratics by factoring.

Reduce the equations to the form $ax^2 + bx + c = 0$, and solve by the methods of § 141.

Solve the following equations by factoring:

1. $x^2 - 5x + 6 = 0$.
2. $x^2 - 5x = 24$.
3. $x^2 - 1 = 3(x + 1)$.
4. $2x^2 - 7x + 3 = 0$.
5. $2x^2 - x - 3 = 0$.
6. $3x^2 - 2x - 8 = 0$.
7. $10x^2 - 27x + 5 = 0$.
8. $6(x^2 + 1) = 13x$.
9. $x^2 - (a - b)x = ab$.
10. $2x^2 - 3ax - 2a^2 = 0$.
11. $3(b^2 + x^2) = 10bx$.
12. $x^2 - 2ax + (a + 1)(a - 1) = 0$.
13. Solve the equation $x^2 + 100x + 2491 = 0$.

SOLUTION

Since, § 99, 100 is the sum of the arithmetical terms of the factors of $x^2 + 100x + 2491$, and their product is 2491, $50 + p$ and $50 - p$ may represent the two factors of 2491 whose sum is 100.

$$\begin{aligned} \text{Then,} & (50 + p)(50 - p) = 2491. \\ \text{Expanding,} & 2500 - p^2 = 2491. \\ \text{Transposing, etc.,} & p^2 = 9. \\ \therefore & p = \pm 3. \end{aligned}$$

$$50 + p = 53, \quad 50 - p = 47.$$

$$\begin{aligned} \text{Therefore, § 130,} & x^2 + 100x + 2491 = (x + 53)(x + 47) = 0, \\ \text{and} & x = -53 \text{ or } -47. \end{aligned}$$

Since $p = -3$ gives no new values of $50 + p$ and $50 - p$, the negative value of p may be disregarded.

14. Solve the equation $x^2 + 3x - 208 = 0$.

SOLUTION

$$\begin{aligned} \text{Let} & x^2 + 3x - 208 = 0. \\ \text{Expanding,} & (\frac{x}{2} + p)(\frac{x}{2} - p) = -208. \\ \text{Solving,} & \frac{x}{4} - p^2 = -208. \\ & p = \pm \frac{29}{2}. \\ & \frac{x}{2} + p = 16, \quad \frac{x}{2} - p = -13. \end{aligned}$$

$$\begin{aligned} \text{Factoring the given equation,} & (x + 16)(x - 13) = 0. \\ \therefore & x = -16 \text{ or } +13. \end{aligned}$$

Solve the following equations:

15. $x^2 + 10x + 21 = 0$.

28. $x^2 + x - 756 = 0$.

16. $x^2 + 12x - 28 = 0$.

29. $x^2 - x - 506 = 0$.

17. $x^2 - 20x + 51 = 0$.

30. $x^2 + 2x - 168 = 0$.

18. $x^2 + 60x + 891 = 0$.

31. $x^2 + 6x - 135 = 0$.

19. $x^2 - 44x + 403 = 0$.

32. $x^2 + 3x - 154 = 0$.

20. $x^2 + 20x - 629 = 0$.

33. $x^2 + 5x + 2 = 0$.

21. $x^2 - 30x - 2275 = 0$.

34. $x^2 + x - 10 = 0$.

22. $x^2 + 24x + 119 = 0$.

35. $x^2 - x - 1 = 0$.

23. $x^2 + 2x - 323 = 0$.

36. $x^2 - 2x - 4 = 0$.

24. $x^2 - 6x - 475 = 0$.

37. $x^2 - 3x - 9 = 0$.

25. $x^2 + 8x - 768 = 0$.

38. $x^2 + 4x + 8 = 0$.

26. $x^2 + 3x - 418 = 0$.

39. $x^2 + 6x + 14 = 0$.

27. $x^2 + 5x - 546 = 0$.

40. $x^2 + 8x = -25$.

293. First method of completing the square.

1. What is the square of $x + 3$? of $x + 5$? of $x + 10$?

2. If $x^2 + 20x$ are the first two terms of the square of a binomial, what is the first term of the binomial?

3. Since $20x$ is twice the product of the two terms of the binomial, and the first term of the binomial is x , how may the second term of the binomial be found?

4. Since the second term of the binomial is 10, what must be added to $x^2 + 20x$ to *complete* the square of the binomial?

5. What term must be added to $x^2 + 6x$ to complete the square of some binomial? How is the term found?

6. What term must be added to $x^2 + 8x$ to complete the square? to $x^2 - 10x$? to $x^2 - 14x$?

7. What must be added to both members of the equation $x^2 - 12x = 13$ to make the first member a perfect square?

EXAMPLES

1. Solve the equation $x^2 - 6x = 40$.

PROCESS	EXPLANATION.
$x^2 - 6x = 40$	Completing the square in the first member by adding to each member the square of half the coefficient of x , $x^2 - 6x + 9 = 49$.
$x^2 - 6x + 9 = 49$	Extracting the square root of each member,
$x - 3 = \pm 7$	$x - 3 = \pm 7$.
$x = +7 + 3 = 10$	Using first the upper sign of ± 7 , the simple equation $x - 3 = +7$ gives $x = 10$. Next using the lower sign of ± 7 , the simple equation $x - 3 = -7$ gives
$x = -7 + 3 = -4$	$x = -4$.

Since each of the values 10 and -4 satisfies the given equation when substituted for x , $x = 10$ or -4 .

2. Solve the equation $x^2 - 5x = 14$.

SOLUTION

	$x^2 - 5x = 14$.
Completing the square,	$x^2 - 5x + (\frac{5}{2})^2 = 14 + \frac{25}{4} = \frac{81}{4}$.
Extracting the square root,	$x - \frac{5}{2} = \pm \frac{9}{2}$.
Taking the upper sign,	$x = \frac{5}{2} + \frac{9}{2} = 7$.
Taking the lower sign,	$x = \frac{5}{2} - \frac{9}{2} = -2$.

3. Solve the equation $10x^2 + 19x = 15$.

SOLUTION

	$10x^2 + 19x = 15$.
Dividing by coefficient of x^2 ,	$x^2 + \frac{19}{10}x = \frac{3}{2}$.
Completing the square,	$x^2 + \frac{19}{10}x + (\frac{19}{20})^2 = \frac{3}{2} + \frac{361}{400} = \frac{2401}{400}$.
Extracting the square root,	$x + \frac{19}{20} = \pm \frac{49}{20}$.
Taking the upper sign,	$x = -\frac{19}{20} + \frac{49}{20} = \frac{3}{5}$.
Taking the lower sign,	$x = -\frac{19}{20} - \frac{49}{20} = -\frac{3}{2}$.

RULE. — Transpose so that the terms containing x^2 and x are in one member of the equation and the known terms in the other, and make the coefficient of x^2 unity by dividing both members of the equation by the coefficient of x^2 .

Add to each member of the equation the square of half the coefficient of x , and extract the square root of each member.

Solve the two simple equations thus obtained.

Find the values of x in the following equations:

- | | |
|------------------------|---------------------------|
| 4. $x^2 - 2x = 143.$ | 15. $x^2 + 3x = 10.$ |
| 5. $x^2 + 2x = 168.$ | 16. $x^2 - 3x = 180.$ |
| 6. $x^2 - 4x = 117.$ | 17. $x^2 + 15x = 54.$ |
| 7. $x^2 - 6x = 160.$ | 18. $x^2 - x = 930.$ |
| 8. $x^2 - 8x = 180.$ | 19. $x^2 + 13x = 140.$ |
| 9. $x^2 + 2x = 120.$ | 20. $x^2 - 11x + 28 = 0.$ |
| 10. $x^2 + 6x = 187.$ | 21. $5x^2 - 3x - 2 = 0.$ |
| 11. $x^2 - 12x = 189.$ | 22. $6x^2 - 5x - 6 = 0.$ |
| 12. $x^2 + 10x = 171.$ | 23. $2x^2 + 9x = 35.$ |
| 13. $x^2 - 22x = 48.$ | 24. $3x^2 - 7x = 10.$ |
| 14. $x^2 + 30x = 31.$ | 25. $4x^2 - 19x = 5.$ |

$$26. \frac{1}{x+1} + \frac{3}{x-1} = \frac{10}{3}.$$

$$27. \frac{x^2}{x-2} - \frac{3x-5}{2} = \frac{x+2}{5}.$$

$$28. \frac{1}{x+2} + \frac{x-2}{x} = \frac{x-7}{2x}.$$

$$29. x^2 + (m+n)(m-n) = 2mx.$$

294. Other methods of completing the square.

By the previous method, when x^2 had a coefficient, the equation was divided by that coefficient so that the *term containing x^2 might always be a perfect square*. The same result may be secured in other ways.

Thus, if the term containing x^2 is $3x^2$, it may be made a perfect square by multiplying by 3; if $8x^2$, by multiplying by 2; if ax^2 , by multiplying by a .

In the completed square $a^2x^2 + 2abx + b^2$, it is evident that the third term, b^2 , is the square of the quotient obtained by dividing the second term by twice the square root of the first.

EXAMPLES

1. Solve the equation $5x^2 + 12x = 9$.

SOLUTION

$$5x^2 + 12x = 9.$$

Multiplying by 5, $25x^2 + 60x = 45$.

Completing the square, $25x^2 + 60x + 36 = 81$.

Extracting the square root, $5x + 6 = \pm 9$.

$$5x = -6 \pm 9.$$

$$\therefore x = \frac{3}{5} \text{ or } -3.$$

EXPLANATION. — Since the coefficient of x^2 is not a perfect square, it may be made a perfect square by multiplying the members of the equation by 5.

Since, if to $25x^2 + 60x$ there were added such a term as would make the trinomial a perfect square, $60x$ would be equal to twice the product of the square root of $25x^2$ and the square root of this third term, the square root of the third term is obtained by dividing $60x$ by $2\sqrt{25x^2}$; that is, by $10x$. $60x \div 10x = 6$, and 6^2 , or 36 , added to both members completes the square of $(5x + 6)$.

2. Solve the equation $8x^2 - 10x = 3$.

SOLUTION

$$8x^2 - 10x = 3.$$

Multiplying by 2, $16x^2 - 20x = 6$.

Completing the square, $16x^2 - 20x + (\frac{5}{4})^2 = 6 + \frac{25}{4} = \frac{49}{4}$.

Extracting the square root, $4x - \frac{5}{2} = \pm \frac{7}{2}$.

$$4x = \frac{5}{2} \pm \frac{7}{2} = 6 \text{ or } -1.$$

$$\therefore x = \frac{3}{2} \text{ or } -\frac{1}{4}.$$

GENERAL RULE. — Transpose so that the terms containing x^2 and x are in one member of the equation and the known terms in the other.

If the term containing the second power of the unknown number is not a perfect square, make it such by multiplying or dividing the members of the equation by some number.

Add to each member of the equation the square of the quotient obtained by dividing the term containing the first power of the unknown number by twice the square root of the term containing the second power.

Extract the square root of each member, and solve the two simple equations.

Solve the following equations :

3. $2x^2 - 5x = 42.$

8. $3x^2 + 4x = 95.$

4. $6x^2 - 5x + 1 = 0.$

9. $7x^2 + 2x = 32.$

5. $4x^2 - 12x = 27.$

10. $8x^2 - 18x = 5.$

6. $8x^2 + 20x = 48.$

11. $6x^2 + 5x = 4.$

7. $18x^2 + 6x = 4.$

12. $5x^2 + 6x = 8.$

13. Solve the equation $ax^2 + bx + c = 0.$

SOLUTION

$$ax^2 + bx + c = 0. \quad (1)$$

Transposing c , $ax^2 + bx = -c. \quad (2)$

Multiplying by a , $a^2x^2 + abx = -ac. \quad (3)$

Completing the square, $a^2x^2 + abx + \frac{b^2}{4} = \frac{b^2}{4} - ac. \quad (4)$

Multiplying by 4, $4a^2x^2 + 4abx + b^2 = b^2 - 4ac. \quad (5)$

Extracting the square root, $2ax + b = \pm \sqrt{b^2 - 4ac}. \quad (6)$

$$\therefore x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}. \quad (7)$$

It is evident that (5) can be obtained by multiplying (2) by $4a$ and adding b^2 to both members. Hence, when a quadratic has the general form of (1), if the absolute term is transposed to the second member, as in (2), the square may be completed and fractions avoided by *multiplying by 4 times the coefficient of x^2 and adding to each member the square of the coefficient of x in the given equation.*

This is called the *Hindoo* method of completing the square.

Solve the following equations by the Hindoo method :

14. $2x^2 + 3x = 27.$

21. $4x^2 - x - 3 = 0.$

15. $2x^2 + 5x = 7.$

22. $5x^2 - 2x - 16 = 0.$

16. $2x^2 + 7x = -6.$

23. $3x^2 + 7x - 110 = 0.$

17. $3x^2 - 5x = 2.$

24. $2x^2 - 5x - 150 = 0.$

18. $4x^2 - 15x = 4.$

25. $3x^2 + x - 200 = 0.$

19. $5x^2 - 7x = -2.$

26. $15x^2 - 7x - 2 = 0.$

20. $6x^2 + 5x = -1.$

27. $7x^2 - 20x - 32 = 0.$

295. To solve quadratics by a formula.

Since every quadratic can be reduced to the general form $ax^2 + bx + c = 0$, in which a , b , and c represent any numbers whatever, and since the roots of this equation are

$$\text{Ex. 13, § 294,} \quad x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a},$$

the values of the unknown number in any affected quadratic equation may be found by substituting the coefficient of x^2 for a the coefficient of x for b , and the absolute term for c .

EXAMPLES

1. Solve the equation $6x^2 - x - 15 = 0$.

SOLUTION. — Since $a = 6$, $b = -1$, and $c = -15$, by the above formula,

$$\begin{aligned} x &= \frac{1 \pm \sqrt{(-1)^2 - 4 \times 6(-15)}}{2 \times 6} \\ &= \frac{1 \pm 19}{12} = \frac{5}{3} \text{ or } -\frac{3}{2}. \end{aligned}$$

Solve by the above formula:

2. $2x^2 + 5x + 2 = 0$.

10. $2x^2 + 3x - 1 = 0$.

3. $3x^2 + 11x + 6 = 0$.

11. $3x^2 + 2x - 4 = 0$.

4. $6x^2 - 7x + 2 = 0$.

12. $x^2 - 5x = -3$.

5. $4x^2 + 4x - 15 = 0$.

13. $3x^2 - 6x = -2$.

6. $2x^2 + 3x - 9 = 0$.

14. $4x^2 - 3x - 2 = 0$.

7. $2x^2 + 3x + 1 = 0$.

15. $x^2 - 6x + 10 = 0$.

8. $3x^2 - 13x = 10$.

16. $x^2 + 4x + 12 = 0$.

9. $7x^2 + 9x = 10$.

17. $x^2 - 8x = -20$.

Solve by any method:

18. $x^2 - 6x + 5 = 0$.

22. $x^2 - 12x = 64$.

19. $x^2 - 8x + 7 = 0$.

23. $18x^2 + 6x = 4$.

20. $2x^2 - 5x = 42$.

24. $x^2 - x - 72 = 0$.

21. $7x^2 + 2x = 32$.

25. $4x^2 - 12x = 27$.

26. $x^2 = 3x + 10.$

27. $x^2 - 30 = 13x.$

28. $x^2 - 12x = 28.$

29. $x^2 - 50x = 159.$

30. $x^2 + 8x = 84.$

31. $x + \frac{1}{x} - \frac{5}{2} = 0.$

32. $\frac{x^2}{9} + \frac{x}{3} = \frac{35}{4}.$

33. $\frac{x}{9(x-1)} = \frac{x-2}{6}.$

34. $\frac{4}{x^2 - 2x + 1} = \frac{1}{4}.$

35. $\frac{x^2}{4} - \frac{2x}{3} = 28.$

36. $\frac{9}{2x+1} + \frac{3}{x-3} = 4.$

37. $\frac{3x-1}{x+2} = \frac{x+1}{x-2}.$

38. $\frac{1+x}{x-3} - \frac{x-1}{x-2} = \frac{4}{5}.$

39. $\frac{x}{x-5} - \frac{x-5}{x} = \frac{3}{2}.$

40. $\frac{x+7}{x+5} + \frac{x+12}{x+6} = 7.$

41. $\frac{x+4}{x-2} + 3 = \frac{x+3}{x-3}.$

42. $\frac{4x}{x-1} - \frac{x+3}{x} = 4.$

43. $\frac{x}{x+2} + \frac{1}{2} = \frac{x+2}{2x}.$

44. $\frac{5x}{x+7} + \frac{x+6}{x+3} = 3.$

45. $\frac{x+2}{x-7} - \frac{x+5}{x-5} = 1.$

46. $\frac{x-3}{x+4} + \frac{x+2}{x-2} = \frac{23}{10}.$

47. $\frac{2x+1}{1-2x} - \frac{5}{7} = \frac{x-8}{2}.$

LITERAL EQUATIONS

296. 1. Solve the equation $x^2 - \frac{a}{b}x - \frac{b}{a}x + 1 = 0.$

SOLUTION

$$x^2 - \frac{a}{b}x - \frac{b}{a}x + 1 = 0.$$

Factoring,
$$\left(x - \frac{a}{b}\right)\left(x - \frac{b}{a}\right) = 0.$$

Therefore,
$$x = \frac{a}{b} \text{ or } \frac{b}{a}$$

Solve by any of the preceding methods:

2. $x^2 - ax = ab - bx.$

6. $5x - 2ax = x^2 - 10a.$

3. $x^2 + ax = ac + cx.$

7. $x^2 + 3bx = 5cx + 15bc.$

4. $x^2 = (m - n)x + mn.$

8. $2abx - x^2 = 14ab - 7x.$

5. $x^2 - 3bx = 2ax - 6ab.$

9. $6x^2 + 3ax = 2bx + ab.$

10. $acx^2 - bcx - bd + adx = 0.$

11. $x^2 + 4mx + 3nx + 12mn = 0.$

12. $x^2 - 2ax = a^2.$

26. $\frac{x}{a} + \frac{a}{x} = \frac{5}{2}.$

13. $x^2 + 4bx = b^2.$

27. $\frac{3c^2x^2}{4} + \frac{2cx}{3} = \frac{5}{3}.$

14. $x^2 = 4ax - 2a^2.$

28. $\frac{2x}{x-a} - \frac{3a}{x+a} = 3.$

15. $x^2 - ax + a^2 = 0.$

29. $\frac{a}{x-a} - 2 = \frac{2a}{x}.$

16. $x^2 = bx + b^2.$

17. $x^2 + px + q = 0.$

30. $\frac{1}{ax+4} = 1 - \frac{ax-4}{16}.$

18. $x^2 - 2x + a = 0.$

19. $4ax - x^2 = 3a^2.$

31. $x^2 + \frac{a}{b}x = \frac{a+b}{b}.$

20. $x^2 - a = 1 + ax.$

21. $x^2 - b^2 = a^2 - bx.$

32. $x^2 + 2 = \frac{(2a^2 + 1)x}{a}.$

22. $21b^2 - 4bx = x^2.$

23. $5ax + 6a^2 = 6x^2.$

24. $x^2 - 1 = 4ax - a^2.$

33. $x^2 - \frac{2x}{ab} = \frac{4(ab-1)}{ab}.$

25. $x^2 + b^2 = 4(a^2 + bx).$

34. $x^2 - 2(a-b)x = 4ab.$

35. $x^2 + 2(a+8)x = -32a.$

36. $x^2 + x + bx + b = ax + a.$

37. $2ax - a + 2bx - b = 2x^2 - x.$

38. $x^2 + 4(a-1)x = 8a - 4a^2.$

39. $a(x-2a+b) + a(x+a-b) = x^2 - (a-b)^2.$

$$40. \frac{2a+x}{2a-x} + \frac{a-2x}{a+2x} = \frac{8}{3}.$$

$$41. \frac{1}{a-x} - \frac{1}{a+x} = \frac{3+x^2}{a^2-x^2}.$$

$$42. \frac{1}{x+a} + \frac{1}{x+b} = \frac{1}{a-b}.$$

$$43. \frac{x^2+1}{x} - \frac{a+b}{c} = \frac{c}{a+b}.$$

$$44. \frac{2x-a}{b} + 3 = \frac{4a}{2x-b}.$$

$$45. \frac{bx}{a-x} + b = \frac{a(x+2b)}{a+b}.$$

$$46. \sqrt{a+x} - \sqrt{a-x} = \sqrt{2x}.$$

$$47. \sqrt{x-a} + \sqrt{b-x} = \sqrt{b-a}.$$

$$48. \sqrt{x^2-b^2} = \sqrt{x+b} \sqrt{a+b}.$$

$$49. \sqrt{a-x} + \sqrt{b-x} = \sqrt{a+b-2x}.$$

$$50. \text{ Solve and verify } \sqrt{x+1} + \sqrt{x-2} - \sqrt{2x-5} = 0.$$

SOLUTION

$$\sqrt{x+1} + \sqrt{x-2} - \sqrt{2x-5} = 0.$$

Transposing,

$$\sqrt{x+1} + \sqrt{x-2} = \sqrt{2x-5}.$$

Squaring,

$$x+1 + 2\sqrt{x^2-x-2} + x-2 = 2x-5.$$

Simplifying,

$$\sqrt{x^2-x-2} = -2.$$

Squaring,

$$x^2-x-2 = 4.$$

Solving,

$$x = -2 \text{ or } 3.$$

VERIFICATION. — Substituting -2 for x in the given equation,

$$\sqrt{-1} + \sqrt{-4} - \sqrt{-9} = 0;$$

that is,

$$\sqrt{-1} + 2\sqrt{-1} - 3\sqrt{-1} = 0.$$

Therefore, -2 is a root of the given equation.Substituting 3 for x in the given equation,

$$\sqrt{4} + \sqrt{1} - \sqrt{1} = 0,$$

which is not true according to the convention adopted in the discussion in § 284. Hence, 3 is not to be regarded as a root of the given equation.

NOTE. — In the previous verification, when only positive square roots are taken, the second value obtained for x does not satisfy the given equation, yet this indicates no error in the process of rationalizing, for the equation can be verified by admitting negative square roots. But, as explained in § 283, it is more convenient to regard 3 as the root of the equation $\sqrt{x+1} - \sqrt{x-2} - \sqrt{2x-5} = 0$, which has for its first member one of the rationalizing factors of the given equation.

Solve and verify:

51. $8\sqrt{x} - 8x = \frac{3}{2}$.

53. $x - 1 + \sqrt{x+5} = 0$.

52. $3x + \sqrt{x} = 5\sqrt{4x}$.

54. $x - 5 - \sqrt{x-3} = 0$.

55. $\sqrt{4x+17} + \sqrt{x+1} - 4 = 0$.

56. $\sqrt{x-1} + \sqrt{2x-1} - \sqrt{5x} = 0$.

57. $\sqrt{2x-7} - \sqrt{2x} + \sqrt{x-7} = 0$.

58. $\sqrt{x+3} + \sqrt{4x+1} - \sqrt{10x+4} = 0$.

59. $\sqrt{6+x} + \sqrt{x} - \sqrt{10-4x} = 0$.

60. $\sqrt{4x-3} - \sqrt{2x+2} = \sqrt{x-6}$.

61. $\sqrt{2x+3} - \sqrt{x+1} = \sqrt{5x-14}$.

62. $\sqrt{3x-5} + \sqrt{x-9} = \sqrt{4x-4}$.

63. $\sqrt{x+a^2} - \sqrt{x-2a^2} = \sqrt{2x-5a^2}$.

PROBLEMS

297. 1. The sum of two numbers is 8, and their product is 15. Find the numbers.

SOLUTION

Let

$x =$ one number.

Then,

$8 - x =$ the other.

Since their product is 15,

$$8x - x^2 = 15.$$

Solving,

$$x = 3 \text{ or } 5,$$

and

$$8 - x = 5 \text{ or } 3.$$

Therefore, the numbers are 3 and 5.

2. A party hired a coach for \$12. In consequence of the failure of three of them to pay, each of the others had to pay 20 cents more. How many persons were in the party?

SOLUTION

Let x = the number of persons.
 Then, $x - 3$ = the number who paid,
 $\frac{12}{x - 3}$ = the number of dollars each paid,
 and $\frac{12}{x}$ = the number of dollars each should have paid.
 Therefore, $\frac{12}{x - 3} - \frac{1}{5} = \frac{12}{x}$.
 Solving, $x = 15$ or -12 .

The second value of x is evidently inadmissible. Hence, the number of persons in the party was 15.

3. A cistern can be filled by two pipes in 24 minutes. If it takes the smaller pipe 20 minutes longer to fill the cistern than the larger pipe, in what time can the cistern be filled by each pipe?

SOLUTION

Let x = the number of minutes required by the larger pipe.
 Then, $x + 20$ = the number of minutes required by the smaller.
 Since $\frac{1}{x}$ = the part which the larger pipe fills in one minute,
 and $\frac{1}{x + 20}$ = the part which the smaller pipe fills in one minute,
 and $\frac{1}{24}$ = the part which both pipes fill in one minute,
 $\frac{1}{x} + \frac{1}{x + 20} = \frac{1}{24}$.
 Solving, $x = 40$ or -12 .

Hence, the larger pipe can fill the cistern in 40 minutes, and the smaller in 60 minutes.

4. Divide 20 into two parts whose product is 96.

5. Divide 14 into two parts whose product is 45.

6. A man purchased a flock of sheep for \$75. If he had paid the same sum for a flock containing 3 more, they would have cost \$1.25 less per head. How many did he purchase?

7. A rectangular garden is 12 rods longer than it is wide, and contains 1 acre. What are its dimensions?

8. If each side of a square field were lengthened 4 rods, the area would be increased 136 square rods more than $\frac{1}{4}$ of it. What are the dimensions of the field?

9. A rectangular lot is 8 rods longer than it is wide. What are its dimensions, if it contains $1\frac{1}{2}$ acres?

10. In a column of 600 soldiers each file contained 3 men more than 9 times as many men as each rank. How broad and how deep was the column?

11. A party had a dinner that cost \$60. If there had been 5 persons more, the share of each would have been \$1 less. How many persons were there in the party?

12. A man worked a certain number of days for \$30. If he had received \$1 per day less than he did, he would have been obliged to work 5 days longer to earn the same sum. How many days did he work?

13. Find two consecutive numbers the sum of whose squares is 61.

14. Find two consecutive numbers the sum of whose reciprocals is $\frac{2}{25}$.

15. A picture that was 8 inches by 12 inches was placed in a frame of uniform width. If the area of the frame was equal to that of the picture, what was the width of the frame?

16. A merchant purchased a quantity of flour for \$100, and retailed it at a gain of \$1 per barrel. After he had sold \$100 worth of it, he had 5 barrels of it left. How many barrels did he buy, and at what price?

17. A merchant sold a hunting coat for \$11, and gained a per cent equal to the number of dollars the coat cost him. What was his per cent of gain?

18. A railway train traveled 5 miles an hour slower than usual and was one hour late in making a run of 280 miles. How many miles per hour did it travel?

19. A rectangular park 56 rods long and 16 rods wide was surrounded by a street of uniform width containing 4 acres. What was the width of the street?

20. A boatman rowed 8 miles up a stream and back in 3 hours. If the velocity of the current was 2 miles an hour, what was his rate of rowing in still water?

21. A man who owned a lot 56 rods long and 28 rods wide constructed a road around it, thereby decreasing the area of the lot 2 acres. What was the width of the road?

22. A man bought two lots of cloth and paid 96 shillings for each. There were 20 yards in all, and the number of shillings per yard paid for each was the same as the number of yards of the other. How many yards of each did he buy?

23. Find the price of eggs, when 2 less for 30 cents raises the price 2 cents per dozen.

24. A and B started at the same time and traveled toward a place 75 miles distant. A traveled one mile an hour faster than B and reached the place $2\frac{1}{2}$ hours before B. At what rate did each travel?

25. A person drew a quantity of wine from a cask filled with 81 gallons of pure wine, and replaced it with water. He then drew from the mixture as many gallons as he drew before of pure wine, when it was found that the cask contained only 64 gallons of pure wine. How many gallons did he draw each time?

26. The circumference of the fore wheel of a coach is 5 feet less than that of the hind wheel. If the fore wheel makes 150 more revolutions than the hind wheel in going a mile, what is the circumference of each wheel?

27. Two pipes running together can fill a cistern in $2\frac{1}{2}$ hours. The larger pipe can fill the cistern in 2 hours less time than the smaller. How many hours will it take each pipe alone to fill the cistern?

28. It took a number of men as many days to dig a ditch as there were men. If there had been 6 more men, they would have done the work in 8 days. How many men were there?

EQUATIONS IN THE QUADRATIC FORM

298. An equation that contains but two powers of an unknown number or expression, the exponent of one power being twice that of the other, is in the **Quadratic Form**.

Equations in the quadratic form can be reduced to the general form $ax^{2n} + bx^n + c = 0$, in which n represents any number.

EXAMPLES

1. Given $x^4 + 6x^2 - 40 = 0$, to find the values of x .

SOLUTION

$$x^4 + 6x^2 - 40 = 0.$$

Factoring, $(x^2 - 4)(x^2 + 10) = 0.$

$$\therefore x^2 - 4 = 0 \text{ or } x^2 + 10 = 0,$$

and

$$x = \pm 2 \text{ or } \pm \sqrt{-10}.$$

2. Given $x^{\frac{1}{2}} - x^{\frac{3}{4}} = 6$, to find the values of x .

FIRST SOLUTION

$$x^{\frac{1}{2}} - x^{\frac{3}{4}} = 6.$$

Completing the square, $x^{\frac{1}{2}} - x^{\frac{3}{4}} + (\frac{1}{4})^2 = 2\frac{1}{4}.$

Extracting the square root, $x^{\frac{1}{2}} - \frac{1}{2} = \pm \frac{3}{2}.$

$$\therefore x^{\frac{1}{2}} = 3 \text{ or } -2.$$

Raising to the fourth power, $x = 81 \text{ or } 16.$

SECOND SOLUTION

$$x^{\frac{1}{2}} - x^{\frac{3}{4}} = 6.$$

Let $x^{\frac{1}{4}} = p$, then, $x^{\frac{1}{2}} = p^2$, and $p^2 - p = 6.$

$$\therefore p^2 - p - 6 = 0.$$

Factoring, $(p - 3)(p + 2) = 0.$

$$\therefore p = 3 \text{ or } -2;$$

that is,

$$x^{\frac{1}{4}} = 3 \text{ or } -2.$$

Whence,

$$x = 81 \text{ or } 16.$$

3. Solve the equation $x - 4x^{\frac{1}{2}} + 3x^{\frac{1}{2}} = 0$.

SOLUTION

Let $x^{\frac{1}{2}} = p$, then, $x^{\frac{1}{2}} = p^2$, and $x = p^3$.

Then, $p^3 - 4p^2 + 3p = 0$.

Factoring, $p(p^2 - 4p + 3) = 0$.

Whence, $p = 0$,

or $p^2 - 4p + 3 = 0$.

Factoring, $(p - 1)(p - 3) = 0$.

Whence, $p = 1$ or $p = 3$.

That is, $x^{\frac{1}{2}} = 0, 1, \text{ or } 3$.

$\therefore x = 0, 1, \text{ or } 27$.

4. Given $x^2 - 7x + \sqrt{x^2 - 7x + 18} = 24$, to find the value of x .

SOLUTION

$$x^2 - 7x + \sqrt{x^2 - 7x + 18} = 24. \quad (1)$$

$$\text{Adding 18, } x^2 - 7x + 18 + \sqrt{x^2 - 7x + 18} = 42. \quad (2)$$

$$\text{Put } p \text{ for } (x^2 - 7x + 18)^{\frac{1}{2}} \text{ and } p^2 \text{ for } (x^2 - 7x + 18). \quad (3)$$

$$\text{Then, } p^2 + p - 42 = 0. \quad (4)$$

$$\text{Solving, } p = 6 \text{ or } -7. \quad (5)$$

$$\text{That is, } \sqrt{x^2 - 7x + 18} = 6, \quad (6)$$

$$\text{or } \sqrt{x^2 - 7x + 18} = -7. \quad (7)$$

$$\text{Squaring (6), } x^2 - 7x + 18 = 36.$$

$$\text{Solving, } x = 9 \text{ or } -2.$$

$$\text{Squaring (7), } x^2 - 7x + 18 = 49.$$

$$\text{Solving, } x = \frac{7}{2} \pm \frac{1}{2}\sqrt{173}.$$

$$\text{Hence, } x = 9, -2, \text{ or } \frac{7}{2} \pm \frac{1}{2}\sqrt{173}.$$

5. Solve the equation $x^6 - 9x^3 + 8 = 0$.

SOLUTION

$$x^6 - 9x^3 + 8 = 0. \quad (1)$$

$$\text{Factoring, } (x^3 - 1)(x^3 - 8) = 0. \quad (2)$$

$$\text{Therefore, } x^3 - 1 = 0, \quad (3)$$

$$\text{or } x^3 - 8 = 0. \quad (4)$$

If the values of x are found by transposing the known terms in (3) and (4) and extracting the cube root of each member, only *one* value of x will be obtained from each equation. But if the equations are factored, three values of x are obtained.

$$\text{Factoring,} \quad (x-1)(x^2+x+1)=0, \quad (5)$$

$$\text{and} \quad (x-2)(x^2+2x+4)=0. \quad (6)$$

Writing each factor equal to zero, and solving:

$$\text{From Eq. (5), } x=1, -\frac{1}{2}+\frac{1}{2}\sqrt{-3}, -\frac{1}{2}-\frac{1}{2}\sqrt{-3}. \quad (7)$$

$$\text{From Eq. (6), } x=2, -1+\sqrt{-3}, -1-\sqrt{-3}. \quad (8)$$

Since the values of x in (7) are obtained by factoring $x^3-1=0$, they may be regarded as the three cube roots of the number 1. Also, the values of x in (8) may be regarded as the three cube roots of the number 8 (§ 225).

$$6. \text{ Solve the equation } x^4+4x^3-8x+3=0.$$

SOLUTION

Extracting the square root of the first member as far as possible,

$$\begin{array}{r} x^4+4x^3-8x+3 \quad | \quad x^2+2x-2 \\ \hline x^4 \\ \hline 2x^2+2x \quad | \quad 4x^3 \\ \hline 4x^3+4x^2 \\ \hline 2x^2+4x-2 \quad | \quad -4x^2-8x+3 \\ \hline -4x^2-8x+4 \\ \hline -1 \end{array}$$

Since the first member lacks 1 of being a perfect square, the square may be completed by adding 1 to each member, giving the following equation:

$$x^4+4x^3-8x+4=1.$$

Extracting the square root, $x^2+2x-2=\pm 1$.

$$\therefore x^2+2x-3=0, \text{ and } x^2+2x-1=0.$$

$$\text{Solving,} \quad x=1, -3, -1\pm\sqrt{2}.$$

Solve the following equations:

$$7. \quad x^4-13x^2+36=0.$$

$$11. \quad 5x^4+6x^2-11=0.$$

$$8. \quad x^4-25x^2+144=0.$$

$$12. \quad 2x^4-8x^2-90=0.$$

$$9. \quad x^4-18x^2+32=0.$$

$$13. \quad x^{\frac{1}{2}}-5x^{\frac{1}{4}}+6=0.$$

$$10. \quad 3x^4+5x^2-8=0.$$

$$14. \quad x^{\frac{1}{2}}+3x^{\frac{1}{4}}-28=0.$$

15. $x^{\frac{1}{2}} - 3x^{\frac{1}{2}} = -2.$

16. $x^{\frac{2}{3}} - x^{\frac{1}{3}} = 6.$

17. $x + 2\sqrt{x} = 3.$

18. $x^{\frac{2}{3}} - 2x^{\frac{1}{3}} = 3.$

19. $x^3 + 8x^{\frac{3}{2}} - 9 = 0.$

20. $x^{\frac{3}{2}} + x^{\frac{1}{2}} - 2 = 0.$

21. $\sqrt[4]{x} + 3\sqrt{x} = 30.$

22. $ax^{2a} + bx^a + c = 0.$

23. $x^{\frac{3}{2}} - 4x - 5x^{\frac{1}{2}} = 0.$

24. $x^{\frac{7}{2}} - x^2 - 12x^{\frac{1}{2}} = 0.$

25. $x - 3x^{\frac{3}{2}} + 2x^{\frac{1}{2}} = 0.$

26. $5x = x\sqrt{x} + 6\sqrt{x}.$

27. $3x = x^{\frac{2}{3}}\sqrt{x} + 2\sqrt[3]{x^2}.$

28. $x^{-\frac{1}{2}} - 3 - 4x^{\frac{1}{2}} = 0.$

29. $x^{-\frac{1}{2}} - 6x^{\frac{1}{2}} = 1.$

30. $x^{-3} + x^2 = 2x^{\frac{1}{2}}.$

31. $x + 2x^{\frac{3}{2}} = 3x^{\frac{1}{2}}.$

32. $2x + \sqrt{x} = 15x\sqrt{x}.$

33. $\sqrt{x} + 5 + 6x^{-\frac{1}{2}} = 0.$

34. $x^4 = 8x + 7x^2\sqrt{x}.$

35. $(x-3)^2 + 2(x-3) = 3.$

36. $(x^2+1)^2 + 4(x^2+1) = 45.$

37. $(x^2-4)^2 - 3(x^2-4) = 10.$

38. $(x^2-2x)^2 - 2(x^2-2x) = 3.$

39. $(x^2-x)^2 - (x^2-x) - 132 = 0.$

40. $x - 5 + 2\sqrt{x-5} = 8.$

41. $x^2 - 3x + 6 + 2\sqrt{x^2 - 3x + 6} = 24.$

42. $x^2 - 5x + 2\sqrt{x^2 - 5x - 2} = 10.$

43. $x^2 - x - \sqrt{x^2 - x + 4} - 8 = 0.$

44. $x^2 - 5x + 5\sqrt{x^2 - 5x + 1} = 49.$

45. $x + 10 = 2\sqrt{x+10} + 5.$

49. $x + 2\sqrt{x+3} = 21.$

46. $x - 3 = 21 - 4\sqrt{x-3}.$

50. $2x - 3\sqrt{2x+5} = -5.$

47. $2x - 6\sqrt{2x-1} = 8.$

51. $x^3 + x\sqrt{x} - 72 = 0.$

48. $x = 11 - 3\sqrt{x+7}.$

52. $x^{-\frac{2}{3}} - 5x^{-\frac{1}{3}} + 4 = 0.$

53. $\left(\frac{12}{x} - 1\right)^2 + 8\left(\frac{12}{x} - 1\right) = 33.$

54. $\left(x + \frac{1}{x}\right)^2 - 2\left(x + \frac{1}{x}\right) = \frac{5}{4}.$

$$55. \left(\frac{1+x^2}{x}\right)^2 + 2\left(\frac{1+x^2}{x}\right) = 8.$$

$$56. \left(\frac{1-x}{x^2}\right)^2 - 4\left(\frac{1-x}{x^2}\right) = \frac{17}{16}.$$

$$57. \left(x - \frac{1}{x}\right)^2 + \frac{5}{6}\left(x - \frac{1}{x}\right) = 1.$$

$$58. (x-a)^{\frac{2}{3}} - 3a^{\frac{1}{3}}(x-a)^{\frac{1}{3}} + 2a^{\frac{2}{3}} = 0.$$

$$59. \text{Find the three cube roots of } -1.$$

$$60. \text{Find the three cube roots of } -8.$$

$$61. \text{Find the four fourth roots of } 1.$$

Solve the following equations:

$$62. x^6 - 28x^3 + 27 = 0.$$

$$65. x^4 + 2x^3 - x = 30.$$

$$63. x^3 - \frac{8}{x^3} = 7.$$

$$66. x^4 - 4x^3 + 8x = -3.$$

$$67. x^4 - 2x^3 + x = 132.$$

$$64. x^4 - 16 = 0.$$

$$68. x^4 - 6x^3 + 27x = 10.$$

$$69. x^4 + 2x^3 + 5x^2 + 4x - 60 = 0.$$

$$70. x^4 + 6x^3 + 7x^2 - 6x - 8 = 0.$$

$$71. x^4 - 6x^3 + 15x^2 - 18x + 8 = 0.$$

$$72. x^4 - 10x^3 + 35x^2 - 50x + 24 = 0.$$

$$73. 16x^4 - 8x^3 - 31x^2 + 8x + 15 = 0.$$

$$74. 4x^4 - 4x^3 - 7x^2 + 4x + 3 = 0.$$

$$75. \frac{x^2}{x+1} - \frac{x+1}{x^2} = \frac{7}{12}.$$

SUGGESTION. — Since the second term is the *reciprocal* of the first, put p for the first term and $-\frac{1}{p}$ for the second.

Then,

$$p - \frac{1}{p} = \frac{7}{12}.$$

$$76. \frac{x^2+x}{2} + \frac{2}{x^2+x} = 2.$$

$$78. \frac{x+2}{x^2+4} + \frac{2(x^2+4)}{x+2} = \frac{51}{5}.$$

$$77. \frac{x^2+1}{4} + \frac{4}{x^2+1} = \frac{5}{2}.$$

$$79. \frac{x^2+1}{x-1} - \frac{4(x-1)}{x^2+1} = \frac{21}{5}.$$

$$80. \frac{x}{x^2-1} + \frac{x^2-1}{x} = -\frac{13}{6}. \quad 83. x^2 - 2x + \frac{6}{x^2-2x-6} = 11.$$

$$81. x^2 + x + 1 - \frac{1}{x^2+x+1} = \frac{8}{3}. \quad 84. x^2 - x + \frac{2}{x^2-x-4} = 7.$$

$$82. x^2 - 3x + \frac{2}{x^2-3x+2} = 1. \quad 85. x^2 - 2x + \frac{4}{x^2-2x+1} = 4.$$

$$86. \frac{1}{1+x+x^2} + \frac{2}{\sqrt{1+x+x^2}} - 3 = 0.$$

SIMULTANEOUS EQUATIONS INVOLVING QUADRATICS

299. An equation whose terms are homogeneous with respect to the unknown numbers is called a **Homogeneous Equation**.

$x^2 - xy + y^2 = 6$, $x^3 + y^3 = 12$, and $ax + y = 10$ are homogeneous equations.

300. An equation that is not affected by interchanging the unknown numbers involved in it is called a **Symmetrical Equation**.

$2x^2 + xy + 2y^2 = 32$, $x^3 + y^3 = 28$ are symmetrical equations.

301. Many simultaneous equations involving quadratics may be solved by the rules for quadratics, if they belong to one of the following classes:

1. *When one is simple and the other quadratic.*
2. *When the unknown numbers are involved in a similar manner in each equation.*
3. *When each equation is homogeneous and quadratic.*

I. Simple and quadratic.

1. Solve the equations
$$\begin{cases} x + y = 7, \\ 3x^2 + y^2 = 43. \end{cases}$$

SOLUTION

$$x + y = 7. \quad (1)$$

$$3x^2 + y^2 = 43. \quad (2)$$

From (1), $y = 7 - x. \quad (3)$

Substituting in (2), $3x^2 + (7 - x)^2 = 43. \quad (4)$

Solving, $x = 3$ or $\frac{1}{2}$. (5)

Substituting 3 for x in (3), $y = 4$. (6)

Substituting $\frac{1}{2}$ for x in (3), $y = \frac{1}{2}$. (7)

That is, x and y each have two values $\begin{cases} \text{when } x = 3, y = 4, \\ \text{when } x = \frac{1}{2}, y = \frac{1}{2}. \end{cases}$

Equations of this class may be solved by finding the value of one unknown number in terms of the other in one equation, and then substituting it in the other.

Solve the following equations:

2. $\begin{cases} x^2 + y^2 = 20, \\ x = 2y. \end{cases}$

5. $\begin{cases} x = 6 - y, \\ x^2 + y^2 = 72. \end{cases}$

3. $\begin{cases} 10x + y = 3xy, \\ y - x = 2. \end{cases}$

6. $\begin{cases} xy(x - 2y) = 10, \\ xy = 10. \end{cases}$

4. $\begin{cases} x^2 + xy = 12, \\ x - y = 2. \end{cases}$

7. $\begin{cases} 3x(y + 1) = 12, \\ 3x = 2y. \end{cases}$

II. Unknown numbers similarly involved.

8. Solve the equations $\begin{cases} x + y = 7, \\ xy = 10. \end{cases}$

SOLUTION

$x + y = 7$. (1)

$xy = 10$. (2)

Squaring (1), $x^2 + 2xy + y^2 = 49$. (3)

Multiplying (2) by 4, $4xy = 40$. (4)

Subtracting (4) from (3), $x^2 - 2xy + y^2 = 9$. (5)

Extracting the square root, $x - y = \pm 3$. (6)

From (1) + (6), $x = 5$ or 2 .

From (1) - (6), $y = 2$ or 5 .

Such equations may be solved by the method illustrated in example 1, but the method given above of finding the value of

$x - y$, so that it may be combined with the value of $x + y$ to discover the values of x and y , is preferable.

9. Solve the equations
$$\begin{cases} x^2 + y^2 = 25, \\ x + y = 7. \end{cases}$$

SOLUTION

$$x^2 + y^2 = 25. \quad (1)$$

$$x + y = 7. \quad (2)$$

Squaring (2), $x^2 + 2xy + y^2 = 49. \quad (3)$

Subtracting (1) from (3), $2xy = 24. \quad (4)$

Subtracting (4) from (1), $x^2 - 2xy + y^2 = 1. \quad (5)$

Extracting the square root, $x - y = \pm 1. \quad (6)$

From (2) + (6), $x = 4$ or $3.$

From (2) - (6) $y = 3$ or $4.$

10. Solve the equations
$$\begin{cases} x^4 + y^4 = 97, \\ x + y = 1. \end{cases}$$

SOLUTION

$$x^4 + y^4 = 97. \quad (1)$$

$$x + y = 1. \quad (2)$$

4th power of (2),

$$x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4 = 1. \quad (3)$$

Subtracting (1), $4x^3y + 6x^2y^2 + 4xy^3 = -96 \quad (4)$

Dividing by 2, $2x^3y + 3x^2y^2 + 2xy^3 = -48. \quad (5)$

$2xy \times$ square of (2), $2x^3y + 4x^2y^2 + 2xy^3 = 2xy. \quad (6)$

Subtracting (5) from (6), $x^2y^2 - 2xy = 48. \quad (7)$

Solving for xy , $xy = -6$ or $8. \quad (8)$

Equations (2) and (8) give two pairs of simultaneous equations,

$$\begin{cases} x + y = 1 \\ xy = -6 \end{cases} \quad \text{and} \quad \begin{cases} x + y = 1 \\ xy = 8 \end{cases}$$

Solving these by previous methods,

$$x = 3, \text{ or } -2, \text{ or } \frac{1}{2} + \frac{1}{2}\sqrt{-31}, \text{ or } \frac{1}{2} - \frac{1}{2}\sqrt{-31}.$$

$$y = -2, \text{ or } 3, \text{ or } \frac{1}{2} - \frac{1}{2}\sqrt{-31}, \text{ or } \frac{1}{2} + \frac{1}{2}\sqrt{-31}.$$

Solve the following equations:

$$11. \begin{cases} x^2 + y^2 = 50, \\ xy = 7. \end{cases}$$

$$14. \begin{cases} x^2 + y^2 = 13, \\ x + y + xy = 11. \end{cases}$$

$$12. \begin{cases} x + y = 8, \\ x^2 + y^2 = 34. \end{cases}$$

$$15. \begin{cases} x^4 + y^4 = 17, \\ x + y = 3. \end{cases}$$

$$13. \begin{cases} x + y = 9, \\ x^3 + y^3 = 243. \end{cases}$$

$$16. \begin{cases} x^4 + x^2y^2 + y^4 = 21, \\ x^2 + xy + y^2 = 7. \end{cases}$$

III. Homogeneous quadratics.

$$17. \text{ Solve the equations } \begin{cases} x^2 - xy + y^2 = 21, \\ y^2 - 2xy = -15. \end{cases}$$

SOLUTION

$$x^2 - xy + y^2 = 21. \quad (1)$$

$$y^2 - 2xy = -15. \quad (2)$$

$$\text{Assume} \quad x = vy. \quad (3)$$

$$\text{Substituting in (1),} \quad v^2y^2 - vy^2 + y^2 = 21. \quad (4)$$

$$\text{Substituting in (2),} \quad y^2 - 2vy^2 = -15. \quad (5)$$

$$\text{Solving (4) for } y^2, \quad y^2 = \frac{21}{v^2 - v + 1} \quad (6)$$

$$\text{Solving (5) for } y^2, \quad y^2 = \frac{15}{2v - 1}. \quad (7)$$

$$\text{Comparing the values of } y^2, \quad \frac{15}{2v - 1} = \frac{21}{v^2 - v + 1} \quad (8)$$

$$\text{Clearing, etc.,} \quad 5v^2 - 19v + 12 = 0. \quad (9)$$

$$\text{Factoring,} \quad (v - 3)(5v - 4) = 0. \quad (10)$$

$$\therefore v = 3 \text{ or } \frac{4}{5}. \quad (11)$$

$$\begin{aligned} &\text{Substituting 3 for } v \text{ in (7) or in (6),} && y = \pm\sqrt{3} \\ &\text{and since } x = vy, && x = \pm 3\sqrt{3} \end{aligned}$$

$$\begin{aligned} &\text{Substituting } \frac{4}{5} \text{ for } v \text{ in (7) or in (6),} && y = \pm 5 \\ &\text{and since } x = vy, && x = \pm 4 \end{aligned}$$

$$\text{Hence,} \quad \begin{cases} x = +3\sqrt{3}, \text{ or } -3\sqrt{3}, \text{ or } +4, \text{ or } -4, \\ y = +\sqrt{3}, \text{ or } -\sqrt{3}, \text{ or } +5, \text{ or } -5. \end{cases}$$

Solve the following equations:

$$18. \begin{cases} xy + 3y^2 = 20, \\ x^2 - 3xy = -8. \end{cases}$$

$$21. \begin{cases} x^2 - xy - y^2 = 20, \\ x^2 - 3xy + 2y^2 = 8. \end{cases}$$

$$19. \begin{cases} x^2 + xy = 12, \\ xy + 2y^2 = 5. \end{cases}$$

$$22. \begin{cases} x^2 - xy + y^2 = 21, \\ x^2 + 2y^2 = 27. \end{cases}$$

$$20. \begin{cases} x^2 + 2y^2 = 44, \\ xy - y^2 = 8. \end{cases}$$

$$23. \begin{cases} 2x^2 - 3xy + 2y^2 = 100, \\ x^2 - y^2 = 75. \end{cases}$$

302. Many simultaneous equations that belong to one or more of the preceding classes, and many that belong to none of them, may be readily solved by *special devices*.

$$24. \text{ Solve the equations } \begin{cases} x^2 + xy = 12, \\ xy + y^2 = 4. \end{cases}$$

SOLUTION

$$x^2 + xy = 12. \quad (1)$$

$$xy + y^2 = 4. \quad (2)$$

$$\text{Adding,} \quad x^2 + 2xy + y^2 = 16. \quad (3)$$

$$\therefore x + y = +4 \text{ or } -4. \quad (4)$$

$$\text{Subtracting (2) from (1),} \quad x^2 - y^2 = 8. \quad (5)$$

$$\text{Dividing (5) by (4),} \quad x - y = +2 \text{ or } -2. \quad (6)$$

$$\text{Combining (4) and (6),} \quad x = 3 \text{ or } -3; y = 1 \text{ or } -1.$$

$$25. \text{ Solve the equations } \begin{cases} x^4 + x^2y^2 + y^4 = 48, \\ x^2 - xy + y^2 = 12. \end{cases}$$

SOLUTION

$$x^4 + x^2y^2 + y^4 = 48. \quad (1)$$

$$x^2 - xy + y^2 = 12. \quad (2)$$

$$\text{Dividing (1) by (2),} \quad x^2 + xy + y^2 = 4. \quad (3)$$

$$\text{From (3)-(2),} \quad xy = -4. \quad (4)$$

By adding (4) to (3), and subtracting (4) from (2), the values of $(x+y)^2$ and $(x-y)^2$ may be found and the solution readily completed.

26. Solve the equations
$$\begin{cases} x^3 - y^3 = 26, \\ x - y = 2. \end{cases}$$

SOLUTION

$$x^3 - y^3 = 26. \quad (1)$$

$$x - y = 2. \quad (2)$$

Dividing (1) by (2), $x^2 + xy + y^2 = 13. \quad (3)$

Squaring (2), $x^2 - 2xy + y^2 = 4. \quad (4)$

Subtracting, dividing by 3, $xy = 3. \quad (5)$

By adding (5) to (3), the value of $(x + y)^2$ may be obtained and the solution completed as in previous examples.

27. Solve the equations
$$\begin{cases} x^4 + y^4 = 82, \\ x - y = 2. \end{cases}$$

SOLUTION

$$x^4 + y^4 = 82. \quad (1)$$

$$x - y = 2. \quad (2)$$

Assume $x = u + v, \quad (3)$

and $y = u - v. \quad (4)$

Substituting these values in (1),

$$\begin{aligned} &u^4 + 4u^3v + 6u^2v^2 + 4uv^3 + v^4 \\ &+ u^4 - 4u^3v + 6u^2v^2 - 4uv^3 + v^4 = 82, \end{aligned} \quad (5)$$

and in (2), $2v = 2. \quad (6)$

Dividing (5) by 2, $u^4 + 6u^2v^2 + v^4 = 41. \quad (7)$

Dividing (6) by 2, $v = 1. \quad (8)$

Substituting 1 for v in (7) and solving, $u = \pm 2$ or $\pm \sqrt{-10}. \quad (9)$

Hence, from (3) and (4), $x = 3$ or -1 or $1 \pm \sqrt{-10},$

and $y = 1$ or -3 or $-1 \pm \sqrt{-10}.$

28. Solve the equations
$$\begin{cases} x^2 + y^2 + x + y = 14, \\ xy = 3. \end{cases}$$

SOLUTION

$$x^2 + y^2 + x + y = 14.$$

$$xy = 3.$$

Adding twice the second equation to the first,

$$x^2 + 2xy + y^2 + x + y = 20.$$

Completing the square, $(x + y)^2 + x + y + (\frac{1}{4})^2 = 20\frac{1}{4}$.

Extracting the square root, $x + y + \frac{1}{4} = \pm \frac{1}{2}$.

$$\therefore x + y = 4 \text{ or } -5.$$

The solution may be completed by solving the equations,

$$\begin{cases} x + y = 4 \\ xy = 3 \end{cases} \text{ and } \begin{cases} x + y = -5 \\ xy = 3 \end{cases}$$

The student will doubtless discover many other methods for solving simultaneous equations. All the preceding solutions are but illustrations of devices that are important only because they are often applicable.

Solve the following simultaneous equations:

$$29. \begin{cases} x^2 + y^2 = 53, \\ x - y = 5. \end{cases}$$

$$38. \begin{cases} 2x^2 + xy - 5y^2 = 20, \\ 2x - 3y = 1. \end{cases}$$

$$30. \begin{cases} x^3 + y^3 = 28, \\ x + y = 4. \end{cases}$$

$$39. \begin{cases} \frac{x^2}{y^2} + \frac{4x}{y} = 21, \\ x - y = 2. \end{cases}$$

$$31. \begin{cases} 1 + x = y, \\ x^2 + y^2 = 61. \end{cases}$$

$$40. \begin{cases} x^2 - xy = 48, \\ xy - y^2 = 12. \end{cases}$$

$$32. \begin{cases} 1 + x = y, \\ 1 + x^3 = \frac{y^3}{4}. \end{cases}$$

$$41. \begin{cases} x^2 + xy = -6, \\ xy + y^2 = 15. \end{cases}$$

$$33. \begin{cases} x^2 + y^2 = 40, \\ xy = 12. \end{cases}$$

$$42. \begin{cases} 4xy = 96 - x^2y^2, \\ x + y = 6. \end{cases}$$

$$34. \begin{cases} x(x + y) = x, \\ y(x - y) = -1. \end{cases}$$

$$43. \begin{cases} x^2 - xy = 8, \\ xy + y^2 = 12. \end{cases}$$

$$35. \begin{cases} x^2 + 3xy - y^2 = 43, \\ x + 2y = 10. \end{cases}$$

$$44. \begin{cases} x^2 - xy = 6, \\ x^2 + y^2 = 61. \end{cases}$$

$$36. \begin{cases} x^2 + xy + y^2 = 19, \\ x^3 - y^3 = 19. \end{cases}$$

$$45. \begin{cases} x^2 + xy = 77, \\ xy - y^2 = 12. \end{cases}$$

$$37. \begin{cases} x^2 + 3xy = y^2 + 23, \\ x + 3y = 9. \end{cases}$$

$$46. \begin{cases} 2x - y = 2, \\ 2x^2 + y^2 = \frac{3}{2}. \end{cases}$$

47. $\begin{cases} 2x^2 + 3xy + y^2 = 20, \\ 5x^2 + 4y^2 = 41. \end{cases}$
48. $\begin{cases} 2xy - y^2 = 12, \\ 3xy + 5x^2 = 104. \end{cases}$
49. $\begin{cases} x^2 + xy + y^2 = 151, \\ x^2 + y^2 = 106. \end{cases}$
50. $\begin{cases} x^2 + xy + y^2 = 84, \\ x - \sqrt{xy} + y = 6. \end{cases}$
51. $\begin{cases} 4x^2 - 2xy + y^2 = 13, \\ 8x^2 + y^2 = 65. \end{cases}$
52. $\begin{cases} 6x^2 + 6y^2 = 13xy, \\ x^2 - y^2 = 20. \end{cases}$
53. $\begin{cases} x^2 + y^2 - 3(x + y) = 8, \\ x + y + xy = 11. \end{cases}$
54. $\begin{cases} 3xy + 2x + y = 25, \\ \frac{9x}{y} = \frac{4y}{x}. \end{cases}$
55. $\begin{cases} x^2 + xy = 40, \\ 27 + 2y^2 = 3xy. \end{cases}$
56. $\begin{cases} xy^2 + xy = 24, \\ xy^3 + x = 56. \end{cases}$
57. $\begin{cases} x^4 + y^4 = 82, \\ x + y = 4. \end{cases}$
58. $\begin{cases} x^4 - y^4 = 369, \\ x^2 - y^2 = 9. \end{cases}$
59. $\begin{cases} x^2 + y^2 - 78 = x + y, \\ xy + x + y = 39. \end{cases}$
60. $\begin{cases} x^2 + 2xy + 3y^2 = 43, \\ 2x^2 + 3xy + 4y^2 = 62. \end{cases}$
61. $\begin{cases} x^2 - xy + 2y^2 = 46, \\ x^2 + xy + 3y^2 = 111. \end{cases}$
62. $\begin{cases} x^2 - 7xy + 12y^2 = 0, \\ xy + 3y = 2x + 21. \end{cases}$
63. $\begin{cases} x^2 - y^2 = 37, \\ xy(y - x) = -12. \end{cases}$
64. $\begin{cases} x + y = 25, \\ \sqrt{x} + \sqrt{y} = 7. \end{cases}$
65. $\begin{cases} x^3 + y^3 = 225y, \\ x^2 - y^2 = 75. \end{cases}$
66. $\begin{cases} x^2 + xy + 2y^2 = 11, \\ 2x^2 + 5y^2 = 22. \end{cases}$
67. $\begin{cases} x^2 + y^2 = 3xy + 5, \\ x^4 + y^4 = 2. \end{cases}$
68. $\begin{cases} x^2 + y^2 = \frac{13}{x - y}, \\ x^2y - xy^2 = 6. \end{cases}$
69. $\begin{cases} (x + y)(x^2 + y^2) = 65, \\ (x - y)(x^2 - y^2) = 5. \end{cases}$
70. $\begin{cases} x^2 + y = x - y^2 + 42, \\ xy = 20. \end{cases}$
71. $\begin{cases} x + y + 2\sqrt{x + y} = 24, \\ x - y + 3\sqrt{x - y} = 10. \end{cases}$
72. $\begin{cases} x^2 + y^2 + 6\sqrt{x^2 + y^2} = 55, \\ x^2 - y^2 = 7. \end{cases}$

$$73. \begin{cases} \frac{x+y}{y} - \frac{2x}{x+y} + \frac{x^2-x^2y}{y^3-x^2y} = \frac{3}{8}, \\ x^2 = y^2 + 16. \end{cases}$$

$$74. \begin{cases} x^2 - xy = a^2 + b^2, \\ xy - y^2 = 2ab. \end{cases}$$

$$75. \begin{cases} x - 2y = 2(a+b), \\ xy + 2y^2 = 2b(b-a). \end{cases}$$

$$76. \begin{cases} x^3 + y^3 = 2a(a^2 + 3b^2), \\ x^2y + xy^2 = 2a(a^2 - b^2). \end{cases}$$

$$77. \begin{cases} \frac{a-x}{x} + \frac{b+y}{y} = \frac{ay+bx}{a^2-b^2}, \\ x^2 + y^2 = 2(a^2 + b^2). \end{cases}$$

PROBLEMS

303. 1. The sum of two numbers is 12, and their product is 32. What are the numbers?

2. The sum of two numbers is 17, and the sum of their squares is 157. What are the numbers?

3. The difference of two numbers is 1, and the difference of their cubes 91. What are the numbers?

4. The sum of two numbers is 82, and the sum of their square roots is 10. What are the numbers?

5. It takes 52 rods of fence to inclose a rectangular garden containing 1 acre. How long and how wide is the garden?

6. The sum of the squares of the terms of a fraction is 89, and the fraction is $\frac{3}{4}$ larger than its reciprocal. What is the fraction?

7. Find two numbers such that their product is 8 greater than twice their sum and 48 less than the sum of their squares.

8. If 63 is subtracted from a certain number expressed by two digits, its digits will be transposed; and if the number is multiplied by the sum of its digits, the product will be 729. What is the number?

9. A man expended \$6 for canvas. Had it cost 4 cents less per yard, he would have received 5 yards more. How many yards did he buy, and at what price per yard?

10. If the difference of two numbers multiplied by the greater is 160, and multiplied by the less is 96, what are the numbers?

11. A rectangular flower garden containing 54 square rods was enlarged to twice its former size by making an addition of $1\frac{1}{2}$ rods on all sides. What were the original dimensions of the garden?

12. If it requires 200 rods of fence to inclose a rectangular field of 15 acres, what are its dimensions?

13. A rectangular field contains 20 acres. If its length were 20 rods less and its width 8 rods less, its area would be 8 acres less. What are its dimensions?

14. A man found that he could buy 16 more sheep than cows for \$100, and that the cost of 3 cows was \$15 greater than the cost of 4 times as many sheep. What was the price of each?

15. Eight persons contributed \$30 to pay for a set of books. One half of the amount was contributed by women, and the other half by men, each man giving \$2 more than each woman. What did each woman and what did each man contribute?

16. A man loaned \$1000 in two unequal sums at such rates that both sums yielded the same annual interest. The larger sum at the higher rate of interest would have yielded \$36 annually, the smaller sum at the lower rate, \$16 annually. What sums did he invest, and at what rates of interest?

17. If 2 is added to the numerator and subtracted from the denominator of a certain fraction, the result will be the reciprocal of the fraction; if 3 is subtracted from the numerator and added to the denominator, the result will be $\frac{2}{3}$ of the original fraction. What is the fraction?

18. The product of two numbers is 59 greater than their sum, and the sum of their squares is 170. What are the numbers?

19. A purse contained \$ 50.50 in gold and silver coins. If there were fifteen coins, and if each gold coin was worth as many dollars as there were silver coins and each silver coin was worth as many cents as there were gold coins, how many coins of each kind were there?

20. Two men working together can complete a piece of work in $6\frac{2}{3}$ days. If it would take one man 3 days longer than the other to do the work alone, in how many days can each man do the work alone?

21. The fore wheel of a carriage makes 12 revolutions more than the hind wheel in going 240 yards. If the circumference of each wheel were one yard more than it is, the fore wheel would make 8 revolutions more than the hind wheel in going 240 yards. What is the circumference of each wheel?

22. A sum of money on interest for 1 year at a certain per cent amounted to \$ 11130. If the rate had been 1% less and the principal \$ 100 more, the amount would have been the same. Find the principal and rate.

23. A number multiplied by another composed of the same two digits, but reversed, gives a product of 4032. If the first divided by the second is equal to $1\frac{1}{2}$, what are the numbers?

24. The town A is on a lake and 12 miles from B, which is 4 miles from the opposite shore. A man rows across the lake and walks to B in 3 hours. In returning he walks at the same rate as before, but rows 2 miles an hour less than before. If it takes him 5 hours to return, find his rates of walking and rowing.

25. A, B, and C started together to ride a certain distance. A and C rode the whole distance at uniform rates, A two miles an hour faster than C. B rode with C for 20 miles, and then, by increasing his speed two miles an hour, reached his destination 40 minutes earlier than C and 20 minutes later than A. Find the distance, and the rate at which each traveled.

26. Find two numbers such that their product is equal to the difference of their squares, and the difference of their cubes is equal to the sum of their squares.

PROPERTIES OF QUADRATICS

304. Let $ax^2 + bx + c = 0$. (1)

Dividing by a , $x^2 + \frac{b}{a}x + \frac{c}{a} = 0$. (2)

Putting p for $\frac{b}{a}$, and q for $\frac{c}{a}$,
 $x^2 + px + q = 0$ (3)

represents any quadratic equation involving one unknown number.

Solving (3) and denoting its roots by r_1 and r_2 ,

$$r_1 = \frac{-p + \sqrt{p^2 - 4q}}{2} \text{ and } r_2 = \frac{-p - \sqrt{p^2 - 4q}}{2}. \quad (4)$$

Since (4) exhibits the forms of the roots of every quadratic, it is possible to discover whether the roots of any given quadratic are positive or negative, equal or unequal, rational or irrational, real or imaginary, by considering the relative numerical values of its coefficients p and q when it is reduced to the form (3).

In the following discussion of the nature of the roots of a quadratic equation, the student should keep in mind the distinctions between rational and irrational, real and imaginary. For example, 2 and $\sqrt{4}$ are *rational*, and *real* also; $\sqrt{2}$ and $\sqrt{5}$ are *irrational*, but *real*; $\sqrt{-2}$ and $\sqrt{-5}$ are *irrational*, and also *imaginary*.

I. Suppose q is negative.

If q is negative, $-4q$ is positive; and since p^2 is positive whether p is positive or negative, $p^2 - 4q$ is positive and greater than p^2 .

Hence, under this supposition, $\sqrt{p^2 - 4q}$ is real and numerically greater than p .

Since $\sqrt{p^2 - 4q}$ is numerically greater than p , r_1 is positive and r_2 is negative.

If p , then, is positive, the negative root is numerically greater than the positive root.

If p is negative, the positive root is numerically greater than the negative root. Hence,

If q is negative, the roots are real and unequal, with opposite signs, the numerically greater root having the sign opposite to that of p .

If $p = 0$, the roots are numerically equal, but with opposite signs.

II. Suppose $q = 0$.

Then, since $\sqrt{p^2 - 4q}$ reduces to $\sqrt{p^2}$,

If q is equal to zero, one root is equal to zero, and the other to $-p$.

III. Suppose q is positive and $p^2 > 4q$.

If q is positive and $p^2 > 4q$, $p^2 - 4q < p^2$, but positive.

Hence, $\sqrt{p^2 - 4q}$ is numerically less than p , and real.

Therefore, both roots have the sign opposite to that of p , are real, and are unequal, since they differ in the sign of the radical part. Hence,

If q is positive and $p^2 > 4q$, the roots are real and unequal, and each has the sign opposite to that of p .

IV. Suppose q is positive and $p^2 = 4q$.

Then, since $\sqrt{p^2 - 4q}$ becomes equal to 0,

If q is positive and $p^2 = 4q$, the roots are real and equal, and have the sign opposite to that of p .

V. Suppose q is positive and $p^2 < 4q$.

If q is positive and $p^2 < 4q$, $p^2 - 4q$ is negative and $\sqrt{p^2 - 4q}$ is imaginary. Hence,

If q is positive and $p^2 < 4q$, the roots are imaginary.

VI. If $p^2 - 4q$ is a perfect square or equal to zero, the roots are rational. Otherwise they are irrational.

EXAMPLES

1. What is the nature of the roots of $x^2 - 7x - 8 = 0$?

SOLUTION. — Since in the general equation $q = -8$, by I, the roots are real and unequal, with opposite signs.

Also, since $p = -7$, by I, the positive root is the numerically greater root. Since $p^2 - 4q = 49 + 32 = 81$, by VI, both roots are rational.

2. What is the nature of the roots of $3x^2 + 5x + 3 = 0$?

SOLUTION. — Dividing by 3, $x^2 + \frac{5}{3}x + 1 = 0$.

Since $p^2 - 4q = \frac{25}{9} - 4 = -\frac{11}{9}$, by V, both roots are imaginary.

Find the nature of the roots of the following equations:

3. $x^2 - 5x - 75 = 0$.

9. $x^2 + x - 2 = 0$.

4. $x^2 + 5x + 6 = 0$.

10. $4x^2 - 4x + 1 = 0$.

5. $x^2 + 7x - 30 = 0$.

11. $4x^2 + 6x - 4 = 0$.

6. $x^2 - 3x + 5 = 0$.

12. $2x^2 - 9x + 4 = 0$.

7. $x^2 + 3x - 5 = 0$.

13. $4x^2 + 16x + 7 = 0$.

8. $x^2 + x + 2 = 0$.

14. $9x^2 + 12x + 4 = 0$.

305. Formation of quadratic equations.

§ 304, $r_1 = \frac{-p + \sqrt{p^2 - 4q}}{2}$ and $r_2 = \frac{-p - \sqrt{p^2 - 4q}}{2}$.

Adding the roots, $r_1 + r_2 = \frac{-2p}{2} = -p$.

Multiplying the roots, $r_1 r_2 = \frac{p^2 - (p^2 - 4q)}{4} = q$. Hence,

306. PRINCIPLE. — *The sum of the roots of a quadratic equation having the form $x^2 + px + q = 0$ is equal to the coefficient of x with its sign changed, and their product is equal to the absolute term.*

Substituting $-(r_1 + r_2)$ for p , and $r_1 r_2$ for q in the general equation,

$$x^2 - (r_1 + r_2)x + r_1 r_2 = 0.$$

Expanding, $x^2 - r_1 x - r_2 x + r_1 r_2 = 0$.

Factoring, $(x - r_1)(x - r_2) = 0$.

Hence, to form a quadratic equation when the roots are given:

RULE. — *Subtract each root from x and place the product of the remainders equal to zero.*

EXAMPLES

1. Form an equation whose roots are -5 and 2 .

SOLUTION. $(x + 5)(x - 2) = 0$, or $x^2 + 3x - 10 = 0$.

Or, since the sum of the roots with their signs changed is $+5 - 2$, or 3 , and the product of the roots is -10 (Prin.), the equation is $x^2 + 3x - 10 = 0$.

Form equations whose roots are

- | | | |
|-----------------------------------|---------------------------------------|--------------------------------------|
| 2. 6, 4. | 8. $a, -3a$. | 14. $3 + \sqrt{2}, 3 - \sqrt{2}$. |
| 3. 5, -3. | 9. $a + 2, a - 2$. | 15. $-2 - \sqrt{5}, -2 + \sqrt{5}$. |
| 4. $3, -\frac{1}{3}$. | 10. $b + 1, b - 1$. | 16. $2 \pm 3\sqrt{\frac{1}{3}}$. |
| 5. $\frac{2}{3}, \frac{5}{3}$. | 11. $a + b, a - b$. | 17. $-\frac{1}{2}(3 \pm \sqrt{6})$. |
| 6. $-2, -\frac{1}{2}$. | 12. $\sqrt{a} - \sqrt{b}, \sqrt{b}$. | 18. $\frac{1}{2}(-1 \pm \sqrt{2})$. |
| 7. $-\frac{1}{2}, -\frac{3}{2}$. | 13. $\frac{1}{2}(a \pm \sqrt{b})$. | 19. $a(2 \pm 2\sqrt{5})$. |

GENERAL REVIEW

307. 1. Add $x\sqrt{y} + y\sqrt{x} + \sqrt{xy}$, $x^{\frac{1}{2}}y^{\frac{1}{2}} - \sqrt{x^2y} - \sqrt{xy^2}$, $\sqrt{x^2y} - \sqrt{xy^2} - \sqrt{xy}$, and $y\sqrt{x} - x\sqrt{4y} - \sqrt{9xy}$.

2. From the sum of $2a + 3b - 3y$ and $2y - a - 3b$ subtract $(a - b - y) - (a + b + y)$.

3. What number must be added to a to give $b - a$?

4. If $a = 2$ and $b = 3$, find the value of

$$\frac{a + 2b}{a} + \frac{b}{a - b} - \frac{a^3 - a^2b}{2a - b}.$$

5. What number must be subtracted from $a - b$ to give $b - a + c$?

6. Simplify $a - \{b - a - [a - b - (2a + b) + (2a - b) - a] - b\}$.

7. A grocer sold m pounds of sugar at a cents a pound, and a pounds of tea at b cents a pound. If the sugar cost him b cents a pound and the tea m cents a pound, what was his gain by the transaction?

8. Multiply $a^3 + 2a^2b + 2ab^2 + b^3$ by $a^3 - 2a^2b + 2ab^2 - b^3$.

9. Multiply $x^{m-1} - 2y^{n-2}$ by $2x + y^2$.

10. Multiply $x\sqrt{x} + x\sqrt{y} + y\sqrt{x} + y\sqrt{y}$ by $\sqrt{x} - \sqrt{y}$.

11. Multiply $2x^{\frac{a}{2b}} - 5y^{\frac{a+b}{2}}$ by $2x^{\frac{a}{2b}} + 5y^{\frac{a+b}{2}}$.

12. Expand $(x^a - y^n)(x^n + y^n)(x^{2n} + y^{2n})$.

13. Divide $x^4 - y^4$ by $x - y$.

14. Divide $x^5 - 3x^2 - 20$ by $x - 2$, using detached coefficients.
15. Prove that $x^{50} - b^{50}$ is divisible by $x + b$.
16. Divide $(a + b) + x$ by $(a + b)^{\frac{1}{2}} + x^{\frac{1}{2}}$.
17. Factor $9x^2 - 12x + 4$.
18. Factor $9x^2 + 9x + 2$.
19. Factor $x^3 - 3x + 2$.
20. Prove that $x - a$ is a factor of $x^n + 3ax^{n-1} - 4a^n$.
21. Separate $a^{12} - 1$ into six rational factors.
22. Factor $4(ad + bc)^2 - (a^2 + b^2 + c^2 + d^2)^2$.
23. Find the H. C. D. of $x^2 - y^2$, $x^2 + 2xy + y^2$, and $y^2 + xy$.
24. Find the H. C. D. of $3x^2 - x - 2$ and $6x^2 + x - 2$.
25. Find the H. C. D. of $4x^4 - 11x^3 + 11x - 12$,
 $2x^4 + x^3 - 4x^2 + 7x - 15$, and $2x^4 + x^3 - x - 12$.
26. Find the L. C. M. of $4a^2bx$, $6ab^2y^2$, and $2axy$.
27. Find the L. C. M. of $x^2 - y^2$, $x + y$, and $xy - y^2$.
28. Reduce $\frac{x^3 - 5x + 6}{3x^2 - 4x - 4}$ to its lowest terms.
29. Reduce $\frac{x^3 - 5x^2 + 4}{x^3 - 2x^2 + 1}$ to its lowest terms.
30. Reduce $\frac{a^2 - b^2 - c^2 - 2bc}{a^2 - b^2 + c^2 + 2ac}$ to its lowest terms.
31. Simplify $\frac{x}{x+1} - \frac{x}{1-x} + \frac{x^2}{x^2-1}$.
32. Simplify $\frac{x+y}{2x-2y} + \frac{x-y}{2x+2y} + \frac{4x^2y^2}{y^4-x^4}$.
33. Simplify $\frac{1}{(a-b)(b-c)} - \frac{1}{(c-b)(c-a)} + \frac{1}{(c-a)(a-b)}$.
34. Simplify $\left(a + \frac{1}{a}\right)\left(a^2 + \frac{1}{a^2}\right)\left(a - \frac{1}{a}\right)$.
35. Simplify $\frac{1}{x - \frac{1}{x + \frac{1}{x}}} - \frac{1}{x + \frac{1}{x - \frac{1}{x}}}$.

36. Simplify $\left[\frac{x}{1+\frac{1}{x}} + 1 - \frac{1}{x+1} \right] + \left[\frac{x}{1-\frac{1}{x}} - x - \frac{1}{x-1} \right]$.
37. Prove that $\frac{a}{b} \times \frac{p}{q} = \frac{ap}{bq}$.
38. Prove that $\frac{a}{b} + \frac{m}{n} = \frac{an}{bm}$.
39. Divide $\frac{\sqrt{x}}{\sqrt{y}} - \frac{\sqrt{y}}{\sqrt{x}}$ by $\frac{1}{\sqrt{y}} - \frac{1}{\sqrt{x}}$.
40. Raise $a - b$ to the seventh power.
41. Expand $(2a + 3b)^4$.
42. Expand $(\sqrt{x} + \sqrt[3]{y})^3$.
43. Square $\sqrt{a+b-c}$.
44. Extract the square root of $a^2 + 2a\sqrt{ab} + 3ab + 2b\sqrt{ab} + b^2$.
45. Extract the cube root of $8a^3 - 36a^2b + 54ab^2 - 27b^3$.
46. Extract the square root of $a + b$ to four terms.
47. Find the sixth root of 4826809.
48. Reduce $\sqrt{\frac{2}{3}}$ to its simplest form.
49. Reduce $\sqrt[4]{25a^4}$ to its simplest form.
50. Find the approximate value of $\frac{1}{\sqrt{2}}$.
51. Multiply $2 + \sqrt{8}$ by $1 - \sqrt{2}$.
52. Simplify $\frac{\sqrt{3} + 3\sqrt{2}}{\sqrt{6} + 2}$.
53. Prove that $x^0 = 1$.
54. Prove that $ax^{-5} = \frac{a}{x^5}$.
55. Prove that $x^{\frac{2}{3}} = \sqrt[3]{x^2}$; also that $x^{\frac{2}{3}} = (\sqrt[3]{x})^2$.
56. Find the value of $125^{\frac{2}{3}}$; of $\left(\frac{x^3}{32}\right)^{-\frac{1}{2}}$.

Solve the following equations :

$$57. \frac{1}{a-b} + \frac{a-b}{x} = \frac{1}{a+b} + \frac{a+b}{x}.$$

$$58. \frac{x-\frac{1}{a}}{c} + \frac{x-\frac{1}{b}}{a} + \frac{x-\frac{1}{c}}{b} = 0.$$

$$59. mx^2 - nx = mn.$$

$$63. \sqrt{x-9} = \sqrt{x} - 1.$$

$$60. x^4 + \frac{1}{2} = \frac{3x^2}{2}.$$

$$64. x^2 + \sqrt{x^2 + 16} = 14.$$

$$61. x^6 + 8 = 9x^2.$$

$$65. \left(\frac{4}{x} + x\right)^2 - \left(\frac{4}{x} + x\right) = 20.$$

$$62. (1+x)^5 + (1-x)^5 = 242.$$

$$66. x + x^2 + (1+x+x^2)^2 = 55.$$

$$67. \frac{1+x}{1+x+\sqrt{1+x^2}} = a - \frac{1+x}{1-x+\sqrt{1+x^2}}.$$

$$68. \begin{cases} x+y=8, \\ y+z=4, \\ z+x=6. \end{cases}$$

$$75. \begin{cases} x^2 + x = 26 - y^2 - y, \\ xy = 8. \end{cases}$$

$$69. \begin{cases} \frac{1}{x} + \frac{1}{y} = 10, \\ \frac{3}{x} + \frac{2}{y} = 10. \end{cases}$$

$$76. \begin{cases} \sqrt{xy} = 12, \\ x+y-\sqrt{x+y} = 20. \end{cases}$$

$$77. \begin{cases} x^2 - y^2 = 7, \\ x^4 - y^4 = 175. \end{cases}$$

$$78. \begin{cases} xy - xy^2 = -6, \\ x - xy^2 = 9. \end{cases}$$

$$70. \begin{cases} 2x + 3y + z = 9, \\ x + 2y + 3z = 13, \\ 3x + y + 2z = 11. \end{cases}$$

$$79. \begin{cases} xy = x + y, \\ x^2 + y^2 = 8. \end{cases}$$

$$71. \begin{cases} ax + y + z = 2(a+1), \\ x + ay + z = 3a + 1, \\ x + y + az = a^2 + 3. \end{cases}$$

$$80. \begin{cases} x^2y^2 - 4xy = 5, \\ x^2 + 4y^2 = 29. \end{cases}$$

$$81. \begin{cases} 2x^3 + 2y^3 = 9xy, \\ x + y = 3. \end{cases}$$

$$72. \begin{cases} x^2 + xy = 24, \\ y^2 + xy = 12. \end{cases}$$

$$82. \begin{cases} x^2 + xy + y^2 = 189, \\ x + \sqrt{xy} + y = 21. \end{cases}$$

$$73. \begin{cases} x^2 + 3xy = 7, \\ xy + 4y^2 = 18. \end{cases}$$

$$83. \begin{cases} x^{\frac{1}{2}} + y^{\frac{1}{2}} = 4, \\ x^{\frac{3}{2}} + y = 16. \end{cases}$$

$$74. \begin{cases} x^2y + xy^2 = 6, \\ x^3 + y^3 = 9. \end{cases}$$

$$84. \begin{cases} \sqrt{x} - \sqrt{y} = \frac{2}{3}(x-y), \\ \sqrt{xy} = \frac{1}{6}. \end{cases}$$

85. A and B hired a carriage for themselves and four friends. If all had paid, A and B would each have had 4 dollars less to pay. What was the cost of hiring the carriage?

86. What number is that to which if 12 is added and from $\frac{1}{12}$ of the sum 12 is subtracted, the remainder is 12?

87. A grocer has two kinds of sirup worth 50 and 80 cents per gallon respectively. How many gallons of each must he take to make a mixture of 45 gallons worth 60 cents a gallon?

88. How many dimes and how many quarters must be taken so that 18 coins are worth \$3?

89. In a certain weight of gunpowder the saltpeter was 5 pounds more than half the weight, the sulphur 2 pounds less than a fifth, and the charcoal 1 pound more than a tenth. Find the number of pounds of each.

90. How far down a river whose current runs 3 miles an hour can a steamboat go and return in 8 hours, if its rate of sailing in still water is 12 miles an hour?

91. A woman being asked what she paid for her eggs, replied, "Six dozen cost as many cents as I can buy eggs for 32 cents." What was the price per dozen?

92. A gentleman had not room in his stables for 8 of his horses, so he built an additional stable $\frac{1}{2}$ the size of the other, and then had room for 8 horses more than he had. How many horses had he?

93. In a mass of copper, lead, and tin, the copper was 5 pounds less than half the whole in weight, and the lead and tin each 5 pounds more than $\frac{1}{3}$ of the remainder. Find the weight of each.

94. At what time between 4 and 5 o'clock do the hands of a clock make a straight line?

95. A person who can walk n miles an hour has a hours at his disposal. How far may he ride in a coach that travels m miles an hour and return on foot within the allotted time?

96. A merchant sold half a car load more than half his grain; then he sold half a car load more than half the remainder, when he found that if he could sell half a car load more than half of what he still had, he would have none left. How many car loads of grain had he?

97. Four years ago A's age was $\frac{1}{2}$ of B's, and 4 years hence it will be $\frac{2}{3}$ of B's age. What is the age of each?

98. A person being asked the time of day replied that the time past noon was $\frac{2}{3}$ of the time to midnight. What was the time of day?

99. If 3 is added to each term of a certain fraction, the value of the fraction will be $\frac{5}{7}$; if 3 is subtracted from each term, the value will be $\frac{2}{3}$. What is the fraction?

100. A boatman rows such a distance down a stream that it takes him 4 hours to return. If it takes him 2 hours to row down and the current is 2 miles an hour, what is his rate of rowing in still water?

101. A man received \$ 2.50 per day for every day he worked, and he agreed to forfeit \$ 1.50 for every day he was idle. If he worked 3 times as many days as he was idle and received \$ 24, how many days did he work?

102. A jeweler has two silver cups, and a cover worth \$ 1.50. The first cup with the cover on it is worth $1\frac{1}{2}$ times as much as the second cup, and the second cup with the cover on it is worth $\frac{1}{2}$ as much as the first cup. Find the value of each cup.

103. Some smugglers discovered a cave that would exactly hold their cargo, which consisted of 13 bales of cotton and 33 casks of wine. While they were unloading, a revenue cutter hove in sight, and they sailed away with 9 casks and 5 bales, leaving the cave two thirds full. How many bales, or how many casks, would the cave hold?

104. Twenty-eight tons of goods are to be transported in carts and wagons, and it is found that it will require 15 carts and 12 wagons, or else 24 carts and 8 wagons. How much can each cart and each wagon carry?

105. There is a number whose three digits are the same. If 7 times the sum of the digits is subtracted from the number, the remainder is 180. What is the number?

106. A and B can do a piece of work in m days, B and C in n days, A and C in p days. In what time can all together do it? How long will it take each alone to do it?

107. Two passengers together have 400 pounds of baggage and are charged, for the excess above the weight allowed free, 40 and 60 cents respectively. If the baggage had belonged to one of them, he would have been charged \$ 1.50. How much baggage is one passenger allowed without charge?

108. Divide 20 into two parts such that the sum of the two fractions formed by dividing each part by the other is $4\frac{1}{2}$.

109. It takes 1000 square tiles of a certain size to pave a hall, or 1440 square tiles whose dimensions are one inch less. Find the area of the hall floor.

110. The sum of two numbers is 16, and the difference of their squares is 128. What are the numbers?

111. Find two numbers such that their sum, their product, and the difference of their squares are all equal.

112. Divide 25 into two parts such that the difference of their square roots is 1.

113. The difference of two numbers is 6, and their product is equal to twice the cube of the less number. What are the numbers?

114. It took a number of men as many days to pave a sidewalk as there were men. Had there been 3 men more, the work would have been done in 4 days. How many men were there?

115. The product of two numbers is 8, and the sum of their squares is 14 greater than the sum of the numbers. What are the numbers?

116. A rectangular lawn 50 feet long and 40 feet wide has a path of uniform width around it. If the area of the walk is 64 square yards, what is its width?

117. A merchant sold goods for 56 dollars and gained as many hundredths of the cost as there were dollars in the cost. Find the cost of the goods.

118. A person swimming in a stream that runs $1\frac{1}{2}$ miles per hour finds that it takes him 3 times as long to swim a certain distance up the stream as it does to swim the same distance down. What is his rate of swimming in still water?

119. A drover bought some oxen for \$ 900. After 5 had died, he sold the rest at a profit of \$ 20 each and thereby gained \$ 350. How many oxen did he buy?

120. A detachment from an army was marching in regular column with 5 men more in depth than in front. On approaching the enemy, the front was increased by 845 men, and the whole was thus drawn up in 5 lines. Find the number of men in the detachment.

121. A round iron bar weighed 36 pounds. If it had been 1 foot longer and of uniform diameter, each foot of it would have weighed $\frac{1}{2}$ a pound less. Find the length of the iron bar and its weight per foot.

122. A farmer has two cubical granaries. The side of one is 3 yards longer than the side of the other, and the difference of their solid contents is 117 cubic yards. What is the length of the side of each?

123. Two workmen, A and B, were employed at different wages. At the end of a certain number of days A received \$ 30, but B, who had been idle two days in the meantime, received only \$ 19.20. If B had worked the whole time, and A had been idle two days, they would have received equal sums. Find the number of days, and the daily wages of each.

124. By traveling 5 miles an hour less than its usual rate a train was 50 minutes late in running 300 miles. Find the usual rate of speed and the time required to make the trip.

125. Find two numbers such that their sum, their product, and the sum of their squares are all equal.

126. A merchant bought two lots of tea, paying for both \$34. One lot was 20 pounds more than the other, and the number of cents paid per pound was in each case equal to the number of pounds bought. How many pounds of each did he buy?

127. A and B hired a pasture into which A put 4 horses, and B as many as cost him 18 shillings per week. Afterward B put in 2 additional horses, and found that he must pay 20 shillings per week. How much was paid for the pasture per week?

128. By lowering the selling price of apples 1 cent a dozen, an apple woman finds that she can sell 60 more than she used to sell for 60 cents. At what price per dozen did she sell them at first?

129. A and B are two stations 300 miles apart. Two trains start at the same time, one from A, the other from B, and travel to the opposite station. If the first train reaches B 9 hours after the trains meet, and the second train reaches A 4 hours after they meet, when do they meet, and what is the rate of each train?

130. If a carriage wheel $14\frac{2}{3}$ feet in circumference takes one second longer to revolve, the rate of traveling will be $2\frac{1}{3}$ miles less per hour. How fast is the carriage traveling?

131. A railway train, after traveling 2 hours, is detained 1 hour by an accident. It then proceeds at $\frac{2}{3}$ of its former rate, and arrives $7\frac{1}{3}$ hours behind time. If the accident had occurred 50 miles farther on, the train would have arrived $6\frac{1}{3}$ hours behind time. What was the whole distance traveled by the train?

132. A person rents a certain number of acres of land for \$200. He retains 5 acres for his own use and sublets the rest at \$1 an acre more than he gave. If he receives \$10 more than he pays for the whole, how many acres does he rent, and at what rate per acre?

133. A and B left Chicago and walked in the same direction at uniform rates. B started 2 hours after A and overtook him at the 30th milestone. Had each traveled half a mile more per hour, B would have overtaken A at the 42d milestone. At what rate did each travel?

RATIO AND PROPORTION

308. 1. What is the relation of $10x$ to $5x$? of $3x$ to $12x$? of $8a$ to $2a$? of $14m$ to $7m$? of $4a$ to $8a$? of $2b$ to $6b$?

2. In finding the relation, or *ratio*, of $10a$ to $5a$, which is the dividend, the number that *precedes*, or the number that *follows*? Which is the divisor?

3. What is the ratio of a to b ? Since b may not be exactly contained in a , how may the ratio be expressed?

4. Since the ratio of two numbers may be expressed in the form of a fraction, what operations may be performed upon the terms of a ratio without changing the ratio?

309. The relation of two numbers that is expressed by the quotient of the first divided by the second is called their **Ratio**.

310. The **Sign of Ratio** is a colon ($:$).

A ratio is also expressed in the form of a fraction.

The ratio of a to b is written $a : b$ or $\frac{a}{b}$.

The colon is sometimes regarded as derived from the sign of division by omitting the line.

311. The first term of a ratio is called the **Antecedent**.

It corresponds to a dividend, or numerator.

312. The second term of a ratio is called the **Consequent**.

It corresponds to a divisor, or denominator.

313. The antecedent and consequent form a **Couplet**.

In the ratio $a : b$, or $\frac{a}{b}$, a is the antecedent, b the consequent, and the terms a and b form a couplet.

314. The ratio of the reciprocals of two numbers is called the **Reciprocal**, or **Inverse Ratio** of the numbers.

It may be expressed by interchanging the terms of the ratio of the numbers.

The inverse ratio of a to b is $\frac{1}{a} : \frac{1}{b}$. Since $\frac{1}{a} + \frac{1}{b} = \frac{b}{a}$, the inverse ratio of a to b may be written $\frac{b}{a}$, or $b : a$.

315. The ratio of the squares of two numbers is called the **Duplicate** ratio; the ratio of their cubes, the **Triplicate** ratio; the ratio of their square roots, the **Subduplicate** ratio; the ratio of their cube roots, the **Subtriplicate** ratio of the numbers.

The duplicate ratio of a to b is $a^2 : b^2$; the triplicate ratio, $a^3 : b^3$; the subduplicate ratio, $\sqrt{a} : \sqrt{b}$; the subtriplicate ratio, $\sqrt[3]{a} : \sqrt[3]{b}$.

316. PRINCIPLE. — *Multiplying or dividing both terms of a ratio by the same number does not change the ratio.*

EXAMPLES

1. What is the ratio of $8m$ to $4m$? of $4m$ to $8m$?
2. Express the ratio of $6:9$ in its lowest terms; $12x:16y$; $am:bm$; $20ab:10bc$; $(m+n):(m^2-n^2)$.
3. Which is the greater ratio, $2:3$ or $3:4$? $4:9$ or $2:5$?
4. What is the ratio of $\frac{1}{2}$ to $\frac{1}{4}$? $\frac{1}{2}$ to $\frac{1}{3}$? $\frac{2}{3}$ to $\frac{3}{4}$?

SUGGESTION. — When fractions have a common denominator, they have the ratio of their numerators.

5. Reduce $a:b$ and $x:y$ to ratios having the same consequent.
6. When the antecedent is $6x$ and the ratio is $\frac{1}{3}$, what is the consequent?

317. It is evident from § 316, that the ratio of two rational fractions may be expressed by the ratio of two integers.

For example, $\frac{m}{n} : \frac{b}{y}$ may be reduced to the form $\frac{m}{n} \times ny : \frac{b}{y} \times ny$, or $my : bn$.

But the ratio of two numbers, when one is rational and the other irrational or when they are dissimilar surds, cannot be expressed by the ratio of two integers.

Thus, the ratio $\sqrt{2} : 3$ cannot be expressed by any two integers.

318. If the ratio of two numbers can be expressed by the ratio of two integers, the numbers are called **Commensurable Numbers**, and their ratio a **Commensurable Ratio**.

319. If the ratio of two numbers cannot be expressed by the ratio of two integers, the numbers are called **Incommensurable Numbers**, and their ratio an **Incommensurable Ratio**.

The ratio $\sqrt{2} : 3 = \frac{\sqrt{2}}{3} = \frac{1.414213+}{3}$ cannot be expressed by any two integers, because there is no number that, used as a common *measure*, will be contained in both $\sqrt{2}$ and 3 an integral number of times. Hence, $\sqrt{2}$ and 3 are incommensurable, and $\sqrt{2} : 3$ is an incommensurable ratio.

It is evident that by continuing the process of extracting the square root of 2, the ratio $\sqrt{2} : 3$ may be expressed by two integers to any desired degree of approximation, but never with absolute accuracy.

320. 1. What two numbers have the same relation to each other as 2 to 3?

2. Name several couplets that express the same ratio as 2 : 5. How may it be indicated that the ratio of 2 to 5 is the *same* as that of $2a$ to $5a$?

3. What number has the same ratio to $12a$ that $5b$ has to $3b$?

4. What number has the same ratio to $10a$ that $10a$ has to 2? How does this number times 2 compare with $10a$?

321. An equality of ratios is called a **Proportion**.

$3 : 10 = 6 : 20$ and $a : x = b : y$ are proportions.

The double colon ($::$) is often used instead of the sign of equality.

The double colon has been supposed to represent the extremities of the lines that form the sign of equality.

The proportion $a : b = c : d$, or $a : b :: c : d$, is read, the ratio of a to b is equal to the ratio of c to d , or a is to b as c is to d .

322. The antecedents and consequents of the ratios that form a proportion are called the **Antecedents** and **Consequents**, respectively, of the proportion.

In $a : b = c : d$, the antecedents of the proportion are a and c , and the consequents are b and d .

323. The first and fourth terms of a proportion are called the **Extremes** of the proportion.

In the proportion $a : b = c : d$, the extremes are a and d .

324. The second and third terms of a proportion are called the **Means** of the proportion.

In the proportion $a : b = c : d$, the means are b and c .

325. The terms of a proportion are also called **Proportionals**. In the proportion $a : b = b : c$, b is called a *Mean Proportional* between a and c , and c is called a *Third Proportional* to a and b .

In the proportion $a : b = c : d$, d is called a *Fourth Proportional* to a , b , and c .

Since a proportion is an equality of ratios each of which may be expressed as a fraction, a proportion may be expressed as an equation each member of which is a fraction. Hence, it follows that:

326. GENERAL PRINCIPLE.—*The changes that may be made upon a proportion without destroying the equality of its ratios are based upon the changes that may be made upon the members of an equation without destroying their equality and upon the terms of a fraction without altering the value of the fraction.*

PRINCIPLES OF PROPORTION

- 327.** 1. Let any four numbers form a proportion, as $a : b = c : d$.
 2. Express the proportion as a fractional equation.
 3. If this equation is cleared of fractions, what terms of the proportion does the first member contain? the second member?

PRINCIPLE 1.—*In any proportion the product of the extremes is equal to the product of the means.*

If $a : b = c : d$, then, $ad = bc$.

Since a mean proportional serves as both means of a proportion, if $a : b = b : c$, $b^2 = ac$, or $b = \sqrt{ac}$. Hence,

The mean proportional between two numbers is equal to the square root of their product.

Principle 1 may be established as follows:

Let $a : b = c : d$ represent any proportion.

Then, § 310, $\frac{a}{b} = \frac{c}{d}$.

Clearing of fractions, $ad = bc$.

Therefore, the product of the extremes is equal to the product of the means.

NUMERICAL ILLUSTRATION

$$2 : 5 = 6 : 15.$$

$$2 \times 15 = 5 \times 6.$$

$$30 = 30.$$

328. 1. Transform the proportion $a : b = c : d$ in accordance with Prin. 1.

2. Since $ad = bc$, how may the value of a be found? the value of d ? What terms of the proportion are a and d ?

3. How, then, may either extreme of a proportion be found? How may either mean be found?

PRINCIPLE 2. — *Either extreme of a proportion is equal to the product of the means divided by the other extreme. Either mean is equal to the product of the extremes divided by the other mean.*

If $a : b = c : d$, then, $a = \frac{bc}{d}$, $b = \frac{ad}{c}$, etc.

Demonstrate Prin. 2, and give numerical illustrations.

329. 1. If $ad = bc$, what will be the resulting proportion when both members are divided by bd and reduced?

2. What do the factors of ad , the first member of the equation, form in the proportion? What do the factors of bc form?

PRINCIPLE 3. — *If the product of two numbers is equal to the product of two other numbers, one pair of them may be made the extremes and the other pair the means of a proportion.*

If $ad = bc$, then, $a : b = c : d$, or $b : a = d : c$, etc.

Demonstrate Prin. 3, and give numerical illustrations.

330. 1. Change the proportion $a : b = c : d$ into an integral equation by Prin. 1.

2. Divide the members of this equation by cd and reduce.

3. What terms of the given proportion now form the first couplet? the second couplet?

PRINCIPLE 4.—*If four numbers are in proportion, the ratio of the antecedents is equal to the ratio of the consequents ; that is, the numbers are in proportion by **Alternation**.*

If $a : b = c : d$, then, $a : c = b : d$.

Demonstrate Prin. 4, and give numerical illustrations.

331. 1. Change the proportion $a : b = c : d$ into an integral equation by Prin. 1.

2. Divide the members of this equation, $bc = ad$, by ac , and reduce.

3. What change has taken place in the order of the terms of each couplet?

PRINCIPLE 5.—*If four numbers are in proportion, the ratio of the second to the first is equal to the ratio of the fourth to the third ; that is, the numbers are in proportion by **Inversion**.*

If $a : b = c : d$, then, $b : a = d : c$.

Demonstrate Prin. 5, and give numerical illustrations.

332. 1. Express the proportion $a : b = c : d$ as a fractional equation.

2. Add 1 to each member and reduce the mixed numbers to fractional form. Write in the form of a proportion.

3. How may the terms of this proportion be formed from the terms of the given proportion?

4. Since, when $a : b = c : d$, $b : a = d : c$, if the changes just indicated are made in the second proportion, how may the terms of the resulting proportion be obtained from the terms of the original proportion?

PRINCIPLE 6.—*If four numbers are in proportion, the sum of the terms of the first ratio is to either term of the first ratio as the sum of the terms of the second ratio is to the corresponding term of the second ratio ; that is, the numbers are in proportion by Composition.*

If $a : b = c : d$, then, $a + b : b = c + d : d$ and $a + b : a = c + d : c$.

Demonstrate Prin. 6, and give numerical illustrations.

333. 1. Express the proportion $a : b = c : d$ as a fractional equation.

2. Subtract 1 from each member, and reduce the mixed numbers to fractional form. Write in the form of a proportion.

3. How may the terms of this proportion be formed from the terms of the given proportion ?

4. Since, when $a : b = c : d$, $b : a = d : c$, if the changes just indicated are made in the second proportion, how may the terms of the resulting proportion be obtained from the terms of the original proportion ?

PRINCIPLE 7.—*If four numbers are in proportion, the difference between the terms of the first ratio is to either term of the first ratio as the difference between the terms of the second ratio is to the corresponding term of the second ratio ; that is, the numbers are in proportion by Division.*

If $a : b = c : d$, then, $a - b : b = c - d : d$ and $a - b : a = c - d : c$.

Demonstrate Prin. 7, and give numerical illustrations.

334. 1. Change the proportion $a : b = c : d$ according to Prin. 6, and also according to Prin. 7, using the same consequents in each transformation. Express in fractional form.

2. Divide the first equation by the second.

3. How may the terms of the resulting proportion be formed from the terms of the given proportion ?

PRINCIPLE 8.—*If four numbers are in proportion, the sum of the terms of the first ratio is to their difference as the sum of the*

terms of the second ratio is to their difference; that is, the numbers are in proportion by Composition and Division.

If $a : b = c : d$, $a + b : a - b = c + d : c - d$ and $a + b : b - a = c + d : d - c$.

Demonstrate Prin. 8, and give numerical illustrations.

335. 1. Express the proportion $a : b = c : d$ as a fractional equation.

2. Raise each member to the n th power.

3. Also express the n th root of each member.

4. How may these proportions be formed from the given proportion?

PRINCIPLE 9. — *If four numbers are in proportion, their like powers, and also their like roots, will be in proportion.*

• If $a : b = c : d$, then, $a^n : b^n = c^n : d^n$ and $a^{\frac{1}{n}} : b^{\frac{1}{n}} = c^{\frac{1}{n}} : d^{\frac{1}{n}}$.

Demonstrate Prin. 9, and give numerical illustrations.

336. 1. Express $a : b = c : d$ as a fractional equation.

2. What may be done to a fraction without changing its value?

3. Multiply the terms of the first fraction by m and the terms of the second fraction by n . Write as a proportion.

4. How may the terms of this proportion be formed from the terms of the given proportion?

5. Take the given proportion by alternation and multiply the terms of the first couplet by m and those of the second couplet by n .

6. How may the terms of this proportion be formed from the terms of the given proportion?

7. How may the given proportion be formed from the proportions $ma : mb = nc : nd$ and $ma : nb = mc : nd$?

PRINCIPLE 10. — *In a proportion, if both terms of a couplet, or both antecedents, or both consequents are multiplied or divided by the same number, the resulting four numbers form a proportion.*

If $a : b = c : d$, then, $ma : mb = nc : nd$ and $ma : nb = mc : nd$; also, if $ma : mb = nc : nd$, or if $ma : nb = mc : nd$, then, $a : b = c : d$.

Demonstrate Prin. 10, and give numerical illustrations.

337. 1. Express the proportions $a:b=c:d$ and $x:y=z:w$ as fractional equations.

2. How may two equations be combined (Ax. 4 and 5)? Combine these two equations and write the results as proportions.

3. How may these proportions be formed from the given proportions?

PRINCIPLE 11. — *The products, and also the quotients, of corresponding terms of two proportions form a proportion.*

If $a:b=c:d$ and $x:y=z:w$, $ax:by=cz:dw$, also $\frac{a}{x}:\frac{b}{y}=\frac{c}{z}:\frac{d}{w}$.

Demonstrate Prin. 11, and give numerical illustrations.

338. If $a:b=c:d$ and $c:d=e:f$, how does the ratio $a:b$ compare in value with the ratio $e:f$?

PRINCIPLE 12. — *If two proportions have a common couplet, the other two couplets will form a proportion.*

If $a:b=c:d$ and $c:d=e:f$, then, $a:b=e:f$.

Demonstrate Prin. 12, and give numerical illustrations (Ax. 1).

339. A proportion that consists of three or more equal ratios is called a **Multiple Proportion**.

$2:4=3:6=5:10$ and $a:b=c:d=e:f$ are multiple proportions.

340. A multiple proportion in which each consequent serves also as the antecedent of the following ratio is called a **Continued Proportion**.

$2:4=4:8=8:16$ and $a:b=b:c=c:d$ are continued proportions.

341. 1. Form a multiple proportion, as

$$2:4=3:6=5:10=10:20.$$

2. How does the ratio of the sum of the antecedents to the sum of the consequents compare with the first ratio? with the second ratio? with the ratio of any antecedent to its consequent?

3. Investigate other multiple proportions.

PRINCIPLE 13.—*In any multiple proportion the sum of the antecedents is to the sum of the consequents as any antecedent is to its consequent.*

Principle 13 may be established as follows:

Let $a : b = c : d = e : f = g : h$.

It is to be proved that

$$a + c + e + g : b + d + f + h = a : b, \text{ or } c : d, \text{ etc.}$$

Denoting each of the equal ratios by r ,

$$\frac{a}{b} = r, \quad \frac{c}{d} = r, \quad \frac{e}{f} = r, \quad \frac{g}{h} = r. \quad (1)$$

$$\text{Hence,} \quad a = br, \quad c = dr, \quad e = fr, \quad g = hr. \quad (2)$$

Adding equations (2),

$$a + c + e + g = (b + d + f + h)r. \quad (3)$$

Dividing by $(b + d + f + h)$,

$$\frac{a + c + e + g}{b + d + f + h} = r.$$

Replacing r by any of the equal ratios,

$$\frac{a + c + e + g}{b + d + f + h} = \frac{a}{b} = \frac{c}{d}, \text{ etc.}$$

That is, $a + c + e + g : b + d + f + h = a : b$, or $c : d$, etc.

EXAMPLES

342. 1. In the proportion $3 : 5 = x : 55$, what is the value of x ?

FIRST SOLUTION

$$3 : 5 = x : 55.$$

$$\text{Prin. 2,} \quad x = \frac{3 \cdot 55}{5} = 33.$$

SECOND SOLUTION

$$3 : 5 = x : 55.$$

$$\text{Prin. 10,} \quad 3 : 1 = x : 11.$$

$$\text{Prin. 1,} \quad x = 33.$$

Find the value of x in each of the following proportions:

$$2. \quad 2 : 3 = 6 : x.$$

$$4. \quad 1 : x = x : 9.$$

$$3. \quad 5 : x = 4 : 3.$$

$$5. \quad 8 : 5 = x : 10.$$

6. $x + 1 : x = 8 : 6$.

8. $x + 2 : x = 10 : 6$.

7. $x : x - 1 = 15 : 12$.

9. $x + 2 : x - 2 = 3 : 1$.

10. If $x + 5 : x - 5 = 5 : 3$, find the value of x .

11. What two numbers are mean proportionals between 1 and 25?

12. Show that a mean proportional between any two numbers has the sign \pm .

When $a : b = c : d$, prove that the following proportions are true by deriving them from $a : b = c : d$:

13. $d : b = c : a$.

16. $a^2 : b^2 c^2 = 1 : d^2$.

14. $c : d = \frac{1}{b} : \frac{1}{a}$.

17. $ma : \frac{b}{2} = mc : \frac{d}{2}$.

15. $b^3 : d^3 = a^3 : c^3$.

18. $ac : bd = c^2 : d^2$.

19. $\sqrt{ad} : \sqrt{b} = \sqrt{c} : 1$.

20. $a + b : c + d = a - b : c - d$.

21. $a^3 + a^2b + ab^2 + b^3 : a^3 = c^3 + c^2d + cd^2 + d^3 : c^3$.

22. $2a + 3c : 2a - 3c = 8b + 12d : 8b - 12d$.

23. Solve the equation $\frac{a^2 + x^2}{ax} = \frac{41}{20}$.

SOLUTION

$$\frac{a^2 + x^2}{ax} = \frac{41}{20}$$

Dividing by 2,

$$\frac{a^2 + x^2}{2ax} = \frac{41}{40}$$

Regarding this equation as a proportion,

by composition and division, $\frac{a^2 + 2ax + x^2}{a^2 - 2ax + x^2} = \frac{81}{1}$.

Extracting the square root, $\frac{a + x}{a - x} = \frac{9}{1}$.

By composition and division, $\frac{2a}{2x} = \frac{10}{8}$.

$$\therefore x = \frac{4}{5}a.$$

24. Solve the equation $\frac{\sqrt{x+7} + \sqrt{x}}{4 + \sqrt{x}} = \frac{\sqrt{x+7} - \sqrt{x}}{4 - \sqrt{x}}$.

SOLUTION

$$\frac{\sqrt{x+7} + \sqrt{x}}{4 + \sqrt{x}} = \frac{\sqrt{x+7} - \sqrt{x}}{4 - \sqrt{x}}$$

By alternation, Prin. 4, $\frac{\sqrt{x+7} + \sqrt{x}}{\sqrt{x+7} - \sqrt{x}} = \frac{4 + \sqrt{x}}{4 - \sqrt{x}}$.

By composition and division, Prin. 8,

$$\frac{2\sqrt{x+7}}{2\sqrt{x}} = \frac{8}{2\sqrt{x}}$$

Since the consequents are equal, the antecedents are equal.

Therefore, $2\sqrt{x+7} = 8$.

Whence, reducing, $x = 9$.

25. Given $\frac{\sqrt{x+11} + 2}{\sqrt{x+11} - 2} = \frac{\sqrt{2x+14} + 2\frac{1}{2}}{\sqrt{2x+14} - 2\frac{1}{2}}$, to find x .

SOLUTION

$$\frac{\sqrt{x+11} + 2}{\sqrt{x+11} - 2} = \frac{\sqrt{2x+14} + 2\frac{1}{2}}{\sqrt{2x+14} - 2\frac{1}{2}}$$

By composition and division, Prin. 8,

$$\frac{2\sqrt{x+11}}{4} = \frac{2\sqrt{2x+14}}{1\frac{1}{2}}$$

Dividing both terms of each ratio by 2, Prin. 10,

$$\frac{\sqrt{x+11}}{2} = \frac{\sqrt{2x+14}}{\frac{1}{2}}$$

Dividing the consequents by $\frac{1}{2}$, Prin. 10,

$$\frac{\sqrt{x+11}}{3} = \frac{\sqrt{2x+14}}{4}$$

By alternation, Prin. 4, $\frac{\sqrt{x+11}}{\sqrt{2x+14}} = \frac{3}{4}$.

Squaring, and applying Prin. 7, $\frac{x+11}{x+3} = \frac{9}{7}$.

Solving, $x = 25$.

Solve by the principles of proportion :

$$26. \frac{\sqrt{x} + \sqrt{m}}{\sqrt{x} - \sqrt{m}} = \frac{m}{n}.$$

$$29. \frac{\sqrt{x+b} + \sqrt{x-b}}{\sqrt{x+b} - \sqrt{x-b}} = a.$$

$$27. \frac{\sqrt{x} + \sqrt{2a}}{\sqrt{x} - \sqrt{2a}} = \frac{2}{1}.$$

$$30. \frac{\sqrt{a} + \sqrt{a-x}}{\sqrt{a} - \sqrt{a-x}} = \frac{1}{a}.$$

$$28. \frac{x + \sqrt{x-1}}{x - \sqrt{x-1}} = \frac{13}{7}.$$

$$31. \frac{\sqrt{ax} - b}{\sqrt{ax} + b} = \frac{3\sqrt{ax} - 2b}{3\sqrt{ax} + 5b}.$$

$$32. \text{ Given } \frac{\sqrt{x+2} + \sqrt{x+1}}{\sqrt{x+2} + \sqrt{x-1}} = \frac{3}{1}, \text{ to find } x.$$

$$33. \text{ Given } \frac{\sqrt{a} + \sqrt{a+x}}{\sqrt{a} - \sqrt{a+x}} = \frac{\sqrt{b} + \sqrt{x-b}}{\sqrt{b} - \sqrt{x-b}}, \text{ to find } x.$$

$$34. \text{ Given } \frac{a - \sqrt{2ax - x^2}}{a + \sqrt{2ax - x^2}} = \frac{a-b}{a+b}, \text{ to find } x.$$

$$35. \text{ Given } \frac{\sqrt{x+1} + \sqrt{x-2}}{\sqrt{x+1} - \sqrt{x-2}} = \frac{\sqrt{x-3} + \sqrt{x-4}}{\sqrt{x-3} - \sqrt{x-4}}, \text{ to find } x.$$

36. Divide \$35 between two men so that their shares shall be in the ratio of 3 to 4.

37. Two numbers are in the ratio of 3 to 2, and if each is increased by 4, the sums will be in the ratio of 4 to 3. What are the numbers?

38. Divide 16 into two parts such that their product is to the sum of their squares as 3 is to 10.

39. Divide 25 into two parts such that the greater increased by 1 is to the less decreased by 1 as 4 is to 1.

40. The sum of two numbers is 4, and the square of their sum is to the sum of their squares as 8 is to 5. What are the numbers?

41. A dealer had two casks of wine. From the larger he drew 34 gallons, and from the smaller 8 gallons, after which their contents were as 5 to 4. When half the original contents of each cask had been drawn, he put 8 gallons into the larger and 6 into the smaller. If the ratio of their contents was then 5 to 3, what was the capacity of each?

VARIATION

343. One quantity or number is said to **vary directly** as another, or simply to **vary** as another, when they depend on each other in such a manner that if one is changed the other is changed *in the same ratio*.

Thus, if a man earns a certain sum per day, the amount of wages he earns *varies* as the number of days he works.

344. The **Sign of Variation** is \propto . It is read "varies as."

345. The expression $x \propto y$ means that if x is doubled, y is doubled, or if x is divided by a number, y is divided by the same number, etc.; that is, that the ratio of x to y is always the same, or constant.

If the constant ratio is represented by k , then when $x \propto y$, $\frac{x}{y} = k$, or $x = ky$. Hence,

If x varies as y , x is equal to y multiplied by a constant.

346. One quantity or number varies **inversely** as another when it varies as the *reciprocal* of the *other*.

Thus, the time required to do a certain piece of work varies *inversely* as the number of men employed. For, if it takes 10 men 4 days to do a piece of work, it will take 5 men 8 days, or 4 men 10 days, to do it.

In $x \propto \frac{1}{y}$, if the constant ratio of x to $\frac{1}{y}$ is k , $\frac{x}{\frac{1}{y}} = k$, or $xy = k$. Hence,

If x varies inversely as y , their product is constant.

347. One quantity or number varies **jointly** as two others when it varies as their product.

Thus, the amount of money a man earns varies *jointly* as the number of days he works and the sum he receives per day. For, if he should work three times as many days, and receive twice as many dollars per day, he would receive six times as much money.

In $x \propto yz$, if the constant ratio of x to yz is k ,

$$\frac{x}{yz} = k, \text{ or } x = kyz. \text{ Hence,}$$

If x varies jointly as y and z , x is equal to their product multiplied by a constant.

348. One quantity or number varies *directly* as a second and *inversely* as a third when it varies *jointly* as the second and the reciprocal of the third.

Thus, the time required to dig a ditch varies *directly* as the length of the ditch and *inversely* as the number of men employed. For, if the ditch were 10 times as long and 5 times as many men were employed, it would take twice as long to dig it.

In $x \propto y \cdot \frac{1}{z}$, or $x \propto \frac{y}{z}$, if k is the constant ratio,

$$x \div \frac{y}{z} = k, \text{ or } x = k \frac{y}{z}. \text{ Hence,}$$

If x varies directly as y and inversely as z , x is equal to $\frac{y}{z}$ multiplied by a constant.

349. *If x varies as y when z is constant, and x varies as z when y is constant, then x varies as yz when both y and z are variable.*

PROOF. — Since the variation of x depends on the variations of y and z , suppose the latter variations to take place in succession, each in turn producing a corresponding variation in x .

While z remains constant, let y change to y_1 ,* thus causing x to change to x' .

Then,
$$\frac{x'}{x} = \frac{y_1}{y}. \quad (1)$$

Now while y keeps the value y_1 , let z change to z_1 , thus causing x' to change to x_1 .

Then,
$$\frac{x_1}{x'} = \frac{z_1}{z}. \quad (2)$$

* In algebraic notation x_1 , x_2 , x_3 , etc., are read 'x sub one,' 'x sub two,' 'x sub three,' etc.

Multiplying (1) by (2),

$$\frac{x}{x_1} = \frac{yz}{y_1 z_1}$$

$$\therefore x = \frac{x_1}{y_1 z_1} \cdot yz = kyz,$$

where k is the constant $\frac{x_1}{y_1 z_1}$.

Hence,

$$x \propto yz.$$

Thus, the area of a triangle varies as the base when the altitude is constant, varies as the altitude when the base is constant, and varies as the product of the base and altitude when both vary.

Similarly, if x varies as each of three or more numbers, y, z, v, \dots when the others are constant, when all vary x varies as their product.

That is,

$$x \propto yzv \dots$$

Thus, the volume of a parallelopiped varies as the length, if the width and thickness are constant; as the width, if the length and thickness are constant; as the thickness, if the length and width are constant; as the product of any two dimensions, if the other is constant; or as the product of the three dimensions, if all vary.

EXAMPLES

350. 1. If x varies inversely as y , and $x = 6$ when $y = 8$, what is the value of x when $y = 12$?

SOLUTION

Since $x \propto \frac{1}{y}$, let k be the constant ratio of x to $\frac{1}{y}$.

Then, § 346,

$$xy = k. \quad (1)$$

Hence, when $x = 6$ and $y = 8$,

$$k = 6 \times 8, \text{ or } 48. \quad (2)$$

Since k is constant,

$$k = 48 \text{ when } y = 12.$$

Hence, Eq. (1) becomes

$$12x = 48.$$

Therefore, when $y = 12$,

$$x = 4.$$

2. The volume of a cone varies jointly as its altitude and the square of the diameter of its base. When the altitude is 15 and the diameter of the base is 10, the volume is 392.7. What is the volume, when the altitude is 5 and the diameter of the base is 20?

SOLUTION

Let V , H , and D denote the volume, altitude, and diameter of the base, respectively, of any cone, and v the volume of a cone whose altitude is 5 and the diameter of whose base is 20.

Since $V \propto HD^2$, or $\dot{V} = kHD^2$,
 and $V = 392.7$ when $H = 15$ and $D = 10$,

$$392.7 = k \times 15 \times 100. \quad (1)$$

Also, since V becomes v when $H = 5$ and $D = 20$,

$$v = k \times 5 \times 400. \quad (2)$$

Dividing (2) by (1),
$$\frac{v}{392.7} = \frac{5 \times 400}{15 \times 100} = \frac{4}{3} \quad (3)$$

$$\therefore v = \frac{4}{3} \text{ of } 392.7 = 523.6.$$

3. If $x \propto y$ and $y \propto z$, prove that $x \propto z$.

PROOF

Since $x \propto y$ and $y \propto z$, let m represent the constant ratio of x to y , and n the constant ratio of y to z .

Then, § 345,
$$x = my, \quad (1)$$

and
$$y = nz. \quad (2)$$

Substituting nz for y in (1),
$$x = mnz. \quad (3)$$

Hence, since mn is constant, $x \propto z$.

4. The circumference of a circle varies as its diameter. If the circumference of a circle whose diameter is 1 foot is 3.1416 feet, what is the circumference of a circle whose diameter is 100 feet?

5. The area of a circle varies as the square of its diameter. If the area of a circle whose diameter is 10 feet is 78.54 square feet, what is the area of a circle whose diameter is 20 feet?

6. The distance a body falls from rest varies as the square of the time of falling. If a stone falls 64.32 feet in 2 seconds, how far will it fall in 5 seconds?

7. The area of a triangle varies jointly as its base and altitude. The area of a triangle whose base is 12 inches and altitude 6 inches is 36 square inches. What is the area of a triangle whose base is 8 inches and altitude 10 inches? What is the constant ratio?

8. A wrought iron bar 1 square inch in cross section and 1 yard long weighs 10 pounds. If the weight of a uniform bar of given material varies jointly as its length and the area of its cross section, what is the weight of a wrought iron bar 36 feet long, 4 inches wide, and 4 inches thick?

9. The weight of a beam varies jointly as the length, the area of the cross section, and the material of which it is composed. If wood is $\frac{1}{12}$ as heavy as wrought iron, what is the weight of a wooden beam 24 feet long, 12 inches wide, and 12 inches thick?

10. What is the weight of a brick 2 in. \times 4 in. \times 8 in., if the material is $\frac{1}{4}$ as heavy as wrought iron?

11. If 10 men can do a piece of work in 20 days, how long will it take 25 men to do it?

12. If a men can do a piece of work in b days, how many men will be required to do it in c days?

13. The distances, from the fulcrum of a lever, of two weights that balance each other vary inversely as the weights. If two boys weighing 80 and 90 pounds, respectively, are balanced on the ends of a board $8\frac{1}{2}$ feet long, how much of the board has each?

14. A water carrier carries two buckets of water suspended from the ends of a 4-foot stick that rests on his shoulder. If one bucket weighs 60 pounds and the other 100 pounds, and they balance each other, what point of the stick rests on his shoulder?

15. The weight of a body near the earth varies inversely as the square of its distance from the center of the earth. If the radius of the earth is 4000 miles, what would be the weight of a 4-lb. brick at the distance of 4000 miles from the earth's surface?

16. A boy wishes to ascertain the height of a tower. He knows that it is 31 feet 6 inches from his window to the pavement below, and that the distance through which a body falls varies as the square of the time of falling. He drops a marble from his window and finds that it strikes the pavement in 1.4 seconds. Then he throws a stone to the top of the tower and observes that it takes just 3 seconds for it to descend. What is the height of the tower?

17. A horse tied with a rope 45 feet long to a stake in the center of a pasture eats all the grass within reach in $1\frac{1}{2}$ days. If his rope were 15 feet longer, how many days would it take him to eat all the grass within reach?

18. The illumination from a source of light varies inversely as the square of the distance. How far must a screen that is 10 feet from a lantern be moved so as to receive one fourth as much light?

19. The number of times a pendulum oscillates in a given time varies inversely as the square root of its length. If a pendulum 39.1 inches long oscillates once a second, what is the length of a pendulum that oscillates twice a second?

20. How long must a pendulum be to oscillate once in three seconds?

21. If $x \propto \frac{y}{z}$, and if $x = 2$ when $y = 12$ and $z = 2$, what is the value of x when $y = 84$ and $z = 7$?

22. If $x \propto \frac{y}{z}$, and if $x = 60$ when $y = 24$ and $z = 2$, what is the value of y when $x = 20$ and $z = 7$?

23. If x varies jointly as y and z and inversely as the square of w , and if $x = 30$ when $y = 3$, $z = 5$, and $w = 4$, what is the value of x expressed in terms of y , z , and w ?

24. If $x \propto \frac{1}{y}$ and $y \propto \frac{1}{z}$, prove that $x \propto z$.

25. If $x \propto y$ and $z \propto y$, prove that $(x \pm z) \propto y$.

26. Three spheres of lead whose radii are 6, 8, and 10 in., respectively, are united into one. Find the radius of the resulting sphere, if the volume of a sphere varies as the cube of its radius.

27. The volume of a cone varies jointly as its altitude and the square of the diameter of its base. The altitudes of three cones, S , P , and R , are 30 ft., 10 ft., and 5 ft., respectively. The diameter of the base of P is 5 ft. and that of R is 10 ft. If the volume of S is equivalent to that of P and R combined, what is the diameter of the base of S ?

PROGRESSIONS

351. 1. How does each of the numbers 2, 4, 6, 8, 10, 12, ... compare with the number that follows it? How may any *term* after the first be obtained from the preceding term?

2. How may any term of 2, $2\frac{1}{2}$, 3, $3\frac{1}{2}$, ... after the first be obtained from the preceding term?

3. Write a *series* of six terms beginning with a and increasing by a constant number d .

4. How may any term, after the first, of the series 3, 6, 12, 24, 48, ... be obtained from the preceding term?

5. How may any term, after the first, of the series $1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots$ be obtained from the preceding term?

6. Write a series of six terms beginning with a and increasing by a constant multiplier r .

352. A succession of numbers, each of which after the first is derived from the preceding number or numbers according to some fixed law, is called a **Series**.

353. The successive numbers are called the **Terms** of the series. The first and last terms of a series are the *Extremes*, the intervening terms the *Means*.

In the series $a, a + d, a + 2d, a + 3d, a + 4d$, the terms a and $a + 4d$ are the extremes and the other terms are the means.

354. A series consisting of a limited number of terms is called a **Finite Series**.

355. A series consisting of an unlimited number of terms is called an **Infinite Series**.

ARITHMETICAL PROGRESSION

356. A series each term of which after the first is derived from the preceding by the addition of a constant number is called an **Arithmetical Series**, or an **Arithmetical Progression**.

357. The number added to any term to produce the next is called the **Common Difference**.

2, 4, 6, 8, ... and 15, 12, 9, 6, ... are arithmetical progressions. In the first, the common difference is 2 and the series is ascending; in the second, the common difference is - 3 and the series is descending.

A.P. is an abbreviation of the words *Arithmetical Progression*.

358. To find the n th, or last term.

1. In the arithmetical progression $x, x + 2, x + 4, x + 6$, what is the common difference? How many times does it enter into the second term? into the third term? into the fourth term?

2. From the first term of the series $a, a + d, a + 2d, a + 3d, \dots$ how is the second term formed? the third term? the fourth term? the fifth term? the n th term, or any term?

3. What is the n th term of the series $a, a - d, a - 2d, \dots$?

359. When a represents the first term of an A.P., d the common difference, l the n th, or last term, and n the number of terms,

$$l = a + (n - 1)d. \quad (\text{I})$$

EXAMPLES

1. What is the 10th term of the series 3, 6, 9, ...?

<p>PROCESS</p> $l = a + (n - 1)d$ $l = 3 + (10 - 1)3$ $l = 30$	<p>EXPLANATION. — Since the series 3, 6, 9, ... is an A.P. the common difference of whose terms is 3, substituting 3 for a, 3 for d, and 10 for n in the formula for the last term, the last term is found to be 30.</p>
--	---

2. Find the 20th term of the series 7, 11, 15, ...

3. Find the 16th term of the series 2, 7, 12, ...

4. Find the 24th term of the series 1, 16, 31, ...

5. Find the 18th term of the series 1, 8, 15, ...
6. Find the 13th term of the series - 3, 1, 5, ...
7. Find the 49th term of the series 1, $1\frac{1}{3}$, $1\frac{2}{3}$, ...
8. Find the 15th term of the series 45, 43, 41, ...

SUGGESTION. — The common difference is - 2.

9. Find the 10th term of the series 5, 1, - 3, ...
10. Find the 16th term of the series a , $3a$, $5a$, ...
11. Find the 12th term of the series $a - b$, $a + b$, $a + 3b$, ...
12. Find the 7th term of the series $x - 3y$, $x - 2y$, ...
13. A body falls $16\frac{1}{2}$ feet the first second, 3 times as far the second second, 5 times as far the third second, etc. How far will it fall during the 10th second?

360. To find the sum of n terms of a series.

1. Express 5 terms of the series a , $a + d$, $a + 2d$, ...
2. How may the term before the last term be obtained from the last term? If l represents the last term and d the common difference, what will be the term next to the last? the second term from the last? the third term from the last?
3. How, then, may the series a , $a + d$, ... be written in reverse order, if the last term is l ?

361. Let a represent the first term of an A.P., d the common difference, l the last term, n the number of terms, and s the sum of the terms.

Writing the sum of n terms in the usual order and then in the reverse order, and adding the two equal series,

$$s = a + (a + d) + (a + 2d) + (a + 3d) + \cdots + l$$

$$s = l + (l - d) + (l - 2d) + (l - 3d) + \cdots + a$$

$$2s = (a + l) + (a + l) + (a + l) + (a + l) + \cdots + (a + l)$$

$$\therefore 2s = n(a + l).$$

$$s = \frac{n}{2}(a + l), \text{ or } n\left(\frac{a + l}{2}\right). \quad (\text{II})$$

EXAMPLES

1. What is the sum of 20 terms of the series 2, 5, 8, ... ?

PROCESS

$$l = a + (n - 1)d = 2 + (20 - 1) \times 3 = 59$$

$$s = n \left(\frac{a + l}{2} \right) = 20 \left(\frac{2 + 59}{2} \right) = 610$$

EXPLANATION.—Since the last term is not given, it is found by the previous case and substituted for l in the formula for the sum.

2. What is the sum of 16 terms of the series 1, 5, 9, ... ?
3. What is the sum of 10 terms of the series $-2, 0, 2, \dots$?
4. What is the sum of 6 terms of the series $1, 3\frac{1}{2}, 6, \dots$?
5. What is the sum of 8 terms of the series $a, 3a, 5a, \dots$?
6. What is the sum of n terms of the series 1, 7, 13, ... ?
7. What is the sum of a terms of the series $x, x + 2a, \dots$?
8. What is the sum of 7 terms of the series 4, 11, 18, ... ?
9. What is the sum of 10 terms of the series $1, -1, -3, \dots$?
10. What is the sum of 10 terms of the series $1, \frac{1}{2}, 0, \dots$?
11. How many times does a common clock strike in 12 hours ?
12. A body falls $16\frac{1}{2}$ feet the first second, 3 times as far the second second, 5 times as far the third second, etc. How far will it fall in 10 seconds ?
13. Thirty flower pots are arranged in a straight line 4 feet apart. How far must a lady walk who, after watering each plant, returns to a well 4 feet from the first plant and in line with the plants ?
14. A boy took a 30-day job on the following terms: he was to receive 5 cents the first day, 10 cents the second day, 15 cents the third day, etc. How much was he paid for the thirtieth day, and what was the whole amount of his earnings ?

362. The two fundamental formulæ,

$$(I) \ l = a + (n - 1)d \text{ and } (II) \ s = \frac{n}{2}(a + l),$$

contain *five elements*, a , d , l , n , and s . Since these formulæ are independent simultaneous equations, if they contain but two unknown elements they may be solved. Hence, if any *three* of the five elements *are known*, the other two may be found.

EXAMPLES

1. The last term of an A. P. is 58, the common difference is 3, and the sum of the series is 260. Find the number of terms and the first term.

SOLUTION

Substituting 58 for l , 3 for d , and 260 for s in both (I) and (II)

$$(I) \text{ becomes } 58 = a + (n - 1)3. \quad (1)$$

$$(II) \text{ becomes } 260 = \frac{n}{2}(a + 58). \quad (2)$$

$$(1) \times n, \quad 58n = na + 3n^2 - 3n. \quad (3)$$

$$(2) \times 2, \quad 520 = na + 58n. \quad (4)$$

$$(3) - (4), \quad 58n - 520 = 3n^2 - 61n. \quad (5)$$

$$3n^2 - 119n + 520 = 0.$$

$$(n - 5)(3n - 104) = 0.$$

$\therefore n = 5$, the number of terms.

Substituting in (1),

$a = 46$, the first term.

Since the number of terms must be expressed by a positive integer, fractional or negative values of n are rejected.

2. How many terms are there in the A. P. 2, 5, 8, ..., if the sum is 610?

SOLUTION

Since a , d , and s are given, and n , but not l , is required, n may be found by eliminating l from (I) and (II) and solving the resulting equation.

$$\text{From (I) and (II), } l = a + (n - 1)d = \frac{2s - an}{n},$$

Substituting 2 for a , 3 for d , and 610 for s , and solving,

$$n = 20.$$

3. How many terms are there in the series 2, 6, 10, ... 66?
4. What is the sum of the series 1, 6, 11, ... 61?
5. How many terms are there in the series $-1, 2, 5, \dots$, if the sum is 221?
6. Determine the series 2, 9, 16, ... 86.
7. Determine the series $-10, -8\frac{1}{2}, -7, \dots$ to 10 terms.
8. The sum of the series ... 22, 27, 32, ... is 714. If there are 17 terms, what are the first and last terms?
9. If $s = 113\frac{3}{8}$, $a = \frac{1}{8}$, and $d = 2$, find n .
10. What is the sum of the series $-16, -11, -6, \dots 34$?
11. What is the sum of the series $\dots -1, 3, 7, \dots 23$, if the number of terms is 16?
12. What are the extremes of the series ... 7, 9, 11, ..., if $s = 300$, and $n = 20$?
13. How many terms are there in the series 1, 5, 9, ... l ?
14. What is the sum of an A.P. whose extremes are x and y , if the number of terms is b ?

363. To insert arithmetical means.

EXAMPLES

1. Insert 5 arithmetical means between 1 and 31.

SOLUTION. — Since there are 5 means, there must be 7 terms. Hence, in $l = a + (n - 1)d$, $l = 31$, $a = 1$, $n = 7$, and d is unknown.

Solving, $d = 5$.

Or, since there are 5 means, there must be 6 terms after the first.

Hence,
$$d = \frac{31 - 1}{6} = 5.$$

$\therefore 1, 6, 11, 16, 21, 26, 31$, is the series.

2. Insert 9 arithmetical means between 1 and 6.
3. Insert 10 arithmetical means between 24 and 2.
4. Insert 7 arithmetical means between 10 and -14 .
5. Insert 6 arithmetical means between -1 and 2.

6. Insert 14 arithmetical means between 15 and 20.
 7. Insert 3 arithmetical means between $a - b$ and $a + b$.
 8. Deduce the formula for the common difference when m arithmetical means are to be inserted between a and l . Find the first mean.
 9. What is the arithmetical mean between 2 and 6? between 10 and 20? between -3 and 5 ? between a and b ?

364. PRINCIPLE. — *The arithmetical mean between two numbers is equal to half their sum.*

The above principle may be established as follows:

Let a and b represent any two numbers, and A their arithmetical mean.

It is to be proved that $A = \frac{a+b}{2}$.

Since the two numbers and their arithmetical mean form the arithmetical progression a, A, b ,

§ 356,

$$A - a = b - A,$$

$$2A = a + b.$$

$$\therefore A = \frac{a+b}{2}.$$

EXAMPLES

Find the arithmetical mean between

- | | |
|--------------------------------------|--|
| 1. $\frac{2}{3}$ and $\frac{1}{3}$. | 4. $\frac{x+y}{x-y}$ and $\frac{x-y}{x+y}$. |
| 2. $a + b$ and $a - b$. | 5. $1 - x$ and $\frac{(1-x)^2}{1+x}$. |
| 3. $(a+b)^2$ and $(a-b)^2$. | |

PROBLEMS

365. Problems in Arithmetical Progression involving two unknown elements commonly suggest series of the form

$$x, x + y, x + 2y, x + 3y, \text{ etc.}$$

Frequently, however, the solution of problems is more readily accomplished by representing the series as follows:

1. When there are *three* terms, the series may be written

$$x - y, x, x + y.$$

2. When there are *five* terms, the series may be written,

$$x - 2y, x - y, x, x + y, x + 2y.$$

3. When there are *four* terms, the series may be written,

$$x - 3y, x - y, x + y, x + 3y.$$

The sum of the terms of a series represented as above evidently contains but one unknown number.

1. The sum of three numbers in arithmetical progression is 30, and the sum of their squares is 462. What are the numbers?

SOLUTION

Let the series be $x - y, x, x + y.$

Then, $(x - y) + x + (x + y) = 30,$ (1)

and $(x - y)^2 + x^2 + (x + y)^2 = 462.$ (2)

From (1), $3x = 30.$ (3)

$\therefore x = 10.$ (4)

From (2), $3x^2 + 2y^2 = 462.$ (5)

Substituting 10 for $x,$ $2y^2 = 162.$ (6)

Solving, $y = \pm 9.$

Forming the series from $x = 10$ and $y = \pm 9$, the terms are

$$1, 10, 19, \text{ or } 19, 10, 1.$$

2. The sum of three numbers in arithmetical progression is 18, and their product is 120. What are the numbers?

3. The sum of three numbers in arithmetical progression is 21, and the sum of their squares is 155. What are the numbers?

4. There are three numbers in arithmetical progression the sum of whose squares is 93. If the third is 4 times as large as the first, what are the numbers?

5. The product of the extremes of an arithmetical progression of 3 terms is 4 less than the square of the mean. What are the numbers, if their sum is 24?

6. The sum of four numbers in arithmetical progression is 14, and the product of the means is 12. What are the numbers?

7. The sum of seven numbers in arithmetical progression is 98, and the sum of their squares is 1484. What are the numbers?

8. The sum of five numbers in arithmetical progression is 15, and the product of the extremes is 3 less than the product of the terms next to the extremes. What are the numbers?

9. A number is expressed by three digits in arithmetical progression. If the number is divided by the sum of its digits, the quotient is $20\frac{1}{3}$; and if the number is increased by 594, the result is the number with its digits in the reverse order. What is the number?

10. Find the sum of the odd numbers from 1 to 100.

11. The product of the extremes of an arithmetical progression of 10 terms is 70, and the sum of the series is 95. What are the extremes?

12. Fifty-five logs are to be piled so that the top layer shall consist of 1 log, the next layer of 2 logs, the next layer of 3 logs, etc. How many logs must be placed in the bottom layer?

13. It cost Mr. Smith \$19.00 to have a well dug. If the cost of digging was \$1.50 for the first yard, \$1.75 for the second, \$2.00 for the third, etc., how deep was the well?

14. The product of the extremes of an arithmetical progression of 15 terms is 93, and the sum of the first and last means is 34. What is the progression?

15. How many arithmetical means must be inserted between 5 and 37, so that the ratio of the first mean to the last mean may be $\frac{1}{11}$?

16. How many arithmetical means must be inserted between 4 and 25, so that the sum of the series may be 116?

17. Prove that the equimultiples of the terms of an arithmetical progression are in arithmetical progression.

18. Prove that the difference of the squares of consecutive integers are in arithmetical progression, and that the common difference is 2.

19. Prove that the sum of n consecutive odd integers, beginning with 1, is n^2 .

GEOMETRICAL PROGRESSION

366. A series of numbers each of which after the first is derived by multiplying the preceding number by some constant multiplier is called a **Geometrical Series**, or a **Geometrical Progression**.

2, 4, 8, 16, 32 and a^4, a^3, a^2, a are geometrical progressions.

In the first series the constant multiplier is 2; in the second it is $\frac{1}{a}$.

G. P. is an abbreviation of the words *Geometrical Progression*.

367. The constant multiplier is called the **Ratio**.

It is evident that the terms of a geometrical progression increase or decrease numerically according as the ratio is numerically greater or less than 1.

368. To find the n th, or last term.

1. In the geometrical progression 3, 6, 12, 24, what is the ratio of 6 to 3? of 12 to 6? of 24 to 12?

2. In the geometrical progression a, ar, ar^2, ar^3, \dots what is the ratio? How many times does the ratio enter as a factor into the second term? into the third term? into the fourth term?

369. When a represents the first term of a G. P., r the ratio, and l the last or n th term,

$$l = ar^{n-1}. \quad (I)$$

EXAMPLES

1. Find the 9th term of the series 1, 3, 9, ...

PROCESS

$$l = ar^{n-1}$$

$$l = 1 \times 3^8$$

$$l = 6561$$

EXPLANATION. — In this example $a = 1, r = 3$, and $n = 9$.

Substituting these values in the formula for l , the last term is 6561.

2. Find the 10th term of the series 1, 2, 4, ...

3. Find the 8th term of the series $\frac{1}{4}, \frac{1}{2}, 1, \dots$

4. Find the 9th term of the series 6, 12, 24, ...

5. Find the 11th term of the series $\frac{1}{2}, 1, 2, \dots$.
6. Find the 7th term of the series $2, 6, 18, \dots$.
7. Find the 6th term of the series $4, 20, 100, \dots$.
8. Find the 6th term of the series $6, 18, 54, \dots$.
9. Find the 10th term of the series $1, \frac{1}{2}, \frac{1}{4}, \dots$.
10. Find the 10th term of the series $1, \frac{2}{3}, \frac{4}{9}, \dots$.
11. Find the 8th term of the series $\frac{1}{2}, \frac{1}{8}, \frac{2}{9}, \dots$.
12. Find the 11th term of the series $a^{19}b, a^{18}b^2, \dots$.
13. Find the n th term of the series $2, \sqrt{2}, 1, \dots$.
14. A man worked for 25 cents the first day, 50 cents the second day, \$1 the third day, and so on for 10 days. How much did he receive the tenth day?
15. If a man begins business with a capital of \$200 and doubles it every year for 6 years, how much will he have at the end of the sixth year?
16. If the population of the United States is 76 millions in 1900 and doubles itself every 25 years, what will it be in the year 2000?
17. A man's salary was raised $\frac{1}{4}$ every year for 5 years. If his salary was \$512 the first year, what was it the fifth year?
18. The population of a city at a certain time was 20,736, and increased in geometrical progression 25% each decade. What was the population at the end of 40 years?
19. A man who wanted 10 bushels of wheat thought \$1 a bushel too high a price. But he agreed to pay 2 cents for the first bushel, 6 cents for the second, 18 cents for the third, and so on. What did the last bushel cost him?
20. From a grain of corn there grew a stalk that produced an ear of 150 grains. These grains were planted, and each produced an ear of 150 grains. This process was repeated until there were 4 harvestings. If 75 ears of corn make 1 bushel, how many bushels were there the fourth year?

370. To find the sum of a finite series.

Let a represent the first term, l the n th term, or the last term, r the ratio, n the number of terms, and s the sum of the terms.

$$\text{Then,} \quad s = a + ar + ar^2 + ar^3 + \cdots + ar^{n-1}. \quad (1)$$

$$(1) \times r, \quad rs = ar + ar^2 + ar^3 + \cdots + ar^{n-1} + ar^n. \quad (2)$$

$$(2) - (1), \quad s(r - 1) = ar^n - a.$$

$$s = \frac{ar^n - a}{r - 1}, \text{ or } \frac{a(r^n - 1)}{r - 1}. \quad (\text{II})$$

But, since $ar^{n-1} = l$, $ar^n = rl$.

Substituting rl for ar^n in (II),

$$s = \frac{rl - a}{r - 1}, \text{ or } \frac{a - rl}{1 - r}. \quad (\text{III})$$

EXAMPLES

1. Find the sum of 6 terms of the series 3, 9, 27, ...

PROCESS

$$s = \frac{ar^n - a}{r - 1}$$

$$s = \frac{3 \times 3^6 - 3}{3 - 1} = 1092$$

EXPLANATION. — Since the first term a , the ratio r , and the number of terms n , are given, and formula II gives the sum in terms of a , r , and n , formula II is used.

2. Find the sum of 8 terms of the series 1, 2, 4, ...
3. Find the sum of 8 terms of the series 1, $\frac{1}{2}$, $\frac{1}{4}$, ...
4. Find the sum of 10 terms of the series 1, $1\frac{1}{2}$, $2\frac{1}{4}$, ...
5. Find the sum of 7 terms of the series 2, $-\frac{2}{3}$, $\frac{2}{9}$, ...
6. Find the sum of 12 terms of the series $-\frac{1}{2}$, $\frac{1}{4}$, $-\frac{1}{8}$, ...
7. Find the sum of 7 terms of the series 1, $2x$, $4x^2$, ...
8. Find the sum of 7 terms of the series 1, $-2x$, $4x^2$, ...
9. Find the sum of n terms of the series 1, x^2 , x^4 , ...
10. Find the sum of n terms of the series 1, 2, 4, ...

11. Find the sum of n terms of the series $1, \frac{1}{3}, \frac{1}{9}, \dots$.
12. The extremes of a geometrical series are 1 and 729, and the ratio is 3. What is the sum of the series?
13. What is the sum of the series $3, 6, 12, \dots, 192$?
14. What is the sum of the series $7, -14, 28, \dots, -224$?

371. To find the sum of an infinite geometrical series.

If the ratio r is numerically less than 1, it is evident that the successive terms of a geometrical series become numerically less and less. Hence, in an infinite decreasing geometrical series, the n th term l , or ar^{n-1} , can be made less than any assignable number, though not absolutely equal to zero.

$$(III) \text{ may be written } s = \frac{a}{1-r} - \frac{rl}{1-r}.$$

Since, by taking enough terms, l and consequently rl can be made less than any assignable number, the second fraction may be neglected.

Hence, the formula for the sum of an infinite decreasing geometrical series is

$$s = \frac{a}{1-r}. \quad (IV)$$

EXAMPLES

1. Find the sum of the series $1, \frac{1}{10}, \frac{1}{100}, \dots$.

SOLUTION

Substituting 1 for a and $\frac{1}{10}$ for r in (IV),

$$s = \frac{1}{1 - \frac{1}{10}} = \frac{1}{\frac{9}{10}} = \frac{10}{9}.$$

2. Find the value of .185185185 ...

SOLUTION

Since .185185185 ... = .185 + .000185 + .000000185 + ..., $a = .185$ and $r = .001$.

Substituting in (IV), .185185185 ... = $s = \frac{.185}{1 - .001} = \frac{5}{27}$.

Find the value of

3. $1 + \frac{1}{2} + \frac{1}{4} + \dots$

6. $.407407 \dots$

4. $3 + \frac{3}{4} + \frac{3}{16} + \dots$

7. $.363636 \dots$

5. $1 - \frac{1}{3} + \frac{1}{9} - \dots$

8. $1.94444 \dots$

372. To insert geometrical means between two terms.

EXAMPLES

1. Insert 3 geometrical means between 2 and 162.

PROCESS EXPLANATION. — Since there are three means, there are
 $l = ar^{n-1}$ five terms, and $n - 1 = 4$. Solving for r and neglecting
 162 = $2r^4$ imaginary values, $r = \pm 3$.
 Therefore, the series is either 2, 6, 18, 54, 162, or 2, -6,
 $r = \pm 3$ 18, -54, 162.

2. Insert 3 geometrical means between 1 and 625.

3. Insert 5 geometrical means between $4\frac{1}{2}$ and $204\frac{1}{2}$.

4. Insert 4 geometrical means between $\frac{343}{16}$ and $\frac{64}{16}$.

5. Insert 4 geometrical means between 5120 and 5.

6. Insert 4 geometrical means between $4\sqrt{2}$ and 1.

7. Insert 5 geometrical means between a^6 and b^6 .

8. Insert 6 geometrical means between -2 and $\frac{1}{8}\sqrt{2}$.

9. Insert 4 geometrical means between x and $-y$.

373. PRINCIPLE. — *The geometrical mean between two numbers is equal to the square root of their product.*

The above principle may be established as follows:

Let a and b represent any two numbers, and G their geometrical mean.

It is to be proved that $G = \sqrt{ab}$.

Since the two numbers and their geometrical mean form the geometrical progression a, G, b ,

§ 366.

$$\frac{G}{a} = \frac{b}{G},$$

$$G^2 = ab.$$

$$\therefore G = \sqrt{ab}.$$

Find the geometrical mean between

1. 8 and 50.
2. $\frac{1}{2}$ and $3\frac{5}{8}$.
3. $1\frac{1}{8}$ and $\frac{3}{4}$.
4. $(a+b)^2$ and $(a-b)^2$.
5. $\frac{a^2+ab}{a^2-ab}$ and $\frac{ab+b^2}{ab-b^2}$.
6. $25x^2 - 10x + 1$ and $x^2 + 10x + 25$.

374. Since formulæ I and II, or III, which is equivalent to II, are two independent simultaneous equations containing five elements, if *three* elements are *known*, the other *two* may be found by elimination.

PROBLEMS

- 375.** 1. Given r , l , and s , to find a .
2. The ratio of a geometrical progression is 5, the last term is 625, and the sum is 775. What is the first term?
3. The ratio of a geometrical progression is $\frac{1}{10}$, the sum is $\frac{1}{3}$, and the series is infinite. What is the first term?
4. Find l in terms of a , r , and s .
5. Find the last term of the series 5, 10, 20, ..., the sum of whose terms is 155.
6. If $\frac{1}{8} + \frac{1}{8}\sqrt{2} + \frac{1}{4} + \dots = 1\frac{7}{8}(1 + \sqrt{2})$, what is the last term, and the number of terms?
7. Deduce the formula for r in terms of a , l , and s .
8. If the sum of the geometrical progression 32 ... 243 is 665, what is the ratio? Write the series.
9. The sum of a geometrical progression is 700 greater than the first term and 525 greater than the last term. What is the ratio? If the first term is 81, what is the progression?
10. Deduce the formula for r in terms of a , n , and l .
11. The first term of a geometrical progression is 3, the last term is 729, and the number of terms is 6. What is the ratio? Write the series.
12. Find l in terms of r , n , and s .

13. The sum of the 12 terms of a geometrical progression whose ratio is 2 is 4095. What is the 12th term?

14. The velocity of a sled at the bottom of a hill is 100 feet per second. How far will it go on the level, if its velocity decreases each second $\frac{1}{2}$ of that of the previous second?

15. From a cask of vinegar $\frac{1}{3}$ was drawn off and the cask was filled by pouring in water. Show that if this is done 6 times, the contents of the cask will be more than $\frac{9}{16}$ water.

16. A ball thrown vertically into the air 100 feet falls and rebounds 40 feet the first time, 16 feet the second time, and so on. What is the whole distance through which the ball will have passed when it finally comes to rest?

17. Show that the amount of \$1 for 1, 2, 3, 4, 5 years at compound interest varies in geometrical progression.

18. Show that equimultiples of numbers in geometrical progression are also in geometrical progression.

19. The sum of three numbers in geometrical progression is 19, and the sum of their squares is 133. What are the numbers?

SUGGESTION. — When there are but three terms in the series they may be represented by x^2 , xy , y^2 , or by x , \sqrt{xy} , y .

20. The product of three numbers in geometrical progression is 8, and the sum of their squares is 21. What are the three numbers?

21. If 4 is a geometrical mean between two numbers whose sum is 10, what are the numbers?

22. The product of three numbers in geometrical progression is 64, and the sum of their cubes is 584. What are the numbers?

23. The sum of the first and second of four numbers in geometrical progression is 15, and the sum of the third and fourth is 60. What are the numbers?

SUGGESTION. — Four unknown numbers in geometrical progression may be represented by $\frac{x^2}{y}$, x , y , $\frac{y^2}{x}$.

24. The sum of the first and third of three numbers in geometrical progression is 130, and their product is 625. What are the numbers?

25. Divide \$700 among three persons so that the first shall receive \$300 more than the third, and the share of the second shall be a geometrical mean between the shares of the first and third.

26. If a , b , and c are in geometrical progression, show that their reciprocals also are in geometrical progression.

27. The difference between two numbers is 24, and their arithmetical mean exceeds their geometrical mean by 6. What are the numbers?

HARMONICAL PROGRESSION

376. 1. Examine the series $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$. Has it a constant difference? Has it a constant ratio?

2. Take the reciprocal of each term. What kind of a series is thus formed? How, then, may the series $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$, be described?

377. A series the reciprocals of whose terms form an arithmetical progression is called a **Harmonical Series**, or a **Harmonical Progression**.

$3, \frac{3}{2}, 1, \frac{2}{3}, \frac{1}{2}, \dots$ is a harmonical progression, because $\frac{1}{3}, \frac{2}{3}, 1, \frac{4}{3}, \frac{5}{3}, 2, \dots$ the reciprocals of its terms form an arithmetical progression.

H. P. is an abbreviation for the words *Harmonical Progression*.

378. Problems in harmonical progression are commonly solved by taking the reciprocals of the terms and employing the principles of arithmetical progression. There is no general method, however, for finding the sum of the terms of a harmonical progression.

379. PRINCIPLE 1. — *The harmonical mean between two numbers is equal to twice their product divided by their sum.*

The above principle may be established as follows:

Let H represent the harmonical mean between a and b .

It is to be proved that $H = \frac{2ab}{a+b}$.

§ 377, $\frac{1}{a}, \frac{1}{H}, \frac{1}{b}$ are in arithmetical progression.

Hence, § 356,
$$\frac{1}{b} - \frac{1}{H} = \frac{1}{H} - \frac{1}{a}.$$

Clearing of fractions,
$$aH - ab = ab - bH.$$

Transposing,
$$aH + bH = 2ab.$$

$$\therefore H = \frac{2ab}{a+b}.$$

380. PRINCIPLE 2. — *The geometrical mean between two numbers is also the geometrical mean between their arithmetical and harmonical means.*

The above principle may be established as follows :

§ 364,
$$A = \frac{a+b}{2}. \quad (1)$$

§ 373,
$$G = \sqrt{ab}. \quad (2)$$

§ 379,
$$H = \frac{2ab}{a+b}. \quad (3)$$

Multiplying (1) by (3),
$$AH = ab. \quad (4)$$

Taking the square root,
$$\sqrt{AH} = \sqrt{ab}. \quad (5)$$

From (2) and (5), Ax. 1,
$$G = \sqrt{AH}.$$

Hence, § 373, G is the geometrical mean between A and H .

EXAMPLES

1. Find the 12th term of the H. P. 6, 3, 2,

SOLUTION. — The reciprocals of the terms form the arithmetical progression

$$\frac{1}{6}, \frac{1}{3}, \frac{1}{2}, \dots$$

in which $a = \frac{1}{6}$ and $d = \frac{1}{6}$.

Substituting $\frac{1}{6}$ for a , $\frac{1}{6}$ for d , and 12 for n in (I),

§ 359,
$$l = \frac{1}{6} + (12-1)\frac{1}{6} = 2.$$

Therefore, § 377, the 12th term of the given harmonical progression is $\frac{1}{2}$.

2. Find the 10th term of the harmonical series $\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots$.

3. Insert 6 harmonical means between $1\frac{1}{2}$ and 12.

4. Insert 2 harmonical means between 2 and 5.

5. Insert 7 harmonical means between $12\frac{1}{2}$ and $2\frac{1}{2}$.

6. Insert 3 harmonical means between b and a .
7. Find the n th term of the H. P. $\frac{1}{5}, \frac{1}{8}, \frac{1}{11}, \dots$.
8. The 3d and 4th terms of a H. P. are $2\frac{1}{2}$ and $1\frac{1}{2}$. Write the first 6 terms.

Find the harmonical mean between

9. 2 and 3.
10. $\frac{1}{2}$ and $\frac{1}{3}$.
11. $2\frac{1}{2}$ and $1\frac{1}{2}$.
12. $2\frac{1}{2}$ and 10.
13. $a - c$ and $a + c$.
14. $1 - \sqrt{a}$ and $1 + \sqrt{a}$.
15. a and $\frac{1}{a}$.
16. $\sqrt{6}$ and $\sqrt{3}$.
17. The 5th term of a harmonical progression is $\frac{1}{12}$, and the 11th term is $\frac{1}{24}$. What is the first term?
18. The arithmetical mean between two numbers is 5, and their harmonical mean is $3\frac{1}{2}$. What are the numbers?
19. If one number exceeds another by 2, and their arithmetical mean exceeds their harmonical mean by $\frac{1}{3}$, what are the numbers?
20. If a , b , and c are in harmonical progression, prove that $a - b : b - c = a : c$.
21. If a , b , c , and d are in harmonical progression, prove that $ab : cd = b - a : d - c$.
22. If b is the harmonical mean between a and c , prove that

$$\frac{1}{b-a} + \frac{1}{b-c} = \frac{1}{a} + \frac{1}{c}.$$

23. When $b - a : c - b = a : x$, prove that $x = a$, if a , b , and c are in arithmetical progression; that $x = b$, if a , b , and c are in geometrical progression; and that $x = c$, if a , b , and c are in harmonical progression.

24. The harmonical mean between two numbers is $5\frac{7}{8}$ and their arithmetical mean is $6\frac{1}{2}$. What is their geometrical mean?

25. Prove that $x + xy$, $2xy$, and $xy + xy^2$ are in harmonical progression.

26. If $b + c$, $c + a$, and $a + b$ are in harmonical progression, prove that a^2 , b^2 , c^2 are in arithmetical progression.

IMAGINARY AND COMPLEX NUMBERS

381. 1. If from $\sqrt{-25}$ the rational factor $\sqrt{25}$ is removed, what irrational factor remains?

2. Simplify $\sqrt{-25}$, $\sqrt{-16}$, $\sqrt{-a^2}$. What common part, or *unit*, have the indicated square roots of negative numbers?

3. What is the square of $\sqrt{4}$? of $\sqrt{5}$? of $\sqrt{9}$? of $\sqrt{2x}$?

What is the effect of squaring a radical of the second degree? What, then, is the square of $\sqrt{-4}$? of $\sqrt{-a}$? of $\sqrt{-1}$?

382. Up to this point the only numbers whose nature has been discussed have been numbers that differ from arithmetical numbers in having a sign, + or -, to indicate quality or direction. These numbers are called real numbers, and may be briefly described as *numbers whose squares are positive*.

There are numbers, however, *whose squares are negative*. They constitute one class of imaginary numbers, defined in § 257. In this chapter only imaginary numbers of the second degree are treated.

Let $-a$ be any negative real number.

Then, $\sqrt{-a}$ will represent an imaginary number.

Since $+\sqrt{-a} = +\sqrt{a} \cdot \sqrt{-1} = +\sqrt{-1} \cdot \sqrt{a}$

and $-\sqrt{-a} = -\sqrt{a} \cdot \sqrt{-1} = -\sqrt{-1} \cdot \sqrt{a}$,

the positive imaginary unit is $+\sqrt{-1}$, and the negative imaginary unit is $-\sqrt{-1}$.


Since the square of the square root of a number is the number itself,

$$(\sqrt{-1})^2 = -1.$$

This relation is sufficient to explain operations with imaginary numbers.

383. Relation between the units $+1$, -1 , $\sqrt{-1}$, and $-\sqrt{-1}$.

In the accompanying figure the *length* of any radius of the circle represents the arithmetical unit 1. The line drawn from O to A , called the line OA ,



represents the positive unit $+1$, and the line OA'' represents the negative unit -1 . Every real number lies somewhere on the line $X'X$, which is supposed to extend indefinitely in both directions from O . $X'X$ is called the axis of real numbers.

The *direction* of any line drawn from O , as OB , that is, the quality or *direction sign* of the number represented by that line, is determined with reference to the fixed

line OA by finding what part of a revolution is required to swing the line from the position OA to the required position. By common consent revolution of the line OA is performed in a direction opposite to that of the hands of a clock, as shown by the arrows. OB is reached after $\frac{1}{4}$ of a revolution, OA' after $\frac{1}{2}$ of a revolution, OA'' after $\frac{3}{4}$ of a revolution, etc.

Since OA'' , or -1 , represents $\frac{1}{2}$ of a revolution of OA , the square of OA'' , or $(-1)^2$, represents 1 revolution of OA , which produces OA , or $+1$. Hence, OA'' , or -1 , represents the square root of $+1$, or $(+1)^{\frac{1}{2}}$.

Similarly, since OA' represents $\frac{1}{4}$ of $\frac{1}{2}$ of a revolution of OA , and OA'' represents $\frac{1}{2}$ of a revolution of OA , OA' represents the square root of OA'' , or of -1 ; that is, $OA' = \sqrt{-1}$.

If OA'' is swung $\frac{1}{4}$ of a revolution from the position OA'' to the position OA''' , OA''' will be multiplied by $\sqrt{-1}$ just as OA is multiplied by $\sqrt{-1}$ to produce OA' . Hence, the result $OA''' = -1 \cdot \sqrt{-1} = -\sqrt{-1}$.

$+1$, represented by OA , and -1 , represented by OA'' , are the units for real numbers, that is, are *real units*. Just as the real number $+a$ is represented by a line a units long extending from O toward X , and the real number $-a$ by a line a units long extending from O in the opposite direction, so the imaginary number $+a\sqrt{-1}$, or $(+\sqrt{-1}) \times a$, is represented by a line a units long extending from O toward Y , and the imaginary number $-a\sqrt{-1}$, or $(-\sqrt{-1}) \times a$, by a line a units long extending from O in the opposite direction, toward Y' . Hence, $+\sqrt{-1}$ and $-\sqrt{-1}$ are the units for imaginary numbers, that is, they are *imaginary units*; $+a\sqrt{-1}$ is called a *positive imaginary number* and $-a\sqrt{-1}$ a *negative imaginary number*.

$Y'Y$ is called the axis of imaginary numbers. If $Y'Y$ is taken as the axis of real numbers, then $X'X$ becomes the axis of imaginary numbers. Hence, it is seen that imaginary numbers have as much reality as real numbers. Imaginary numbers were named before their nature was understood.

384. In the graphical illustration of the relation between real and imaginary numbers it was assumed that $+1 \cdot \sqrt{-1} = \sqrt{-1}$, or $\sqrt{-1} \cdot 1$; that is, the Commutative Law for multiplication was assumed to apply when imaginary numbers were involved. It is evident, if the discussion of number is to proceed, that in any operation imaginary numbers must obey all the laws of real numbers except those which determine the quality of the result; and that the quality of the result is determined, as far as the imaginary numbers are concerned, by the relation $(\sqrt{-1})^2 = -1$.

385. Powers of $\sqrt{-1}$.

$$(\sqrt{-1}) = +\sqrt{-1};$$

$$(\sqrt{-1})^2 = -1;$$

$$(\sqrt{-1})^3 = (\sqrt{-1})^2 \sqrt{-1} = (-1)\sqrt{-1} = -\sqrt{-1};$$

$$(\sqrt{-1})^4 = (\sqrt{-1})^2(\sqrt{-1})^2 = (-1)(-1) = +1;$$

$$(\sqrt{-1})^5 = (\sqrt{-1})^4 \sqrt{-1} = (+1)\sqrt{-1} = +\sqrt{-1};$$

and so on. Hence, if $n = 0$ or a positive integer,

$$(\sqrt{-1})^{4n+1} = +\sqrt{-1}; \quad (\sqrt{-1})^{4n+2} = -1;$$

$$(\sqrt{-1})^{4n+3} = -\sqrt{-1}; \quad (\sqrt{-1})^{4n+4} = +1.$$

Hence, *any even power of $\sqrt{-1}$ is real and any odd power is imaginary.*

386. Operations involving imaginary numbers.

EXAMPLES

Find the value of

1. $(\sqrt{-1})^6$ 3. $(\sqrt{-1})^{10}$ 5. $(\sqrt{-1})^{18}$ 7. $(-\sqrt{-1})^3$.

2. $(\sqrt{-1})^7$ 4. $(\sqrt{-1})^{21}$ 6. $(\sqrt{-1})^{15}$ 8. $(-\sqrt{-1})^8$.

9. Add $\sqrt{-a^4}$ and $\sqrt{-16a^4}$.

SOLUTION

$$\sqrt{-a^4} + \sqrt{-16a^4} = a^2\sqrt{-1} + 4a^2\sqrt{-1} = 5a^2\sqrt{-1}.$$

Simplify the following:

10. $\sqrt{-4} + \sqrt{-49}$. 13. $\sqrt{-12} + 4\sqrt{-3}$.
 11. $\sqrt{-9} + \sqrt{-64}$. 14. $5\sqrt{-18} - \sqrt{-72}$.
 12. $2\sqrt{-4} + 3\sqrt{-1}$. 15. $3\sqrt{-20} - \sqrt{-80}$.
 16. $(\sqrt{-a} + 3\sqrt{-b}) + (\sqrt{-a} - 3\sqrt{-b})$.
 17. $(\sqrt{-9xy} - \sqrt{-xy}) - (\sqrt{-4xy} + \sqrt{-xy})$.
 18. $\sqrt{-x^2} + \sqrt{-4x^2} - \sqrt{-x^2} + 3x\sqrt{-x}$.
 19. $\sqrt{-16} - 3\sqrt{-4} + \sqrt{-18} + \sqrt{-50} + \sqrt{-25}$.
 20. $\sqrt{-8} + a\sqrt{-2} - \sqrt{-98} - 5\sqrt{-2a^2}$.
 21. $\sqrt{-16a^2x^2} + \sqrt{-a^2x^2} - \sqrt{-9a^2x^2}$.
 22. $\sqrt{1-5} - 3\sqrt{1-10} + 2\sqrt{5-30}$.
 23. Multiply $3\sqrt{-10}$ by $2\sqrt{-8}$.

PROCESS

$$\begin{aligned} 3\sqrt{-10} \times 2\sqrt{-8} &= 3\sqrt{10}\sqrt{-1} \times 2\sqrt{8}\sqrt{-1} \\ &= 6\sqrt{10} \times 8 \times (-1) \\ &= -6\sqrt{80} = -24\sqrt{5} \end{aligned}$$

EXPLANATION. — In order to determine the sign of the product, each imaginary number is reduced to the form $b\sqrt{-1}$. The numbers are then multiplied together as ordinary radicals, observing, however, that $\sqrt{-1} \times \sqrt{-1} = -1$.

24. Multiply $\sqrt{-2} + 3\sqrt{-3}$ by $4\sqrt{-2} - \sqrt{-3}$.

FIRST SOLUTION

$$\begin{aligned} \sqrt{-2} + 3\sqrt{-3} &= (\sqrt{2} + 3\sqrt{3})\sqrt{-1} \\ 4\sqrt{-2} - \sqrt{-3} &= (4\sqrt{2} - \sqrt{3})\sqrt{-1} \\ \hline (\sqrt{2} + 3\sqrt{3})(4\sqrt{2} - \sqrt{3})(-1) \\ &= (8 + 12\sqrt{6} - \sqrt{6} - 9)(-1) \\ &= 1 - 11\sqrt{6} \end{aligned}$$

SECOND SOLUTION

$$\begin{aligned} \sqrt{-2} + 3\sqrt{-3} \\ 4\sqrt{-2} - \sqrt{-3} \\ \hline -4\sqrt{4} - 12\sqrt{6} \\ + 3\sqrt{9} + \sqrt{6} \\ \hline 1 - 11\sqrt{6} \end{aligned}$$

Multiply:

25. $3\sqrt{-5}$ by $2\sqrt{-15}$. 28. $8\sqrt{-1}$ by $\sqrt{-b^2}$.
 26. $4\sqrt{-27}$ by $\sqrt{-12}$. 29. $\sqrt{-125}$ by $\sqrt{-108}$.
 27. $2\sqrt{-8}$ by $5\sqrt{-3}$. 30. $\sqrt{-100}$ by $\sqrt{-30}$.

31. $\sqrt{-6} + \sqrt{-3}$ by $\sqrt{-6} - \sqrt{-3}$.
 32. $\sqrt{-ab} + \sqrt{-a}$ by $\sqrt{-ab} - \sqrt{-a}$.
 33. $\sqrt{-xy} + \sqrt{-x}$ by $\sqrt{-xy} - \sqrt{-x}$.
 34. $\sqrt{-50} - \sqrt{-12}$ by $\sqrt{-8} - \sqrt{-75}$.
 35. $\sqrt{-a} + \sqrt{-b} + \sqrt{-c}$ by $\sqrt{-a} + \sqrt{-b} - \sqrt{-c}$.
 36. Divide $\sqrt{-12}$ by $\sqrt{-3}$.

SOLUTION

$$\frac{\sqrt{-12}}{\sqrt{-3}} = \frac{\sqrt{12}\sqrt{-1}}{\sqrt{3}\sqrt{-1}} = \frac{\sqrt{12}}{\sqrt{3}} = \sqrt{4} = 2.$$

37. Divide $\sqrt{12}$ by $\sqrt{-3}$.

SOLUTION

$$\begin{aligned}\frac{\sqrt{12}}{\sqrt{-3}} &= \frac{\sqrt{12}}{\sqrt{3}\sqrt{-1}} = \frac{\sqrt{4}}{\sqrt{-1}} = \frac{2}{\sqrt{-1}} \\ &= \frac{2\sqrt{-1}}{-1} = -2\sqrt{-1}.\end{aligned}$$

38. Divide 5 by $(\sqrt{-1})^3$.

SOLUTION

$$\frac{5}{(\sqrt{-1})^3} = \frac{5(+1)}{(\sqrt{-1})^3} = \frac{5(\sqrt{-1})^4}{(\sqrt{-1})^3} = 5\sqrt{-1}.$$

Divide:

- | | |
|--|---|
| 39. $\sqrt{-18}$ by $\sqrt{-3}$. | 48. -2 by $\sqrt{-1}$. |
| 40. $\sqrt{27}$ by $\sqrt{-3}$. | 49. $(\sqrt{-1})^5$ by $\frac{1}{3}\sqrt{-1}$. |
| 41. $14\sqrt{-5}$ by $2\sqrt{-7}$. | 50. $(\sqrt{-1})^3$ by $(\sqrt{-1})^{15}$. |
| 42. $-\sqrt{-a^2}$ by $\sqrt{-b^2}$. | 51. $\sqrt{4ab}$ by $\sqrt{-bc}$. |
| 43. 1 by $\sqrt{-1}$. | 52. $\sqrt{-20} - \sqrt{-2}$ by $2\sqrt{-1}$. |
| 44. $(\sqrt{-1})^4 - \sqrt{-1}$ by $\sqrt{-1}$. | 53. $\sqrt{-16} - \sqrt{-6}$ by $2\sqrt{-2}$. |
| 45. $\sqrt{-3} + (\sqrt{-1})^2$ by $\sqrt{-1}$. | 54. $(\sqrt{-1})^{14}$ by $-\frac{1}{2}\sqrt{-1}$. |
| 46. $\sqrt{8} - 3\sqrt{14}$ by $\sqrt{-2}$. | 55. $(\sqrt{-1})^{10}$ by $(\sqrt{-1})^{-2}$. |
| 47. $\sqrt{12} + \sqrt{3}$ by $\sqrt{-3}$. | 56. $\sqrt{-a^2} + b\sqrt{-1}$ by $\sqrt{-ab}$. |
| 57. $\sqrt{-4}$ by $\sqrt{-2} \cdot \sqrt{-2} \cdot \sqrt{-1}$. | |

395. Operations involving complex numbers.

EXAMPLES

1. Add
- $3 - 2\sqrt{-1}$
- and
- $2 + 5\sqrt{-1}$
- .

SOLUTION

$$\begin{aligned} 3 - 2\sqrt{-1} + 2 + 5\sqrt{-1} &= (3 + 2) + (-2\sqrt{-1} + 5\sqrt{-1}) \\ &= 5 + 3\sqrt{-1}. \end{aligned}$$

EXPLANATION. — Since, § 387, the imaginary terms cannot unite with the real terms, the simplest form of the sum is obtained by uniting the real and the imaginary terms separately and indicating the algebraic sum of the results.

Simplify the following:

2. $(5 + \sqrt{-4}) + (\sqrt{-9} - 3).$

3. $(2 - \sqrt{-16}) + (3 + \sqrt{-4}).$

4. $(3 - \sqrt{-8}) + (4 + \sqrt{-18}).$

5. $(\sqrt{-20} - \sqrt{16}) + (\sqrt{-45} + \sqrt{4}).$

6. $(4 + \sqrt{-25}) - (2 + \sqrt{-4}).$

7. $(3 - 2\sqrt{-5}) - (2 - 3\sqrt{-5}).$

8. $(2 - 2\sqrt{-1} + 3) - (\sqrt{16} - \sqrt{-16}).$

9. $\sqrt{-49} - 2 - 3\sqrt{-4} - \sqrt{-1} + 6.$

10. Expand
- $(a + b\sqrt{-1})(a + b\sqrt{-1}).$

SOLUTION

§ 91, $(a + b\sqrt{-1})(a + b\sqrt{-1}) = a^2 + 2ab\sqrt{-1} + (b\sqrt{-1})^2.$

§ 384, $= a^2 + 2ab\sqrt{-1} - b^2.$

11. Expand
- $(\sqrt{5} - \sqrt{-3})^2.$

SOLUTION

$$\begin{aligned} (\sqrt{5} - \sqrt{-3})^2 &= 5 - 2\sqrt{-15} + (-3) \\ &= (5 - 3) - 2\sqrt{-15} \\ &= 2 - 2\sqrt{-15}. \end{aligned}$$

Expand the following:

12. $(2 + 3\sqrt{-1})(1 + \sqrt{-1}).$

15. $(2 + 3i)^2.$

13. $(5 - \sqrt{-1})(1 - 2\sqrt{-1}).$

16. $(2 - 3i)^2.$

14. $(\sqrt{2} + \sqrt{-2})(\sqrt{8} - \sqrt{-8}).$

17. $(a - bi)^2.$

18. Show that $(1 + \sqrt{-3})(1 + \sqrt{-3})(1 + \sqrt{-3}) = -8$.
19. Show that $(-1 + \sqrt{-3})(-1 + \sqrt{-3})(-1 + \sqrt{-3}) = 8$.
20. Show that $(-\frac{1}{2} + \frac{1}{2}\sqrt{-3})(-\frac{1}{2} + \frac{1}{2}\sqrt{-3})(-\frac{1}{2} + \frac{1}{2}\sqrt{-3}) = 1$.
21. Divide $8 + \sqrt{-1}$ by $3 + 2\sqrt{-1}$.

FIRST SOLUTION

$$\begin{array}{r} 8 + \sqrt{-1} = 6 + \sqrt{-1} + 2 \quad \left| \begin{array}{l} 3 + 2\sqrt{-1} \\ 2 - \sqrt{-1} \end{array} \right. \\ \underline{6 + 4\sqrt{-1}} \\ -3\sqrt{-1} + 2 \\ \underline{-3\sqrt{-1} + 2} \end{array}$$

The real term of the dividend may always be separated into two parts, one of which will exactly contain the real term of the divisor.

SECOND SOLUTION

$$\frac{8 + \sqrt{-1}}{3 + 2\sqrt{-1}} = \frac{(8 + \sqrt{-1})(3 - 2\sqrt{-1})}{(3 + 2\sqrt{-1})(3 - 2\sqrt{-1})} = \frac{26 - 13\sqrt{-1}}{9 + 4} = 2 - \sqrt{-1}.$$

Divide:

22. 3 by $1 - \sqrt{-2}$. 26. $16 + 4\sqrt{-5}$ by $3 - \sqrt{-5}$.
23. 2 by $1 + \sqrt{-1}$. 27. $a^2 + b^2$ by $a - b\sqrt{-1}$.
24. $4 + \sqrt{4}$ by $2 - \sqrt{-2}$. 28. $a + bi$ by $ai + b$.
25. $9 + \sqrt{-3}$ by $3 - \sqrt{-3}$. 29. $(1 + i)^2$ by $1 - i$.
30. Find by inspection the square root of $3 + 2\sqrt{-10}$.

SOLUTION

$$\begin{aligned} 3 + 2\sqrt{-10} &= (5 - 2) + 2\sqrt{5 \cdot -2} = 5 + 2\sqrt{5 \cdot -2} + (-2). \\ \therefore \sqrt{3 + 2\sqrt{-10}} &= \sqrt{5 + 2\sqrt{5 \cdot -2} + (-2)} = \sqrt{5} + \sqrt{-2}. \end{aligned}$$

Find by inspection the square root of

31. $4 + 2\sqrt{-21}$. 33. $6 - 2\sqrt{-7}$. 35. $12\sqrt{-1} - 5$.
32. $1 + 2\sqrt{-6}$. 34. $9 + 2\sqrt{-22}$. 36. $b^2 + 2ab\sqrt{-1} - a^2$.
37. Verify that $-1 + \sqrt{-1}$ and $-1 - \sqrt{-1}$ are roots of the equation $x^2 + 2x + 2 = 0$.
38. Expand $(\frac{1}{2} + \frac{1}{2}\sqrt{-3})^3$.

INEQUALITIES

396. One number is said to be *greater* than another when the remainder obtained by subtracting the second from the first is *positive*, and to be *less* than another when the remainder obtained by subtracting the second from the first is *negative*.

If $a - b$ is a positive number, a is *greater* than b ; but if $a - b$ is a negative number, a is *less* than b .

Any negative number is regarded as less than 0; and, of two negative numbers, that more remote from 0 is the less.

Thus, -1 is less than 0; -2 is less than -1 ; -3 is less than -2 ; etc.

An algebraic expression indicating that one number is greater or less than another is called an **Inequality**.

397. The **Sign of Inequality** is $>$ or $<$.

It is placed between two unequal numbers with the opening toward the greater.

Thus, a is *greater than* b is written $a > b$; a is *less than* b is written $a < b$.

The expressions on the left and right, respectively, of the sign of inequality are termed the *first* and *second members* of the inequality.

398. When the first members of two inequalities are each greater or each less than the corresponding second members, the inequalities are said to *subsist in the same sense*.

When the first member is greater in one inequality and less in another, the inequalities are said to *subsist in a contrary sense*.

$x > a$ and $y > b$ subsist in the same sense, also $x < 3$ and $y < 4$; but $x > b$ and $y < a$ subsist in a contrary sense.

399. 1. If 2 is added to each member of the inequality $8 > 5$, how will the two inequalities subsist? How will they subsist, if 2 is subtracted from each member?

2. Investigate other inequalities.

PRINCIPLE 1. — *If the same number or equal numbers are added to or subtracted from both members of an inequality, the resulting inequality will subsist in the same sense.*

Let $a > b$.

Then, § 396, $a - b = p$, a positive number.

Adding $c - c = 0$, Ax. 2, $a + c - (b + c) = p$.

Therefore, $a + c > b + c$.

400. 1. What is the effect of adding 2 to each member of the inequality $x - 2 > y$? of subtracting 2 from each member of the inequality $a + 2 > b$?

2. If a term is transposed from one member of an inequality to the other, what must be done to its sign?

PRINCIPLE 2. — *A term may be transposed from one member of an inequality to the other, provided its sign is changed.*

Let $a - b > c - d$.

Adding b to each side, Prin. 1, $a > b + c - d$.

Adding $-c$ to each side, Prin. 1, $a - c > b - d$.

401. PRINCIPLE 3. — *If the signs of all the terms of an inequality are changed, the resulting inequality will subsist in a contrary sense.*

Let $a - b > c - d$.

Transposing every term, Prin. 2,

$$-c + d > -a + b;$$

that is,

$$-a + b < -c + d.$$

402. 1. If both members of the inequality $10 > 8$ are multiplied by 2, how will the two inequalities subsist? How will they subsist, if both members are divided by 2?

2. How will they subsist in each case, if the multiplier or divisor is -2 ?

PRINCIPLE 4. — *If both members of an inequality are multiplied or divided by the same number, the resulting inequality will subsist in the same sense if the multiplier or divisor is positive, but in the contrary sense if the multiplier or divisor is negative.*

Let $a > b$.

Then, § 396, $a - b = p$, a positive number.

Multiplying by m , $ma - mb = mp$.

If m is positive, mp is positive,

and $ma > mb$.

If m is negative, mp is negative,

and $mb > ma$, or $ma < mb$.

Putting $\frac{1}{m}$ for m , the principle holds also for division.

403. 1. If the corresponding members of the inequalities $6 > 5$ and $4 > 2$ are added, how will the resulting inequality subsist? How, if $-5 > -6$ and $-2 > -4$ are added?

PRINCIPLE 5. — *If the corresponding members of any number of inequalities subsisting in the same sense are added together, the resulting inequality will subsist in the same sense.*

Let $a > b$, $c > d$, $e > f$, etc.

Then, § 396, $a - b$, $c - d$, $e - f$, etc., are positive numbers.

Hence, their sum, $a + c + e + \dots - (b + d + f + \dots)$, is a positive number; that is,

$$a + c + e + \dots > b + d + f + \dots$$

404. **PRINCIPLE 6.** — *If each member of an inequality is subtracted from the corresponding members of an equation, the resulting inequality will subsist in a contrary sense.*

Let $a > b$ and let c be any number.

Then, § 396, $a - b$ is a positive number.

Since a number is diminished by subtracting a positive number from it,

$$c - (a - b) < c.$$

Transposing, $c - a < c - b$.

That is, if each member of the inequality $a > b$ is subtracted from the corresponding member of the equation $c = c$, the result is an inequality subsisting in a contrary sense.

405. It is evident that the difference of two inequalities subsisting in the same sense, or the sum of two inequalities subsisting in a contrary sense, or the product, or the quotient of two inequalities, member by member, may have its first member greater than, equal to, or less than its second.

For example, take the inequality $12 > 6$.

Subtracting $8 > 2$, or adding $-8 < -2$, the result is $4 = 4$.

Subtracting $8 > 1$, or adding $-8 < -1$, the result is $4 < 5$.

Multiplying and dividing by $3 > 2$, the results are $36 > 12$ and $4 > 3$.

Multiplying by $-2 > -4$, the result is $-24 = -24$. Dividing by $4 > 2$, the result is $3 = 3$.

Multiplying and dividing by $-2 > -3$, the results are $-24 < -18$ and $-6 < -2$.

EXAMPLES

406. 1. Find one limit of x in the inequality $3x - 10 > 11$.

SOLUTION

$$3x - 10 > 11.$$

$$\text{Prin. 2,} \quad 3x > 21.$$

$$\text{Prin. 4,} \quad x > 7.$$

Therefore, the *inferior* limit of x is 7; that is, x may have any value greater than 7.

2. Find the limits of x in the simultaneous inequalities $3x + 5 < 38$ and $4x < 7x - 18$.

SOLUTION

$$3x + 5 < 38. \quad (1)$$

$$4x < 7x - 18. \quad (2)$$

$$\text{Transposing in (1), Prin. 2,} \quad 3x < 33.$$

$$\therefore \text{Prin. 4,} \quad x < 11.$$

$$\text{Transposing in (2), Prin. 2,} \quad -3x < -18.$$

$$\text{Changing signs, Prin. 3,} \quad 3x > 18.$$

$$\therefore \text{Prin. 4,} \quad x > 6.$$

The *inferior* limit of x is 6, and the *superior* limit is 11; that is, the given inequalities are satisfied simultaneously by any value of x between 6 and 11.

3. Find the limits of x and y in $3x - y > -14$ and $x + 2y = 0$.

SOLUTION

$$3x - y > -14. \quad (1)$$

$$x + 2y = 0. \quad (2)$$

Multiplying (1) by 2, Prin. 4,

$$6x - 2y > -28. \quad (3)$$

Adding (2) and (3),

$$7x > -28.$$

$$\therefore x > -4. \quad (4)$$

Multiplying (2) by 3,

$$3x + 6y = 0. \quad (5)$$

Subtracting (5) from (1), Prin. 1,

$$-7y > -14.$$

Dividing by -7 , Prin. 4,

$$y < 2.$$

That is, x is greater than -4 , and y is less than 2 .

Find the limits of x in the following:

4. $6x - 5 > 13.$

5. $5x - 1 < 6x + 4.$

6. $3x - \frac{1}{2}x < 30.$

7. $4x + 1 < 6x - 11.$

8. $\begin{cases} 4x - 11 > \frac{1}{3}x, \\ 20 - 2x > 10. \end{cases}$

9. $\begin{cases} 3 - 4x < 7, \\ 5x + 10 < 20. \end{cases}$

10. $x + \frac{2x}{3} + \frac{5x}{6} > 25 \text{ and } < 30.$

11. Find the limits of x in $x^2 + 3x > 28$.

SOLUTION

$$x^2 + 3x > 28.$$

Transposing, Prin. 2,

$$x^2 + 3x - 28 > 0.$$

Factoring,

$$(x - 4)(x + 7) > 0.$$

That is, $(x - 4)(x + 7)$ is positive.

If $(x - 4)(x + 7)$ is positive, either both factors are positive or both are negative. Both factors are positive, when $x > 4$; both factors are negative, when $x < -7$.

Therefore, x can have all values except 4 and -7 and intermediate values.

Find the limits of x in the following:

12. $x^2 + 3x > 10$.

16. $x^2 > 9x - 18$.

13. $x^2 + 8x > 20$.

17. $x^2 + 40x > 3(4x - 25)$.

14. $x^2 + 5x > 24$.

18. $x^2 + bx > ax + ab$.

15. $(x-2)(3-x) > 0$.

19. $(x-3)(5-x) > 0$.

Find the limits of x and y in the following, and, if possible, one positive integral value for each unknown number:

20.
$$\begin{cases} 2x - 3y < 2, \\ 2x + 5y = 25. \end{cases}$$

23.
$$\begin{cases} y = 3x + 4, \\ 25 < 2y + 3x. \end{cases}$$

21.
$$\begin{cases} 3x + 2y = 42, \\ 3x - \frac{y}{7} > 16. \end{cases}$$

24.
$$\begin{cases} y - x > 9, \\ \frac{7x}{20} + \frac{y}{15} = 9. \end{cases}$$

22.
$$\begin{cases} x + y = 10, \\ 4x < 3y. \end{cases}$$

25.
$$\begin{cases} x > y + 4, \\ x - 2y = 8. \end{cases}$$

If a , b , and c are positive and unequal,

26. Which is the greater, $\frac{a+b}{a+2b}$ or $\frac{a+2b}{a+3b}$?

27. Prove that $a^2 + b^2 > 2ab$.

SUGGESTION. — Whether $a - b$ is positive or negative, $(a - b)^2$ is positive.

28. Prove that $a^2 + b^2 + c^2 > ab + ac + bc$.

29. Prove that $a^3 + b^3 > a^2b + ab^2$.

30. Which is the greater, $\frac{a^2+b^2}{a+b}$ or $\frac{a^3+b^3}{a^2+b^2}$?

31. Prove that $\frac{a}{3b} + \frac{3b}{4a} > 1$, except when $2a = 3b$.

32. Prove that $(a-2b)(4b-a) < b^2$, except when $a = 3b$.

33. Prove that $a^3 + b^3 + c^3 > 3abc$.

34. Prove that the sum of any positive real number, except 1, and its reciprocal is always greater than 2.

VARIABLES AND LIMITS

407. A number that has the same value throughout a discussion is called a **Constant**.

Arithmetical numbers are constants. A literal number, as a or x , is constant in a discussion, if it keeps the same value throughout that discussion.

408. A number that under the conditions imposed upon it may have a series of different values is called a **Variable**.

Variables are usually represented by x , y , z , etc.

The numbers .3, .33, .333, .3333, . . . are successive values of a variable approaching the constant $\frac{1}{3}$.

The commensurable numbers 1, 1.4, 1.41, 1.414, 1.4142, . . . are successive values of a variable approaching the constant $\sqrt{2}$.

409. An expression whose value depends upon the value of a variable is called a **Function** of that variable.

$2x + 1$ is a function of x , because, if successive values are given to x , $2x + 1$ will take successive values. For example, if $x = 0$, $2x + 1 = 1$; if $x = 1$, $2x + 1 = 3$; if $x = 2$, $2x + 1 = 5$, etc. $x^2 - 2x$ and $\frac{1}{1-x}$ are also functions of x .

$x^2 + 2xz - 5z^2$ is a function of x and z .

Every function of a single variable is a variable.

The variable to which different values may be given at pleasure or according to some law is called the *Independent Variable*, and the function of the independent variable is called the *Dependent Variable*.

In the first illustration x is the independent variable and $2x + 1$, the function of x , is the dependent variable.

410. To discuss functions of a variable it is necessary to suppose that the variable takes its successive values according to some definite law of change.

Where a variable takes a series of values that approach nearer and nearer a given constant, so that by taking a sufficient number of steps the difference between the variable and the constant can be made numerically less than any conceivable number however small, the constant is called the **Limit** of the variable, and the variable is said to *approach its limit*.

The variable .3, .33, .333, .3333, . . . , whose increase at each step is $\frac{1}{10}$ of the previous increase, approaches $\frac{1}{3}$ as a limit. For .3 differs from $\frac{1}{3}$ by less than $\frac{1}{10}$, .33 by less than $\frac{1}{100}$, .333 by less than $\frac{1}{1000}$, etc., and by taking a sufficient number of steps it is possible to obtain a value of the variable differing from $\frac{1}{3}$ by less than any number that can be conceived of, however small.

411. The difference between a variable and its limit is a variable whose limit is zero.

As .3, .33, .333, . . . approaches its limit $\frac{1}{3}$, the variable difference $\frac{1}{3} - .3$, $\frac{1}{3} - .33$, $\frac{1}{3} - .333$, . . . , or $\frac{1}{10}$, $\frac{1}{100}$, $\frac{1}{1000}$, . . . approaches the limit zero.

A variable may approach a constant without approaching it as a limit.

The variable 6.6, 6.66, 6.666, . . . , in approaching $6\frac{2}{3}$ as a limit, approaches 7 also, but not as a limit.

A variable in approaching its limit may be always less than its limit, or always greater, or sometimes greater and sometimes less.

The variable .3, .33, .333, . . . is always less than its limit $\frac{1}{3}$.

The variable .7, .67, .667, . . . is always greater than its limit $1 - \frac{1}{3}$.

The sum of the first n terms of the geometrical series $1, -\frac{1}{2}, \frac{1}{4}, -\frac{1}{8}, \frac{1}{16}, -\frac{1}{32}, \dots$ is a variable whose successive values $1, \frac{1}{2}, \frac{3}{4}, \frac{7}{8}, \frac{15}{16}, \frac{31}{32}, \dots$ are alternately greater and less than the limit $\frac{2}{3}$.

A variable may increase or decrease according to its law of change and become numerically greater or less than any assignable number.

The variable $1, -2, 4, -8, 16, \dots$ may become numerically greater than any number that can be assigned. The variable $1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$ may become numerically less than any number that can be assigned.

412. A variable that may become numerically greater than any assignable number is said to be **Infinite**.

The symbol of an infinite number is ∞ .

413. A variable that may become numerically less than any assignable number is said to be **Infinitesimal**.

An infinitesimal is a variable whose limit is zero.

The character 0 is used as a symbol for an infinitesimal number as well as for absolute zero, the result obtained by subtracting a number from itself.

414. A number that cannot become either infinite or infinitesimal is said to be **Finite**.

415. Interpretation of $\frac{a}{0}$.

If the numerator of the fraction $\frac{a}{x}$ is constant while the denominator decreases regularly until it becomes numerically less than any assignable number, the quotient will increase regularly and become numerically greater than any assignable number.

$$\therefore \frac{a}{0} = \infty. \quad \text{That is,}$$

If a finite number is divided by an infinitesimal number, the quotient will be an infinite number.

416. Interpretation of $\frac{a}{\infty}$.

If the numerator of the fraction $\frac{a}{x}$ is constant while the denominator increases regularly until it becomes numerically greater than any assignable number, the quotient will decrease regularly and become numerically less than any assignable number.

$$\therefore \frac{a}{\infty} = 0. \quad \text{That is,}$$

If a finite number is divided by an infinite number, the quotient will be an infinitesimal number.

417. The symbol \doteq is read 'approaches as a limit.'

The abbreviation *lim.* is used for *limit*.

$x \doteq a$ is read ' x approaches a as its limit.' The expression $\lim. x = a$ is equivalent to $x \doteq a$, and is read 'the limit of x is a .'

Though ∞ represents a variable that may transcend all finite values, it is convenient to use the expression $x \doteq \infty$ to indicate that x increases numerically *without limit*.

Thus, as $x \doteq \infty$, $\frac{1}{x} \doteq 0$.

418. PRINCIPLE 1.—*A variable cannot approach two unequal limits at the same time.*

For in approaching the more remote as a limit the variable would reach a value intermediate between the two, and thereafter in approaching one as a limit it would recede from the other, which, therefore, could not be a limit.

419. PRINCIPLE 2. — *If two variables are always equal and each approaches a limit, their limits are equal.*

For in their values the two variables are but one.

Hence, Prin. 1, the limit of their common values is their common limit.

420. 1. Since the limit of $.333 \dots$ is $\frac{1}{3}$, what is the limit of $2 + .333 \dots$, or $2.333 \dots$? of $4 + .333$, or $4.333 \dots$? of $5.333 \dots$? of $2 - .333 \dots$, or $1.666 \dots$? of $.333 \dots - .13$, or $.203 \dots$?

2. What is the limit of the algebraic sum of a constant and a variable?

PRINCIPLE 3. — *The limit of the algebraic sum of a constant and a variable is the algebraic sum of the constant and the limit of the variable.*

421. 1. Since the limit of $.333 \dots$ is $\frac{1}{3}$, what is the limit of $.666 \dots$? of $.111 \dots$?

2. Since the limit of $1 + \frac{1}{2} + \frac{1}{4} + \dots$ is 2, what is the limit of $3 + \frac{3}{2} + \frac{3}{4} + \dots$? of $\frac{1}{10} + \frac{1}{20} + \frac{1}{40} + \dots$?

3. How may the limit of the product of a variable and a constant be obtained from the limit of the variable?

PRINCIPLE 4. — *The limit of the product of a variable and a finite constant is equal to the product of the constant and the limit of the variable.*

The above principle may be established as follows:

CASE 1. — *When the limit of x is 0.*

Let k be any finite constant.

It is to be proved that $kx \doteq 0$.

However small any number, as q , may be, since $x \doteq 0$, x may be made numerically less than $q + k$.

Hence, kx may be made numerically less than q ; that is, kx may be made numerically less than any number however small.

Therefore, § 410, $kx \doteq 0$.

CASE 2. — *When the limit of x is not 0.*

Let k be any finite constant, a the limit of x , and y the variable that must be added to x to produce a .

It is to be proved that

$$kx \doteq ka.$$

Since $x + y = a$,

$$x = a - y;$$

$$\therefore kx = ka - ky.$$

Prin. 2,

$$\lim. (kx) = \lim. (ka - ky).$$

Prin. 3,

$$= ka - \lim. (ky).$$

But since, § 411, $y \doteq 0$, by Case 1,

$$\lim. ky = 0.$$

Hence,

$$\lim. (kx) = ka - 0 = ka;$$

that is,

$$kx \doteq ka.$$

NOTE. — The principle holds for the limit of a variable *divided* by a constant, since dividing by k is equivalent to multiplying by $\frac{1}{k}$.

422. PRINCIPLE 5. — *The limit of the variable sum of any finite number of variables is equal to the sum of their limits.*

The above principle may be established as follows:

Let $x \doteq a$, $y \doteq b$, $z \doteq c$, etc., to any finite number of variables, as n .

It is to be proved that $\lim. (x + y + z + \dots) = a + b + c + \dots$.

Let v_1, v_2, v_3, \dots be the variable differences between $x, y, z \dots$ and their respective limits.

Then, $x + y + z + \dots = (a + b + c + \dots) - (v_1 + v_2 + v_3 + \dots)$,

and $\lim. (x + y + z + \dots) = \lim. [(a + b + c + \dots) - (v_1 + v_2 + v_3 + \dots)]$

Prin. 3, $= a + b + c + \dots - \lim. (v_1 + v_2 + v_3 + \dots).$

Since, § 411, $v_1 \doteq 0$, $v_2 \doteq 0$, $v_3 \doteq 0$, etc., however small any number, as q , may be, each of the n variables, v_1, v_2, v_3 , etc., may be made less than $q \div n$, and hence their sum may be made less than q .

Therefore, § 410, $\lim. (v_1 + v_2 + v_3 + \dots) = 0$.

Hence, $\lim. (x + y + z + \dots) = a + b + c + \dots$.

423. PRINCIPLE 6. — *The limit of the variable product of two or more variables is equal to the product of their limits.*

The above principle may be established as follows:

Given, $x \doteq a$ and $y \doteq b$.

It is to be proved that $\lim. (xy) = ab$.

Let

$$v_1 = a - x \text{ and } v_2 = b - y.$$

Then,

$$xy = (a - v_1)(b - v_2),$$

and

$$\lim. (xy) = \lim. [ab - bv_1 - av_2 + v_1v_2]$$

Prin. 5, Prin. 4,

$$= ab - b \lim. v_1 - a \lim. v_2 + \lim. (v_1v_2)$$

§ 411,

$$= ab + \lim. (v_1v_2).$$

Since when v_1 and v_2 are near their common limit 0, their product is much less than either v_1 or v_2 ,

$$\lim. (v_1 v_2) = 0.$$

Hence,

$$\lim. (xy) = ab + 0 = ab.$$

Similarly, the principle may be established for any number of variables.

424. PRINCIPLE 7. — *The limit of the variable quotient of two variables is equal to the quotient of their limits, provided the limit of the divisor is not 0.*

The above principle may be established as follows :

Let $x = \frac{y}{z}$; and let $\lim. z$ be not 0.

It is to be proved that $\lim. x = \frac{\lim. y}{\lim. z}$.

Since $x = \frac{y}{z}$, $y = xz$.

Prin. 6, $\lim. y = \lim. x \cdot \lim. z$.

Therefore, $\lim. x = \frac{\lim. y}{\lim. z}$.

The principle has no meaning when $z \doteq 0$, since $\lim. y$ cannot be divided by 0.

425. When by causing a variable x to approach sufficiently near to a it is possible to make the value of a given function of x approach as near as we please to a finite constant l , l is called the **limit of the function when $x \doteq a$** .

Suppose that .1, .11, .111, .1111, ... are successive values of x approaching $\frac{1}{10}$ as a limit. Then, the corresponding values of $1 - 2x$, a function of x , are .8, .78, .778, .7778, values of a variable approaching $\frac{4}{5}$ as a limit. As $x \doteq \frac{1}{10}$, the function of x , $1 - 2x$, $\doteq \frac{4}{5}$; for by causing x to approach sufficiently near to $\frac{1}{10}$ it is possible to make $1 - 2x$ approach as near as we please to $\frac{4}{5}$.

The expression

$$\lim. [\text{function of } x]_{x \doteq a}$$

is read 'limit of function of x as x approaches a as a limit.'

Thus, $\lim. (1 - 2x)_{x \doteq \frac{1}{10}} = \frac{4}{5}$ indicates that as x approaches its limit $\frac{1}{10}$, $1 - 2x$ approaches the limit $\frac{4}{5}$.

426. In finding the limiting values of the functions given in the following examples, the student is expected to apply the principles that have been established above.

Finding the limiting value of a function of x as $x \doteq a$ is called **evaluating** the function for $x = a$.

EXAMPLES

If $x \doteq a$, $y \doteq 2$, and $z \doteq 0$, find the limit of

1. $x + y + z.$

4. $x - \frac{1}{2}y + ax - y^2.$

2. $axy - x^2.$

5. $(x + y)x - (x - y)z.$

3. $\frac{x}{2} - \frac{y}{a}.$

6. $\frac{a + x + z}{x - y} + \frac{x + y}{a}.$

Find the value of

7. $\text{Lim.} \left[\frac{x^2 - 5x + 8}{x - 2} \right]_{x \doteq 2}$

8. $\text{Lim.} \left[\frac{x^3 + 4x + 1}{x^4 + x^2 + 1} \right]_{x \doteq -1}$

9. $\text{Lim.} \left[\frac{x^2 - 3x + 4}{x + 6} \right]_{x \doteq 2}$

10. $\text{Lim.} \left[\frac{1 + x^n}{1 + x} \right]_{x \doteq 1}$

427. Interpretation of $\frac{0}{0}.$

If both terms of a fraction are divided by any number, and both terms of the resulting fraction are divided by any number, and this process is continued indefinitely, though the fraction will not change in value, it will finally assume the form $\frac{0}{0}.$

Since *any* fraction may be reduced in this way to the form $\frac{0}{0}$, $\frac{0}{0}$ may have any value; that is, is *indeterminate*.

The expressions $\frac{\infty}{\infty}$, $0 \times \infty$, and $\infty - \infty$ are indeterminate, for each may be reduced to the form $\frac{0}{0}.$

Thus,
$$\frac{\infty}{\infty} = \frac{a}{0} \div \frac{b}{0} = \frac{a}{0} \times \frac{0}{b} = \frac{a \times 0}{0 \times b} = \frac{0}{0};$$

$$0 \times \infty = 0 \times \frac{a}{0} = \frac{0 \times a}{0} = \frac{0}{0};$$

and
$$\infty - \infty = \frac{a}{0} - \frac{b}{0} = \frac{a \times 0 - b \times 0}{0 \times 0} = \frac{0}{0}.$$

428. Since every function of a single variable is a variable, it is evident that the preceding principles apply to functions of a variable. But, the indeterminate forms having now been introduced, the principles must be understood to be inapplicable to the evaluation of functions when the results obtained involve the forms $\infty - \infty$, $0 \times \infty$, $\frac{0}{0}$, and $\frac{\infty}{\infty}$. Thus, to apply to functions of a variable, Prin. 5, 6, and 7 should be modified as follows:

The limit of the sum of two or more functions of x is equal to the sum of their limits, provided this does not become $\infty - \infty$.

The limit of the product of two or more functions of x is equal to the product of their limits, provided this does not become $0 \times \infty$.

The limit of the quotient of two functions of x is equal to the quotient of their limits, provided this does not become $\frac{0}{0}$ or $\frac{\infty}{\infty}$.

429. The preceding principles of limits lead to the conclusion that the limit of a function is found by *substituting* the limits of the variables for the variables, except when such a substitution gives an indeterminate form (§ 428).

Thus, if $\lim. x = 5$ and $\lim. y = 2$, the limit of $4x - 3y$ is found by substituting 5 for x and 2 for y in the function $4x - 3y$. But if substitution is employed directly to evaluate the functions $\frac{1}{x-1} - \frac{1}{x(x-1)}$, $\frac{a}{y} \times y$, and $(x^2 - 1) \div (x - 1)$ when $x \doteq 1$ and $y \doteq 0$, these functions take the forms $\infty - \infty$, $\infty \times 0$, and $0 \div 0$, respectively.

When the method of evaluation by substitution in the given function fails, the evaluation of the function is performed by the aid of Prin. 2.

Thus, to evaluate $(x^2 - 1) \div (x - 1)$ when $x \doteq 1$, find another function of x , as $x + 1$, equal to the given function $(x^2 - 1) \div (x - 1)$ for all values assumed by x while approaching the limit 1.

If x takes the successive values

$$2, \frac{3}{2}, \frac{5}{4}, \frac{7}{8}, \frac{9}{16}, \frac{11}{32}, \dots, \text{approaching } 1,$$

then both functions $(x^2 - 1) \div (x - 1)$ and $x + 1$, take the successive values

$$3, \frac{5}{2}, \frac{7}{4}, \frac{9}{8}, \frac{11}{16}, \frac{13}{32}, \dots, \text{approaching } 2.$$

Since the two functions are equal for all values of x as x approaches its limit 1 by Prin. 2 they have the same limit. This limit is $\lim. (x + 1)_{x \doteq 1}$, which by substituting $\lim. x$ for x is found to be $1 + 1$, or 2.

EXAMPLES

Find the value of

1. $\text{Lim.} \left[\frac{x^3 - 8}{x^2 - 4} \right]_{x \doteq 2}$

3. $\text{Lim.} \left[\frac{x^4 - a^4}{x^3 + a^3} \right]_{x \doteq -a}$

2. $\text{Lim.} \left[\frac{1 - x^5}{1 - x} \right]_{x \doteq 1}$

4. $\text{Lim.} \left[\frac{x^3 + b^3}{x^2 - b^2} \right]_{x \doteq -b}$

5. $\text{Lim.} \left[\frac{x^2 - 5x + 6}{x^2 - 4} \right]_{x \doteq 2}$

6. $\text{Lim.} \left[\frac{x^4 - 3x^3 + 5x^2 - 2x - 1}{x^3 - 8x^2 + 7} \right]_{x \doteq 1}$

7. Find the limiting value of $\frac{4x^3 - 2x^2 + 3x + 1}{3x^3 - x^2 + x + 2}$ when $x \doteq 0$ and also when $x \doteq \infty$.

SOLUTION.—As x approaches the limit 0, the first three terms of the numerator and also of the denominator become infinitely small as compared with the fourth, and, consequently, may be neglected. Hence, when $x \doteq 0$, the fraction approaches the limiting value $\frac{1}{2}$.

As $x \doteq \infty$, that is, as x becomes indefinitely greater, the last three terms of the numerator and also of the denominator become infinitely small as compared with the first, and, consequently, may be neglected. Hence, when $x \doteq \infty$, the fraction approaches the limiting value $\frac{4x^3}{3x^3}$, or $\frac{4}{3}$.

Find the limiting values of the following when $x \doteq 0$ and when $x \doteq \infty$:

8. $\frac{1 + x^2 + x^4 + x^6}{1 - x^2 - x^4 - 2x^6}$

12. $\frac{5x + 10}{x^2 + 2x + 2}$

9. $\frac{5x^3 - x^2 + 4x + 2}{2x^3 + 3x^2 - x + 1}$

13. $\frac{3x - 4}{x^2 - x - 8}$

10. $\frac{4x^4 - 3x^3 + x + 1}{2x^4 - x^3 - x^2 + x + 1}$

14. $\frac{2x^2 - 4x + 1}{2x - 1}$

11. $\frac{2x^4 - 3x^3 + 2x^2 - 2}{x^4 - 2x^2 + x + 1}$

15. $\frac{4x^3 + 5x^2 + 2x}{2x^3 + x + 1}$

INTERPRETATION OF RESULTS

430. When the roots obtained by solving an *equation* satisfy the equation, the solution has been properly performed; but the results found in solving a *problem* may sometimes be at variance with some condition of the problem. Consequently, the interpretation of results becomes important.

POSITIVE RESULTS

431. Since the numbers sought in the solution of a problem are arithmetical rather than algebraic, when positive results are obtained, it is not likely that they will conflict with the conditions of the problem. Sometimes, however, even a positive result violates one or more of the conditions of a problem.

In such cases the problem is *impossible*.

432. 1. A club consisting of 25 members raised the sum of \$ 13 by assessing the men 80 cents each and the women 40 cents each. How many men were there, and how many women?

SOLUTION

Let	$x =$ the number of men.
Then,	$25 - x =$ the number of women ;
\therefore	$\frac{4}{5}x + \frac{2}{5}(25 - x) = 13 ;$
whence,	$x = 7\frac{1}{2},$
and	$25 - x = 17\frac{1}{2}.$

Though the numbers found will satisfy the equation, yet since the number of men and the number of women cannot be fractional, the problem is *impossible*.

2. The second digit of a number expressed by two digits is twice the first, and 4 times the first digit is 9 greater than the second. What is the number?

SOLUTION

Let	$x =$ the first digit.
Then,	$2x =$ the second digit;
\therefore	$4x = 2x + 9;$
whence,	$x = 4\frac{1}{2}$, the first digit,
and	$2x = 9$, the second digit.

While these numbers satisfy the equation, they fail to satisfy the implied condition that the digits must be integers.

Hence, the problem is *impossible*.

NEGATIVE RESULTS

433. A few examples will suggest the methods to be employed in the interpretation of negative results.

1. If A is 40 years old and B is 30, in how many years will A be twice as old as B?

SOLUTION

Let	$x =$ the number of years after which A will be twice as old as B.
Then,	$40 + x = 2(30 + x);$
whence,	$x = -20.$

Though the result is *algebraically* correct, inasmuch as -20 substituted for x satisfies the equation, nevertheless it is *arithmetically* absurd. Hence, the conditions of the problem are inconsistent with each other. Had the result been $+20$, the statement that A would be twice as old as B in 20 years would have been arithmetically reasonable. However, since $-x = +20$, the equation will give a result arithmetically reasonable, if $-x$ is substituted for x ; that is, if x is taken to represent the number of years *since* A was twice as old as B.

The conditions of the problem should, therefore, be modified as follows: If A is 40 years old and B is 30, how many years ago was A twice as old as B?

2. How much money has A, if $\frac{1}{4}$ of his money is 5 dollars more than $\frac{1}{3}$ of it?

SOLUTION

Let x = the number of dollars A has.

Then $\frac{x}{4} - \frac{x}{3} = 5$.

Solving, $x = -60$.

While the result -60 satisfies the equation, it violates the supposition, made in the problem, that A has some money.

If $-x$ is substituted for x , the equation becomes

$$\frac{x}{3} - \frac{x}{4} = 5; \text{ whence, } x = 60.$$

The problem when modified to express conditions *arithmetically* reasonable will be: How much money has A, if $\frac{1}{4}$ of it is 5 dollars less than $\frac{1}{3}$ of it; or, if -60 dollars is interpreted 60 dollars *in debt*: How much money does A owe, if $\frac{1}{3}$ of what he owes is 5 dollars more than $\frac{1}{4}$ of it?

434. From the above discussions we may infer:

1. *A negative result indicates that some quantity in the problem has been applied in the wrong direction.*

2. *A possible problem analogous to the given problem may be found by changing the absurd conditions to their opposites.*

PROBLEMS

435. Interpret arithmetically the negative results obtained by solving the following:

1. If A is 40 years old and B is 25, in how many years will B be half as old as A?

2. Find the numbers whose sum is 6 and difference 10.

3. What fraction is equal to $\frac{3}{8}$, if 1 is added to its numerator or to $\frac{3}{8}$ if 1 is added to its denominator?

4. A boy bought some apples for 24 cents. Had he received 4 more for that sum, the cost of each would have been 1 cent less. How many did he buy?

5. A man worked 7 days, during which he had his son with him 3 days, and received 22 shillings. He afterwards worked 5 days, during which he had his son with him 1 day, and received 18 shillings. What were the daily wages of each?

ZERO RESULTS

436. When the result obtained by solving a problem is zero, it may sometimes indicate that the problem is impossible, and sometimes it may be the proper answer to the question.

1. A dealer had two kinds of tea worth 75 and 60 cents per pound, respectively. How many pounds of each must he take to make a mixture of 45 pounds worth \$ 27?

SOLUTION

Let x = the number of pounds of the better kind.

Then, $45 - x$ = the number of pounds of the poorer kind;

$$\therefore \frac{3}{4}x + \frac{2}{3}(45 - x) = 27;$$

whence,

$$x = 0.$$

This result means that no such mixture can be made. In fact, 45 pounds of the poorer tea is worth \$ 27.

2. A is 48 years old, and B is 16 years old. After how many years will A be 3 times as old as B?

SOLUTION

Let x = the required number of years.

Then, $48 + x = 3(16 + x)$.

Solving, $x = 0$.

This result indicates that A is *now* 3 times as old as B.

3. What number is equal to the square of itself?

SOLUTION

Let x = the number.

Then, $x = x^2$,

$$x(x - 1) = 0;$$

$$\therefore x = 1 \text{ or } 0.$$

These results indicate that no number except 1 is equal to the square of itself.

INDETERMINATE RESULTS

437. 1. A lady being asked her age replied, "If from 3 times my age you take 4 years and divide the difference by 2, you will have twice my age less half of my age 4 years hence." What was her age?

SOLUTION

Let

x = the number of years.

Then,

$$\frac{3x - 4}{2} = 2x - \frac{x + 4}{2}, \quad (1)$$

$$3x - 4 = 4x - x - 4, \quad (2)$$

$$(3 - 3)x = 0; \quad (3)$$

$$\therefore x = \frac{0}{0}. \quad (4)$$

Since (2) may be reduced to the identity $3x - 4 = 3x - 4$, it may be satisfied by any value of x whatever. This relation, § 427, is expressed by (4). Hence, the problem is *indeterminate*.

PROBLEMS

438. 1. If twice a certain number is subtracted from the square of the number, the result will be 1 less than the square of a number 1 less. What is the number?

2. A father is 30 years older than his son, and the sum of their ages is 30 years less than twice the father's age. What is the son's age?

3. The sum of the first and third of three consecutive integers is equal to twice the second. What are the integers?

4. A bought 400 sheep in two flocks, paying \$1.50 per head for the first flock and \$2 per head for the second. He lost 30 of the first flock and 56 of the second, but by selling the rest of the first flock at \$2 per head and the rest of the second at \$2.50 per head, he neither lost nor gained. How many sheep were there in each flock originally?

5. A and B receive the same monthly salary. A is employed 10 months in the year and his annual expenses are \$600. B is employed 8 months in the year and his annual expenses are \$480. If A saves as much money in 4 years as B saves in 5 years, what is the monthly salary of each?

INFINITE RESULTS

439. An infinite result indicates that the problem is *impossible*.

1. If a man's yearly income is a dollars and his yearly expenses are a dollars, in how many years will he have saved b dollars?

SOLUTION

Let x = the required number of years.

Then,
$$x = \frac{b}{a - a} = \frac{b}{0}, \text{ or } \infty.$$

That is, he will *never* have saved b dollars in this way.

2. A reservoir is fitted with three pipes. One pipe can fill the reservoir in 15 hours, the second can fill it in $\frac{2}{3}$ of that time, and the third pipe can empty it in 6 hours. If the reservoir is full and the three pipes are opened, in what time will it be emptied?

SOLUTION

Let x = the required number of hours.

Then,
$$\frac{1}{15} + \frac{1}{10} - \frac{1}{6} = \frac{1}{x}$$

Solving,
$$x = \frac{60}{0}, \text{ or } \infty.$$

That is, the reservoir will *never* be emptied under these conditions.

3. What number added to both terms of the fraction $\frac{1}{2}$ will make the fraction equal to 1?

SOLUTION

Let x = the number.

Then,
$$\frac{1+x}{2+x} = 1.$$

Clearing,
$$1+x = 2+x.$$


Solving,
$$x = \frac{1}{0} \text{ or } \frac{-1}{0};$$

that is,
$$x = +\infty \text{ or } -\infty.$$

Consequently, there is no such number; but the larger the number in numerical value, the nearer will the resulting fraction approach the value 1.

THE PROBLEM OF THE COURIERS

440. Two couriers, A and B, travel on the same road in the direction from X to Y at the rates of m and n miles an hour, respectively. At a certain time, say 12 o'clock, A is at P , and B is at Q , a miles from P . Find when and where they are together.



SOLUTION

Suppose that time reckoned from 12 o'clock toward a later time is positive, and toward an earlier time, negative; also, that distances measured from P toward the right are positive, and toward the left, negative.

Let x represent the number of hours from 12 o'clock, and y the number of miles from P , when A and B are together. Then, they will be together $y - a$ miles from Q .

Since A travels mx miles and B travels nx miles before they are together,

$$y = mx, \quad (1)$$

and $y - a = nx. \quad (2)$

Solving (1) and (2),

$$x = \frac{a}{m - n}, \text{ the required time.} \quad (3)$$

$$y = \frac{ma}{m - n}, \text{ the required distance.} \quad (4)$$

DISCUSSION

1. When $a > 0$ and $m > n$.

When $a > 0$ and $m > n$, the numerator and denominator in (3) and also in (4) are positive; hence, x and y are positive.

That is, A overtakes B sometime after 12 o'clock, somewhere at the right of P .

2. When $a > 0$ and $m < n$.

When $a > 0$ and $m < n$, both x and y are negative.

That is, at 12 o'clock B is ahead of A and gaining on him, and they were together sometime before 12 o'clock and somewhere at the left of P .

3. When $a > 0$ and $m = n$.

When $a > 0$ and $m = n$, x and y are positive and infinitely great.

That is, at 12 o'clock B is ahead of A and traveling at the same rate; consequently, he will never be overtaken by A.

4. When $a = 0$ and $m > n$ or $m < n$.

When $a = 0$ and $m > n$ or $m < n$, $x = 0$ and $y = 0$.

If $m > n$, $x = +0$ and $y = +0$. That is, at 12 o'clock A and B are together, and A is passing B.

If $m < n$, $x = -0$ and $y = -0$. That is, at 12 o'clock A and B are together, and B is passing A.

5. When $a = 0$ and $m = n$.

When $a = 0$ and $m = n$, $x = \frac{0}{0}$ and $y = \frac{0}{0}$.

That is, A and B are together at 12 o'clock, and since they travel at the same rate they will be together at all times.



INDETERMINATE EQUATIONS



441. While a problem that presents more unknown literal numbers than independent equations involving them is in general indeterminate (§ 214), yet frequently by the introduction of a condition or conditions not leading to equations, the number of values of the unknown numbers may be limited and these values algebraically determined. A common condition is that the results shall be positive integers.

1. Solve the equation $5x + 3y = 35$ in positive integers.

SOLUTION

Since x and y are positive integers, $5x$ must be equal to 5 or a multiple of 5, and $3y$ must be equal to 3 or a multiple of 3. Since the sum of these multiples is 35, if the multiples of 5 are subtracted from 35, one or more of the remainders will be a multiple of 3, if the problem is possible.

The only multiples of 5 that subtracted from 35 leave multiples of 3 are 5 and 4 times 5.

$\therefore x = 1$ or 4 ; whence, $y = 10$ or 5 .

Or, since x must be a positive integer and by transposition $5x = 35 - 3y$, the values of x must be 1, 2, 3, 4, 5, or 6, if the equation is possible. Substituting these values of x in the given equation and rejecting all those that give negative or fractional values for y , the positive integral values are found to be $x = 1$ or 4 , and $y = 10$ or 5 .

2. Solve the equation $5x + 8y = 107$ in positive integers.

SOLUTION

$$5x + 8y = 107. \quad (1)$$

Dividing by 5,
$$x + y + \frac{3y}{5} = 21 + \frac{2}{5}. \quad (2)$$

Collecting integral and fractional terms,

$$x + y - 21 = \frac{2 - 3y}{5}. \quad (3)$$

Since $x + y - 21$ is integral,
$$\frac{2 - 3y}{5} = \text{an integer}. \quad (4)$$

If $\frac{2 - 3y}{5} = w$, an integer, then, $y = \frac{2 - 5w}{3}$, which is in the fractional form. To avoid this, the coefficient of y in the number placed equal to w should be made equal to *unity*. Since $\frac{2 - 3y}{5}$ is equal to an integer, any multiple of it is equal to an integer. Since 5 is contained in 3 times $-3y$, $-2y$ times with a remainder of y , multiplying (4) by 3,

$$\begin{aligned} \frac{6 - 9y}{5} &= \text{an integer} \\ &= 1 - 2y + \frac{1 + y}{5}. \end{aligned} \quad (5)$$

Then, let
$$\frac{1 + y}{5} = w, \text{ an integer}. \quad (6)$$

Solving for y ,
$$y = 5w - 1. \quad (7)$$

Substituting in (3),
$$x = 23 - 8w. \quad (8)$$

Equations (7) and (8) are called the *general solution* of the given equation in integers.

To make y and x *positive* integers, it is evident from (7) that we must take $w > 0$; and from (8) that we must take $w < 3$.

Since w is an integer greater than 0 and less than 3, $w = 1$ or 2.

When $w = 1$,
$$x = 15, y = 4;$$

when $w = 2$,
$$x = 7, y = 9.$$

3. Determine whether the equation $10x + 15 = 53$ may be satisfied by integral values of x and y .

SOLUTION. — Dividing by 5, $2x + 3y = \frac{53}{5}$.

If x and y are integers, the first member is integral.

Since the first member is equal to the fraction $\frac{53}{5}$, it cannot be an integer. Hence, x and y cannot be integers at the same time; that is, the equation is not satisfied by integral values of x and y .

Solve the following equations in positive integers:

4. $5x + 3y = 49.$

8. $12x + 5y = 61.$

5. $3x + 2y = 5.$

9. $6x + 7y = 72.$

6. $2x + 7y = 48.$

10. $5x + 9y = 75.$

7. $8x + 5y = 80.$

11. $6x + 9y = 100.$

Find the least integral values of x and y in the following:

12. $2x = 9 + 3y.$

14. $7x - 2y = 6.$

13. $5y = 2x + 7.$

15. $5x - 3y = 1.$

16. Solve the equations $\begin{cases} x+y+z=6 \\ 2x+y-z=7 \end{cases}$ in positive integers.

SOLUTION

$$x + y + z = 6. \quad (1)$$

$$2x + y - z = 7. \quad (2)$$

Adding, $3x + 2y = 13. \quad (3)$

Solving (3) for positive integers, $x = 3$ and $y = 2.$

Substituting in (1), $z = 1.$

Solve the following equations in positive integers:

$$17. \begin{cases} x + y + z = 8, \\ x - y + 2z = 6. \end{cases}$$

$$19. \begin{cases} 3x + 2y = 17, \\ y + 2z = 14. \end{cases}$$

$$18. \begin{cases} 2x + 3y + z = 15, \\ 3x + y - z = 8. \end{cases}$$

$$20. \begin{cases} y + z = 7, \\ 3x - z = 7. \end{cases}$$

21. Separate 100 into two parts one of which is a multiple of 11, and the other a multiple of 6.

22. In what ways may a weight of 19 pounds be weighed with 5-pound and 2-pound weights?

23. A man has \$300 that he wishes to expend for cows and sheep. If cows cost \$45 apiece and sheep \$6 apiece, how many can he buy of each?

24. If 9 apples and 5 oranges together cost 52 cents, what is the cost of one of each?

25. A grocer sold two packages of sugar for \$1.25. One package contained a certain number of pounds of 7-cent sugar, the other a certain number of pounds of 5-cent sugar. How many pounds were there in each package?

26. A man sold 9 animals—sheep, hogs, and cows—for \$100. If he received \$3 for a sheep, \$6 for a hog, and \$35 for a cow, how many of each did he sell?

27. A woman expended 93 cents for 14 yards of cloth, some at 5, some at 7, and the rest at 10 cents a yard. How many whole yards of each did she buy?

28. Divide 74 into three parts that shall give integral quotients when divided by 5, 6, and 7, respectively, the sum of which quotients shall be 12.

29. A purse contained 30 coins, consisting of half-dollars, quarters, and dimes. How many coins of each kind were there, if their aggregate value was \$6.50?

30. A man bought 100 animals for \$99. There were pigs, sheep, and ducks. If he paid \$6 for a pig, \$4 for a sheep, and 50 cents for a duck, how many of each did he buy?

31. What is the least number that will contain 25 with a remainder of 1, and 33 with a remainder of 2?

32. Find the least number that divided by 10 and by 11 will leave remainders of 3 and 6, respectively.

33. What is the least number that will contain 2, 3, 4, 5, and 6, each with a remainder of 1, and 7 without a remainder?

34. A man selling eggs to a grocer took them out of his basket 4 at a time and there was 1 egg over. The grocer put them into a box 5 at a time and there were 3 over. Both lost the count; but knowing that there were between 6 and 7 dozen eggs, the grocer paid for $6\frac{1}{2}$ dozen. How many eggs did he lose?

35. Four boys have a pile of marbles. A throws away 1 and takes $\frac{1}{4}$ of the remainder; B throws away 1 and takes $\frac{1}{4}$ of the remainder; C throws away 1 and takes $\frac{1}{4}$ of the remainder; D throws away 1, and each boy takes $\frac{1}{4}$ of the remainder. At least how many marbles must have been in the pile, and how many does each boy now have?

THE BINOMIAL THEOREM



442. The Binomial Theorem derives a formula by means of which any power of a binomial may be expanded into a series, whether the index of the power is positive or negative, integral or fractional.

POSITIVE INTEGRAL EXPONENTS

443. The powers of $(a + x)$, expanded in § 221, may be written :

$$(a + x)^2 = a^2 + 2 ax + \frac{2 \cdot 1}{1 \cdot 2} x^2.$$

$$(a + x)^3 = a^3 + 3 a^2 x + \frac{3 \cdot 2}{1 \cdot 2} ax^2 + \frac{3 \cdot 2 \cdot 1}{1 \cdot 2 \cdot 3} x^3.$$

$$(a + x)^4 = a^4 + 4 a^3 x + \frac{4 \cdot 3}{1 \cdot 2} a^2 x^2 + \frac{4 \cdot 3 \cdot 2}{1 \cdot 2 \cdot 3} ax^3 + \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 2 \cdot 3 \cdot 4} x^4.$$

$$(a + x)^5 = a^5 + 5 a^4 x + \frac{5 \cdot 4}{1 \cdot 2} a^3 x^2 + \frac{5 \cdot 4 \cdot 3}{1 \cdot 2 \cdot 3} a^2 x^3 + \frac{5 \cdot 4 \cdot 3 \cdot 2}{1 \cdot 2 \cdot 3 \cdot 4} ax^4 + \frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} x^5.$$

If the law of development revealed in the above is assumed to apply to the expansion of any power of any binomial, as the n th power of $(a + x)$, the result is

$$(a+x)^n = a^n + na^{n-1}x + \frac{n(n-1)}{1 \cdot 2} a^{n-2}x^2 + \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3} a^{n-3}x^3 + \dots (1)$$

From formula (1) it is evident that in any term,

1. The exponent of x is 1 less than the number of the term.

Hence, the exponent of x in the $(r + 1)$ th term is r .

2. The exponent of a is n minus the exponent of x .

Hence, the exponent of a in the $(r + 1)$ th term is $n - r$.

3. The number of factors in each term of the coefficient is 1 less than the number of the term.

Hence, the coefficient of the $(r+1)$ th term has r factors in each term.

Therefore, the $(r+1)$ th, or *general term*, is

$$\frac{n(n-1)(n-2) \cdots \text{to } r \text{ factors}}{1 \cdot 2 \cdot 3 \cdots \text{to } r \text{ factors}} a^{n-r} x^r. \quad (2)$$

Since, when there are two factors in the numerator, the last is $n-1$, when there are three factors, $n-2$, when there are four factors, $n-3$, etc.; when there are r factors, the last is $n-(r-1)$, or $n-r+1$. Hence, (2) may be written

$$\frac{n(n-1)(n-2) \cdots (n-r+1)}{1 \cdot 2 \cdot 3 \cdots r} a^{n-r} x^r. \quad (3)$$

Hence, the full form of (1) is

$$\begin{aligned} (a+x)^n = & a^n + na^{n-1}x + \frac{n(n-1)}{1 \cdot 2} a^{n-2}x^2 + \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3} a^{n-3}x^3 + \cdots \\ & + \frac{n(n-1)(n-2) \cdots (n-r+1)}{1 \cdot 2 \cdot 3 \cdots r} a^{n-r}x^r + \cdots + x^n. \quad (I) \end{aligned}$$

This is called the **Binomial Formula**.

444. Since it has already been proved, by actual multiplication, that the binomial formula is true for the second, third, fourth, and fifth powers of a binomial, it remains to discover whether it is true for powers higher than the fifth.

If the binomial theorem, when *assumed* to be true for the n th power, can be *proved* to be true for the $(n+1)$ th power, it will then have been proved to be true for the sixth power, since it is known to be true for the fifth power; also for the seventh power, being true for the sixth power; and in like manner for each succeeding power.

It then remains to prove that if (I) is true for the n th power, it will hold true for the $(n+1)$ th power.

To find the expansion of $(a+x)^{n+1}$ (I) may be multiplied by $a+x$. But since the $(r+1)$ th term of the product will be the algebraic sum of a times the $(r+1)$ th term of $(a+x)^n$ and x times the r th term of $(a+x)^n$, (I) should be prepared for multiplication

by writing also the r th term, obtained from the $(r+1)$ th term by substituting r for $r+1$, or $r-1$ for r . Then (I) is written

$$\begin{aligned}(a+x)^n &= a^n + na^{n-1}x + \frac{n(n-1)}{1 \cdot 2} a^{n-2}x^2 + \dots \\ &+ \frac{n(n-1)(n-2) \dots (n-r+2)}{1 \cdot 2 \cdot 3 \dots (r-1)} a^{n-r+1}x^{r-1} \\ &+ \frac{n(n-1)(n-2) \dots (n-r+2)(n-r+1)}{1 \cdot 2 \cdot 3 \dots (r-1)r} a^{n-r}x^r + \dots + x^n\end{aligned}$$

Multiplying both members by $a+x$,

$$\begin{aligned}(a+x)^{n+1} &= a^{n+1} + n \left| \begin{array}{c} a^n x + \frac{n(n-1)}{1 \cdot 2} a^{n-1} x^2 + \dots \\ + 1 \end{array} \right| + n \left| \begin{array}{c} a^{n-1} x^2 + \dots \\ + \dots \end{array} \right| + \dots \\ &+ \frac{n(n-1)(n-2) \dots (n-r+2)(n-r+1)}{1 \cdot 2 \cdot 3 \dots (r-1)r} \left| \begin{array}{c} a^{n-r+1} x^r \\ + \dots \end{array} \right| + \dots + x^{n+1} \\ &= a^{n+1} + (n+1)a^n x + \left[\frac{n(n-1)}{2} + n \right] a^{n-1} x^2 + \dots \\ &+ \left(\frac{n-r+1}{r} + 1 \right) \left(\frac{n(n-1)(n-2) \dots (n-r+2)}{1 \cdot 2 \cdot 3 \dots (r-1)} \right) a^{n-r+1} x^r + \dots + x^{n+1} \\ &= a^{n+1} + (n+1)a^n x + \frac{(n+1)n}{1 \cdot 2} a^{n-1} x^2 + \dots \\ &+ \left(\frac{n+1}{r} \right) \left(\frac{n(n-1)(n-2) \dots (n-r+2)}{1 \cdot 2 \cdot 3 \dots (r-1)} \right) a^{n-r+1} x^r + \dots + x^{n+1}.\end{aligned}$$

That is,

$$\begin{aligned}(a+x)^{n+1} &= a^{n+1} + (n+1)a^n x + \frac{(n+1)n}{1 \cdot 2} a^{n-1} x^2 + \dots \\ &+ \frac{(n+1)n(n-1) \dots (n-r+2)}{1 \cdot 2 \cdot 3 \dots r} a^{n-r+1} x^r + \dots + x^{n+1}. \quad (\text{II})\end{aligned}$$

Since upon comparison it may be seen that (II) and (I) have the same form, $n+1$ in one taking the place of n in the other, (II) and (I) must express the same law of formation.

Therefore, if the formula is true for the n th power, it holds true for the $(n+1)$ th power.

Hence, the binomial formula is true for any positive integral exponent.

This method of proof is called a proof by *Mathematical Induction*.

445. If $-x$ is substituted for x in (I), the terms that contain the odd powers of $-x$ will be negative, and those that contain the even powers will be positive. Therefore,

$$(a - x)^n = a^n - na^{n-1}x + \frac{n(n-1)}{1 \cdot 2} a^{n-2}x^2 - \dots \quad (\text{III})$$

If $a = 1$, (I) becomes

$$(1 + x)^n = 1 + nx + \frac{n(n-1)}{1 \cdot 2} x^2 + \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3} x^3 + \dots \quad (\text{IV})$$

446. From (I) it is seen that the last factor in the numerator of the coefficient is n for the 2d term, $n-1$ for the 3d term, $n-2$ for the 4th term, $n-(n-2)$, or 2, for the n th term, and $n-(n-1)$, or 1, for the $(n+1)$ th term; and that the coefficient of the $(n+2)$ th term, and that of each succeeding term, contains the factor $n-n$, or 0, and therefore reduces to 0. Hence,

When n is a positive integer, the series formed by expanding $(a+x)^n$ is finite and has $n+1$ terms.

447. By formula (I), when n is a positive integer,

$$(a+x)^n = a^n + na^{n-1}x + \frac{n(n-1)}{1 \cdot 2} a^{n-2}x^2 + \dots + \frac{n(n-1) \dots 2 \cdot 1}{1 \cdot 2 \dots (n-1)n} x^n.$$

$$(x+a)^n = x^n + nx^{n-1}a + \frac{n(n-1)}{1 \cdot 2} x^{n-2}a^2 + \dots + \frac{n(n-1) \dots 2 \cdot 1}{1 \cdot 2 \dots (n-1)n} a^n.$$

A comparison of the two series shows that:

The coefficients of the latter half of the expansion of $(a+x)^n$, when n is a positive integer, are the same as those of the first half, written in the reverse order.

EXAMPLES

1. Expand $(3a-2b)^4$.

SOLUTION. — Substituting $3a$ for a , $2b$ for x , and 4 for n in (III),

$$\begin{aligned} (3a-2b)^4 &= (3a)^4 - 4(3a)^3(2b) + \frac{4 \cdot 3}{1 \cdot 2} (3a)^2(2b)^2 - \frac{4 \cdot 3 \cdot 2}{1 \cdot 2 \cdot 3} (3a)(2b)^3 \\ &\quad + \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 2 \cdot 3 \cdot 4} (2b)^4 \\ &= 81a^4 - 216a^3b + 216a^2b^2 - 96ab^3 + 16b^4. \end{aligned}$$

2. Expand $\left(\frac{b}{2} + bx\right)^5$.

SOLUTION

Since $\left(\frac{b}{2} + bx\right)^5 = \left[\frac{b}{2}(1 + 2x)\right]^5 = \frac{b^5}{32}(1 + 2x)^5$,

$(1 + 2x)^5$ may be expanded by (IV), and the result multiplied by $\frac{b^5}{32}$.

Expand:

3. $(b - n)^7$.

10. $\left(1 + \frac{2}{x^2}\right)^5$.

15. $\left(\sqrt{2} + \frac{1}{\sqrt{x}}\right)^3$.

4. $(1 + a^{-1})^4$.

11. $\left(\frac{a}{x} - \frac{x}{a}\right)^5$.

16. $\left(x^{\frac{n-1}{n}} - x^{\frac{1}{n}}\right)^4$.

5. $(2 - 3x)^6$.

12. $\left(\frac{1}{x} - \frac{a}{y}\right)^3$.

17. $(ax^{-2} - b\sqrt{x})^6$.

6. $(x^2 - x)^8$.

18. $\left(\frac{\sqrt{a}}{\sqrt[3]{b}} - \frac{\sqrt{b}}{\sqrt{a^3}}\right)^6$.

7. $(x + x^{-1})^6$.

13. $(\sqrt[3]{a^2} + \sqrt[4]{b^5})^3$.

19. $\left(\frac{x}{y}\sqrt{\frac{x}{y}} + \frac{2}{3}\sqrt{\frac{2}{3}}\right)^3$.

8. $(2a + \sqrt{x})^3$.

14. $(2\sqrt{2} - \sqrt[3]{3})^6$.

9. $(a + a\sqrt{a})^4$.

448. To find any term.

Any term of the expansion of a power of a binomial may be obtained by substitution in (2) or (3), § 443.

Since the exponent of x in the $(r + 1)$ th term is r , the sign of the general term is + if r is even, and - if r is odd.

EXAMPLES

1. Find the 12th term of $(a - b)^{14}$.

SOLUTION

$$\begin{aligned} \text{12th term} &= \frac{14 \cdot 13 \cdot 12 \cdot 11 \cdot 10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8 \cdot 9 \cdot 10 \cdot 11} a^2(-b)^{11} \\ &= -\frac{14 \cdot 13 \cdot 12}{1 \cdot 2 \cdot 3} a^2b^{11} = -364 a^2b^{11}. \end{aligned}$$

OR

By § 447, since there are 15 terms, the coefficient of the 12th term, or the 4th term from the end, is equal to that of the 4th term from the beginning.

$$\therefore \text{12th term} = -\frac{14 \cdot 13 \cdot 12}{1 \cdot 2 \cdot 3} a^2b^{11} = -364 a^2b^{11}.$$

2. In the expansion $(x^2 + 2x)^{11}$, find the term containing x^{15} .

SOLUTION

Since $(x^2 + 2x)^{11} = \left[x^2 \left(1 + \frac{2}{x} \right) \right]^{11} = x^{22} \left(1 + \frac{2}{x} \right)^{11}$, every term of the series expanded from $\left(1 + \frac{2}{x} \right)^{11}$ will be multiplied by x^{22} .

Hence, the term sought is that which contains $\left(\frac{2}{x} \right)^7$, or $\frac{2^7}{x^7}$; that is, the $(7 + 1)$ th, or 8th term.

$$\text{8th term} = x^{22} \frac{11 \cdot 10 \cdot 9 \cdot 8}{1 \cdot 2 \cdot 3 \cdot 4} \left(\frac{2}{x} \right)^7 = 42240 x^{15}.$$

3. Find the 4th term of $(a + 2)^{10}$.
4. Find the 4th term of $(x - 3y)^{12}$.
5. Find the 8th term of $(x + y)^{10}$.
6. Find the 5th term of $(x - 2y)^{12}$.
7. Find the 3d term of $(a^2 - a^{-2})^4$.
8. Find the 20th term of $(1 + x)^{24}$.
9. Find the 16th term of $(1 - 2x)^{20}$.
10. Find the middle term of $(a + 3b)^6$.
11. Find the 6th term of $\left(x + \frac{1}{x} \right)^{10}$.
12. Find the middle term of $\left(\frac{x}{y} - \frac{y}{x} \right)^{10}$.
13. Find the two middle terms of $\left(\frac{a}{b} - \frac{b}{a} \right)^9$.
14. Find the coefficient of a^9 in the expansion of $(a^3 + a)^5$.

449. The formula given for the expansion of $(a + x)^n$ is true for all commensurable values of n , whether they are positive, negative, integral, or fractional, and the student will, therefore, be able to expand such expressions, but the proof for negative and fractional exponents will be deferred until page 431.

In the expansion of $(a + x)^n$, if n is negative or fractional, none of the binomial coefficients, $\frac{n(n-1)}{1 \cdot 2}$, $\frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3}$, etc., can become 0; consequently, when such exponents are given, the series developed can have no end.

EXAMPLES

1. Expand
- $(1-x)^{-1}$
- and find its
- $(r+1)$
- th term.

SOLUTION. — Substituting 1 for a , y for x , and -1 for n in (III),

$$\begin{aligned}(1-y)^{-1} &= 1^{-1} - (-1)1^{-2}y + \frac{-1(-2)}{1 \cdot 2}1^{-3}y^2 - \frac{-1(-2)(-3)}{1 \cdot 2 \cdot 3}1^{-4}y^3 + \dots \\ &= 1 + y + y^2 + y^3 + \dots\end{aligned}$$

The $(r+1)$ th term is evidently y^r .

Since $(1-y)^{-1} = \frac{1}{1-y}$, the above expansion of $(1-y)^{-1}$ may be verified by division.

2. Expand
- $(a+x)^{\frac{1}{2}}$
- to five terms and find the 10th term.

Expand to four terms:

- | | | |
|------------------------------------|---|-----------------------------|
| 3. $(a+b)^{\frac{2}{3}}$. | 8. $\sqrt{4+x}$. | 13. $(1+x)^{\frac{2}{3}}$. |
| 4. $(a+b)^{-\frac{1}{2}}$. | 9. $\sqrt{(9-x)^3}$. | 14. $(1+a)^{-1}$. |
| 5. $(a-b)^{\frac{1}{2}}$. | 10. $(a^{\frac{2}{3}} - x^{-1})^{\frac{3}{2}}$. | 15. $(1-a)^{-1}$. |
| 6. $\sqrt[4]{(a-b)^3}$. | 11. $(a^{\frac{1}{2}} - x^{\frac{1}{3}})^{-6}$. | 16. $(1-x)^{-2}$. |
| 7. $\frac{1}{\sqrt[4]{(a-b)^3}}$. | 12. $\left(\frac{1}{\sqrt{a} - \sqrt[3]{x}}\right)^3$. | 17. $(1-x)^{-3}$. |

18. Find the
- $(r+1)$
- th term of
- $(a+x)^{\frac{1}{2}}$
- .

19. Show that

$$(1-x-x^2)^{-1} = 1 + x + 2x^2 + 3x^3 + 5x^4 + 8x^5 + 13x^6 + 21x^7 + \dots$$

20. Find the square root of 24 to three decimal places.

SOLUTION. $\sqrt{24} = (24)^{\frac{1}{2}} = (25-1)^{\frac{1}{2}} = (25)^{\frac{1}{2}}(1-\frac{1}{25})^{\frac{1}{2}} = 5(1-\frac{1}{25})^{\frac{1}{2}}$.

$$\begin{aligned}&5\left(1-\frac{1}{25}\right)^{\frac{1}{2}} \\ &= 5\left[1 - \frac{1}{2}\left(\frac{1}{25}\right) + \frac{\frac{1}{2}(-\frac{1}{2})}{1 \cdot 2}\left(\frac{1}{25}\right)^2 - \frac{\frac{1}{2}(-\frac{1}{2})(-\frac{3}{2})}{1 \cdot 2 \cdot 3}\left(\frac{1}{25}\right)^3 + \dots\right] \\ &= 5 - .1 - .001 - .00002 - \dots = 4.89898 - = 4.899, \text{ nearly.}\end{aligned}$$

Find the values of the following to three decimal places:

- | | | |
|-------------------|----------------------|----------------------|
| 21. $\sqrt{5}$. | 23. $\sqrt{26}$. | 25. $\sqrt[3]{9}$. |
| 22. $\sqrt{17}$. | 24. $\sqrt[3]{25}$. | 26. $\sqrt[5]{30}$. |

LOGARITHMS

450. 1. What power of 3 is 9? 27? 81? 243? 729?

2. What power of 5 is 25? 125? 625? 3125? 5? 1? $\frac{1}{5}$?

3. Express 100 as a power of 10; 1000 as a power of 10; 10,000 as a power of 10; 10 as a power of 10; 1 as a power of 10.

451. The exponent of the power to which a fixed number called the *Base* must be raised in order to produce a given number is called the **Logarithm** of the given number.

When 10 is the base, the logarithm of 100 is 2, for $100 = 10^2$; the logarithm of 1000 is 3, for $1000 = 10^3$; the logarithm of 10,000 is 4, for $10,000 = 10^4$.

452. When a is the base, x the exponent, and m the given number, x is the logarithm of the number m to the base a .

It is written $\log_a m = x$.

When the base is 10, it is not indicated.

Thus, the logarithm of 100 to the base 10 is 2. It is written $\log 100 = 2$.

453. Logarithms may be computed with any arithmetical number except unity as a base, but the base of the **Common** or **Briggs System** of logarithms is 10.

Since $10^0 = 1$, the logarithm of 1 is 0.

Since $10^1 = 10$, the logarithm of 10 is 1.

Since $10^2 = 100$, the logarithm of 100 is 2.

Since $10^3 = 1000$, the logarithm of 1000 is 3.

Since $10^{-1} = \frac{1}{10}$, the logarithm of .1 is -1 .

Since $10^{-2} = \frac{1}{100}$, the logarithm of .01 is -2 .

454. It is evident, then, that the logarithm of any number between 1 and 10 is a number greater than 0 and less than 1. For example, the logarithm of 4 is approximately 0.6021. Again, the logarithm of any number between 10 and 100 is a number greater than 1 and less than 2. For example, the logarithm of 50 is approximately 1.6990.

Most logarithms are incommensurable numbers. All the laws established for commensurable exponents apply also to incommensurable exponents, but the proofs have been omitted as being too difficult for the beginner.

455. The integral part of a logarithm is called the **Characteristic**; the fractional or decimal part, the **Mantissa**.

In $\log 50 = 1.6990$, the characteristic is 1 and the mantissa .6990.

456. The following examples will illustrate the *characteristic* and *mantissa*, and their significance:

$$\log 4580 = 3.6609; \text{ that is, } 4580 = 10^{3.6609}.$$

$$\log 458.0 = 2.6609; \text{ that is, } 458.0 = 10^{2.6609}.$$

$$\log 45.80 = 1.6609; \text{ that is, } 45.80 = 10^{1.6609}.$$

$$\log 4.580 = 0.6609; \text{ that is, } 4.580 = 10^{0.6609}.$$

$$\log .4580 = \bar{1}.6609; \text{ that is, } .4580 = 10^{-1+.6609}.$$

$$\log .0458 = \bar{2}.6609; \text{ that is, } .0458 = 10^{-2+.6609}.$$

$$\log .00458 = \bar{3}.6609; \text{ that is, } .00458 = 10^{-3+.6609}.$$

From the above examples it is evident that:

457. PRINCIPLES. — 1. *The characteristic of the logarithm of a number greater than 1 is positive and 1 less than the number of digits in its integral part.*

2. *The characteristic of the logarithm of a decimal is negative and numerically 1 greater than the number of ciphers immediately following the decimal point.*

458. To avoid writing a negative characteristic before a positive mantissa, it is customary to add 10 or some multiple of 10 to the negative characteristic, and to indicate that the number added is to be subtracted from the whole logarithm.

Thus, $\bar{1}.6609$ is written $9.6609 - 10$; $\bar{2}.3010$ is written $8.3010 - 10$; $\bar{14}.9031$ is written $6.9031 - 20$; $\bar{28}.8062$ is written $2.8062 - 30$; etc.

459. It is evident, also, from the examples, that in the logarithms of numbers expressed by the same figures in the same order, the decimal parts, or *mantissas*, are the same, and that the logarithms differ only in their *characteristics*. Hence, tables of logarithms contain only the *mantissas*.

460. The table of logarithms on the two following pages gives the decimal parts, or mantissas, correct to four places, for the common logarithms of all numbers from 1 to 1000.

461. To find the logarithm of a number.

EXAMPLES

1. Find the logarithm of 765.

SOLUTION. — In the following table the letter **N** designates a vertical column of numbers from 10 to 99 inclusive, and also a horizontal row of figures 0, 1, 2, 3, 4, 5, 6, 7, 8, 9. The first two figures of 765 appear as the number 76 in the vertical column marked **N** on page 409, and the third figure 5 in the horizontal row marked **N**.

In the same horizontal row as 76 are found the mantissas of the logarithms of the numbers 760, 761, 762, 763, 764, 765, etc. The mantissa of the logarithm of 765 is found in this row under 5, the third figure of 765. It is 8837 and means .8837.

By Prin. 1, the characteristic of the logarithm of 765 is 2.

Hence, the logarithm of 765 is 2.8837.

2. Find the logarithm of 4.

SOLUTION. — Although the numbers in the table appear to begin with 100, the table really includes all numbers from 1 to 1000, since numbers expressed by less than three figures may be expressed by three figures by adding decimal ciphers. Since $4 = 4.00$, and since, § 459, the mantissa of the logarithm of 4.00 is the same as that of 400, which is .6021, the mantissa of the logarithm of 4 is .6021.

By Prin. 1, the characteristic of the logarithm of 4 is 0.

Therefore, the logarithm of 4 is 0.6021.

Verify the following from the table:

$$3. \log 10 = 1.0000.$$

$$9. \log .2 = 9.3010 - 10.$$

$$4. \log 100 = 2.0000.$$

$$10. \log 542 = 2.7340.$$

$$5. \log 110 = 2.0414.$$

$$11. \log 345 = 2.5378.$$

$$6. \log 2 = 0.3010.$$

$$12. \log 5.07 = 0.7050.$$

$$7. \log 20 = 1.3010.$$

$$13. \log 78.5 = 1.8949.$$

$$8. \log 200 = 2.3010.$$

$$14. \log .981 = 9.9917 - 10.$$

TABLE OF COMMON LOGARITHMS

N	0	1	2	3	4	5	6	7	8	9
10	0000	0043	0086	0128	0170	0212	0253	0294	0334	0374
11	0414	0453	0492	0531	0569	0607	0645	0682	0719	0755
12	0792	0828	0864	0899	0934	0969	1004	1038	1072	1106
13	1139	1173	1206	1239	1271	1303	1335	1367	1399	1430
14	1461	1492	1523	1553	1584	1614	1644	1673	1703	1732
15	1761	1790	1818	1847	1875	1903	1931	1959	1987	2014
16	2041	2068	2095	2122	2148	2175	2201	2227	2253	2279
17	2304	2330	2355	2380	2405	2430	2455	2480	2504	2529
18	2553	2577	2601	2625	2648	2672	2695	2718	2742	2765
19	2788	2810	2833	2856	2878	2900	2923	2945	2967	2989
20	3010	3032	3054	3075	3096	3118	3139	3160	3181	3201
21	3222	3243	3263	3284	3304	3324	3345	3365	3385	3404
22	3424	3444	3464	3483	3502	3522	3541	3560	3579	3598
23	3617	3636	3655	3674	3692	3711	3729	3747	3766	3784
24	3802	3820	3838	3856	3874	3892	3909	3927	3945	3962
25	3979	3997	4014	4031	4048	4065	4082	4099	4116	4133
26	4150	4166	4183	4200	4216	4232	4249	4265	4281	4298
27	4314	4330	4346	4362	4378	4393	4409	4425	4440	4456
28	4472	4487	4502	4518	4533	4548	4564	4579	4594	4609
29	4624	4639	4654	4669	4683	4698	4713	4728	4742	4757
30	4771	4786	4800	4814	4829	4843	4857	4871	4886	4900
31	4914	4928	4942	4955	4969	4983	4997	5011	5024	5038
32	5051	5065	5079	5092	5105	5119	5132	5145	5159	5172
33	5185	5198	5211	5224	5237	5250	5263	5276	5289	5302
34	5315	5328	5340	5353	5366	5378	5391	5403	5416	5428
35	5441	5453	5465	5478	5490	5502	5514	5527	5539	5551
36	5563	5575	5587	5599	5611	5623	5635	5647	5658	5670
37	5682	5694	5705	5717	5729	5740	5752	5763	5775	5786
38	5798	5809	5821	5832	5843	5855	5866	5877	5888	5899
39	5911	5922	5933	5944	5955	5966	5977	5988	5999	6010
40	6021	6031	6042	6053	6064	6075	6085	6096	6107	6117
41	6128	6138	6149	6160	6170	6180	6191	6201	6212	6222
42	6232	6243	6253	6263	6274	6284	6294	6304	6314	6325
43	6335	6345	6355	6365	6375	6385	6395	6405	6415	6425
44	6435	6444	6454	6464	6474	6484	6493	6503	6513	6522
45	6532	6542	6551	6561	6571	6580	6590	6599	6609	6618
46	6628	6637	6646	6656	6665	6675	6684	6693	6702	6712
47	6721	6730	6739	6749	6758	6767	6776	6785	6794	6803
48	6812	6821	6830	6839	6848	6857	6866	6875	6884	6893
49	6902	6911	6920	6928	6937	6946	6955	6964	6972	6981
50	6990	6998	7007	7016	7024	7033	7042	7050	7059	7067
51	7076	7084	7093	7101	7110	7118	7126	7135	7143	7152
52	7160	7168	7177	7185	7193	7202	7210	7218	7226	7235
53	7243	7251	7259	7267	7275	7284	7292	7300	7308	7316
54	7324	7332	7340	7348	7356	7364	7372	7380	7388	7396
N	0	1	2	3	4	5	6	7	8	9

TABLE OF COMMON LOGARITHMS

N	0	1	2	3	4	5	6	7	8	9
55	7404	7412	7419	7427	7435	7443	7451	7459	7466	7474
56	7482	7490	7497	7505	7513	7520	7528	7536	7543	7551
57	7559	7566	7574	7582	7589	7597	7604	7612	7619	7627
58	7634	7642	7649	7657	7664	7672	7679	7686	7694	7701
59	7709	7716	7723	7731	7738	7745	7752	7760	7767	7774
60	7782	7789	7796	7803	7810	7818	7825	7832	7839	7846
61	7853	7860	7868	7875	7882	7889	7896	7903	7910	7917
62	7924	7931	7938	7945	7952	7959	7966	7973	7980	7987
63	7993	8000	8007	8014	8021	8028	8035	8041	8048	8055
64	8062	8069	8075	8082	8089	8096	8102	8109	8116	8122
65	8129	8136	8142	8149	8156	8162	8169	8176	8182	8189
66	8195	8202	8209	8215	8222	8228	8235	8241	8248	8254
67	8261	8267	8274	8280	8287	8293	8299	8306	8312	8319
68	8325	8331	8338	8344	8351	8357	8363	8370	8376	8382
69	8388	8395	8401	8407	8414	8420	8426	8432	8439	8445
70	8451	8457	8463	8470	8476	8482	8488	8494	8500	8506
71	8513	8519	8525	8531	8537	8543	8549	8555	8561	8567
72	8573	8579	8585	8591	8597	8603	8609	8615	8621	8627
73	8633	8639	8645	8651	8657	8663	8669	8675	8681	8686
74	8692	8698	8704	8710	8716	8722	8727	8733	8739	8745
75	8751	8756	8762	8768	8774	8779	8785	8791	8797	8802
76	8808	8814	8820	8825	8831	8837	8842	8848	8854	8859
77	8865	8871	8876	8882	8887	8893	8899	8904	8910	8915
78	8921	8927	8932	8938	8943	8949	8954	8960	8965	8971
79	8976	8982	8987	8993	8998	9004	9009	9015	9020	9025
80	9031	9036	9042	9047	9053	9058	9063	9069	9074	9079
81	9085	9090	9096	9101	9106	9112	9117	9122	9128	9133
82	9138	9143	9149	9154	9159	9165	9170	9175	9180	9186
83	9191	9196	9201	9206	9212	9217	9222	9227	9232	9238
84	9243	9248	9253	9258	9263	9269	9274	9279	9284	9289
85	9294	9299	9304	9309	9315	9320	9325	9330	9335	9340
86	9345	9350	9355	9360	9365	9370	9375	9380	9385	9390
87	9395	9400	9405	9410	9415	9420	9425	9430	9435	9440
88	9445	9450	9455	9460	9465	9469	9474	9479	9484	9489
89	9494	9499	9504	9509	9513	9518	9523	9528	9533	9538
90	9542	9547	9552	9557	9562	9566	9571	9576	9581	9586
91	9590	9595	9600	9605	9609	9614	9619	9624	9628	9633
92	9638	9643	9647	9652	9657	9661	9666	9671	9675	9680
93	9685	9689	9694	9699	9703	9708	9713	9717	9722	9727
94	9731	9736	9741	9745	9750	9754	9759	9763	9768	9773
95	9777	9782	9786	9791	9795	9800	9805	9809	9814	9818
96	9823	9827	9832	9836	9841	9845	9850	9854	9859	9863
97	9868	9872	9877	9881	9886	9890	9894	9899	9903	9908
98	9912	9917	9921	9926	9930	9934	9939	9943	9948	9952
99	9956	9961	9965	9969	9974	9978	9983	9987	9991	9996
N	0	1	2	3	4	5	6	7	8	9

15. Find the logarithm of 6253.

SOLUTION. — Since the table contains the mantissas not only of the logarithms of numbers expressed by three figures, but also of logarithms expressed by four figures when the last figure is 0, the mantissa of the logarithm of 625 is first found, since, § 459, it is also the mantissa of the logarithm of 6250. It is found to be .7959.

The next greater mantissa found in the table is .7966, which is the mantissa of the logarithm of 626 or of 6260. Since the numbers 6250 and 6260 differ by 10, and the mantissas of their logarithms differ by 7 ten-thousandths, it may be assumed as sufficiently accurate that each increase of 1 unit, as 6250 increases to 6260, produces a corresponding increase of .1 of 7 ten-thousandths in the mantissa of the logarithm. Consequently, 3 added to 6250 will add .3 of 7 ten-thousandths, or 2 ten-thousandths, to the mantissa of the logarithm of 6250 for the mantissa of the logarithm of 6253.

Hence, the mantissa of the logarithm of 6253 is .7959 + .0002, or .7961.

Since the number is an integer expressed by 4 digits, the characteristic is 3 (Prin. 1).

Therefore, the logarithm of 6253 is 3.7961.

NOTE. — The difference between two successive mantissas in the table is called the *Tabular Difference*.

Find the logarithm of

16. 1054.	20. 21.09.	24. .09095.
17. 1272.	21. 3.060.	25. .10125.
18. .0165.	22. 441.1.	26. 54.675.
19. 1906.	23. .7854.	27. .09885.

462. To find a number whose logarithm is given.

EXAMPLES

1. Find the number whose logarithm is 0.9472.

SOLUTION. — The two mantissas adjacent to the given mantissa are .9469 and .9474, corresponding to the numbers 8.85 and 8.86, since the given characteristic is 0. The given mantissa is 3 ten-thousandths greater than the mantissa of the logarithm of 8.85, and the mantissa of the logarithm of 8.86 is 5 ten-thousandths greater than the mantissa of the logarithm of 8.85.

Since the numbers 8.85 and 8.86 differ by 1 one-hundredth, and the mantissas of their logarithms differ by 5 ten-thousandths, it may be assumed as sufficiently accurate that each increase of 1 ten-thousandth in the mantissa is produced by an increase of $\frac{1}{5}$ of 1 one-hundredth in the number. Conse-

quently, an increase of 3 ten-thousandths in the mantissa is produced by an increase of $\frac{3}{1000}$ of 1 one-hundredth, or .006, in the number.

Hence, the number whose logarithm is 0.9472 is 8.856.

2. Find the number whose logarithm is 9.4180 — 10.

SOLUTION. — Given mantissa, .4180

Mantissa next less, .4166 ; figures corresponding, 261.

Difference, 14

'Tabular difference, 17)14(.8

Hence, the figures corresponding to the given mantissa are 2618.

Since the characteristic is 9 — 10, or — 1, the number is a decimal with no ciphers immediately following the decimal point.

Hence, the number whose logarithm is 9.4180 — 10 is .2618.

Find the number corresponding to

3. 0.3010. 8. 3.9545. 13. 9.3685 — 10.

4. 1.6021. 9. 0.8794. 14. 8.9932 — 10.

5. 2.9031. 10. 2.9371. 15. 8.9535 — 10.

6. 1.6669. 11. 0.8294. 16. 7.7168 — 10.

7. 2.7971 12. 1.9039. 17. 6.7016 — 10.

463. Multiplication by logarithms.

Since logarithms are the exponents of the powers to which a constant number is to be raised, it follows that:

464. PRINCIPLE. — *The logarithm of the product of two or more numbers is equal to the sum of their logarithms; that is,*

To any base, $\log(mn) = \log m + \log n$.

The above principle may be established as follows:

Let $\log_a m = x$ and $\log_a n = y$, a being any base.

It is to be proved that $\log_a(mn) = x + y$.

§ 451, $a^x = m$,

and $a^y = n$.

Multiplying, § 240, $a^{x+y} = mn$.

Hence, § 452, $\log_a(mn) = x + y$
 $= \log_a m + \log_a n$.

EXAMPLES

1. Multiply .0381 by 77.

SOLUTION

$$\text{Prin.,} \quad \log (.0381 \times 77) = \log .0381 + \log 77$$

$$\log .0381 = 8.5809 - 10$$

$$\log 77 = 1.8865$$

$$\text{Sum of logs} = 10.4674 - 10$$

$$= 0.4674$$

$$0.4674 = \log 2.934.$$

$$\therefore .0381 \times 77 = 2.934.$$

NOTE. — Three figures of a number corresponding to a logarithm may be found from this table with absolute accuracy, and in most cases the fourth will be correct. In finding the logarithms of numbers or the numbers corresponding to logarithms, allowance should be made for the figures after the fourth, whenever they express .5 or more.

Multiply :

2. 3.8 by 56.

6. 2.26 by 85.

10. 1.414 by 2.829.

3. 72 by 39.

7. 7.25 by 240.

11. 42.37 by .236.

4. 8.5 by 6.2.

8. 3272 by 75.

12. 2912 by .7281.

5. 1.64 by 35.

9. .892 by .805.

13. 289 by .7854.

465. Division by logarithms.

Since the logarithms of two numbers to a common base represent exponents of the same number, it follows that:

466. PRINCIPLE. — *The logarithm of the quotient of two numbers is equal to the logarithm of the dividend minus the logarithm of the divisor; that is,*

$$\text{To any base,} \quad \log (m \div n) = \log m - \log n.$$

The above principle may be established as follows:

$$\text{Let} \quad \log_a m = x \text{ and } \log_a n = y, \text{ } a \text{ being any base.}$$

$$\text{It is to be proved that} \quad \log_a (m \div n) = x - y.$$

$$\S 451, \quad a^x = m \text{ and } a^y = n.$$

$$\text{Dividing, } \S 248, \quad a^{x-y} = m \div n.$$

$$\text{Hence, } \S 452, \quad \log_a (m \div n) = x - y$$

$$= \log_a m - \log_a n.$$

EXAMPLES

1. Divide .00468 by 75.

SOLUTION

$$\begin{aligned} \text{Prin.,} \quad & \log (.00468 \div 75) = \log .00468 - \log 75. \\ & \log .00468 = 7.6702 - 10 \\ & \log 75 = 1.8751 \\ & \hline \text{Difference of logs} & = 5.7951 - 10 \\ & 5.7951 - 10 = \log .00006239. \\ \therefore .00468 \div 75 & = .00006239. \end{aligned}$$

2. Divide 12.4 by 16.

SOLUTION

$$\begin{aligned} \text{Prin.,} \quad & \log (12.4 \div 16) = \log 12.4 - \log 16. \\ & \log 12.4 = 1.0934 = 11.0934 - 10 \\ & \log 16 = 1.2041 \\ & \hline \text{Difference of logs} & = 9.8893 - 10 \\ & 9.8893 - 10 = \log .775. \\ \therefore 12.4 \div 16 & = .775. \end{aligned}$$

SUGGESTION. — The positive part of the logarithm of the dividend may be made to exceed that of the divisor, if necessary, by adding 10 — 10 or 20 — 20, etc.

Divide:

- | | | |
|------------------|--------------------|--------------------|
| 3. 3025 by 55. | 8. 10 by 3.14. | 13. 1 by 40. |
| 4. 4096 by 32. | 9. .6911 by .7854. | 14. 1 by 75. |
| 5. 3249 by 57. | 10. 2.816 by 22.5. | 15. 200 by .5236. |
| 6. .2601 by .68. | 11. 4 by .00521. | 16. 300 by 17.32. |
| 7. 3950 by .250. | 12. 26 by .06771. | 17. .220 by .3183. |

467. Extended operations in multiplication and division.

Since dividing by a number is equivalent to multiplying by its reciprocal, for every operation of division an operation of multiplication may be substituted. In extended operations in multiplication and division with the aid of logarithms, the latter method of dividing is the more convenient.

468. The logarithm of the reciprocal of a number is called the **Cologarithm** of the number.

The cologarithm of 100 is equal to the logarithm of $\frac{1}{100}$, which is -2 .
The cologarithm 100 is -2 is abbreviated, $\text{colog } 100 = -2$.

469. Since the logarithm of 1 is 0 and the logarithm of a quotient is obtained by subtracting the logarithm of the divisor from that of the dividend, it is evident that the cologarithm of a number is 0 minus the logarithm of the number, or the logarithm of the number with the sign of the logarithm changed; that is, if $\log_a m = x$; then, $\text{colog}_a m = -x$.

Since subtracting a number is equivalent to adding it with its sign changed, it follows that:

470. PRINCIPLE. — *Instead of subtracting the logarithm of the divisor from the logarithm of the dividend, the cologarithm of the divisor may be added to the logarithm of the dividend; that is,*

To any base, $\log(m \div n) = \log m + \text{colog } n$.

EXAMPLES

1. Find the value of $\frac{.063 \times 58.5 \times 799}{458 \times 15.6 \times .029}$ by logarithms.

SOLUTION

$$\frac{.063 \times 58.5 \times 799}{458 \times 15.6 \times .029} = .063 \times 58.5 \times 799 \times \frac{1}{458} \times \frac{1}{15.6} \times \frac{1}{.029}$$

$$\log .063 = 8.7993 - 10$$

$$\log 58.5 = 1.7672$$

$$\log 799 = 2.9025$$

$$\text{colog } 458 = 7.3391 - 10$$

$$\text{colog } 15.6 = 8.8069 - 10$$

$$\text{colog } .029 = 1.5376$$

$$\log \text{ of result} = 31.1526 - 30$$

$$= 1.1526.$$

$$\therefore \text{ result} = 14.21.$$

Find the value of

$$2. \frac{110 \times 3.1 \times .653}{33 \times 7.854 \times 1.7}$$

$$5. \frac{15 \times .37 \times 26.16}{11 \times 8 \times .18 \times 6.67}$$

$$3. \frac{6000 \times 5 \times 29}{.7854 \times 25000 \times 81.7}$$

$$6. \frac{78 \times 52 \times 1605}{338 \times 767 \times 431}$$

$$4. \frac{3.516 \times 485 \times 65}{3.33 \times 17 \times 18 \times 73}$$

$$7. \frac{.5 \times .315 \times 428}{.317 \times .973 \times 43.7}$$

471. Involution by logarithms.

Since logarithms are simply exponents, it follows that:

472. PRINCIPLE. — *The logarithm of a power of a number is equal to the logarithm of the number multiplied by the index of the power; that is,*

To any base, $\log m^n = n \log m$.

The above principle may be established as follows:

Let $\log_a m = x$, and let n be any number, a being any base.

It is to be proved that $\log_a m^n = nx$.

§ 451, $a^x = m$.

Raising each member to the n th power,

Ax. 6 and § 249, $a^{nx} = m^n$.

Hence, § 452, $\log_a m^n = nx = n \log_a m$.

EXAMPLES

1. Find the value of $.25^2$.

SOLUTION

Prin., $\log .25^2 = 2 \log .25$.

$\log .25 = 9.3979 - 10$.

$2 \log .25 = 18.7958 - 20$

$= 8.7958 - 10$.

$8.7958 - 10 = \log .06249$.

$\therefore .25^2 = .06249$.

NOTE. — By actual multiplication it is found that $.25^2 = .0625$, whereas the result obtained by the use of the table is $.06249$.

Also, by multiplication, $18^2 = 324$, whereas by the use of the table it is found to be 324.1. Such inaccuracies must be expected when a four-place table is used.

Find by logarithms the value of

- | | | | |
|--------------|----------------|-----------------|------------------------------|
| 2. 7^2 . | 7. $.78^2$. | 12. 4.07^2 . | 17. $(\frac{3}{20})^2$. |
| 3. 11^2 . | 8. 8.05^2 . | 13. $.543^2$. | 18. $(\frac{1}{7})^2$. |
| 4. 47^2 . | 9. 8.33^2 . | 14. 7^4 . | 19. $(\frac{128}{1738})^2$. |
| 5. 4.9^2 . | 10. 6.61^2 . | 15. 1.02^5 . | 20. $(\frac{675}{4121})^2$. |
| 6. 5.2^2 . | 11. $.714^2$. | 16. 1.738^2 . | 21. $(\frac{1}{248})^4$. |

473. Evolution by logarithms.

Since logarithms are simply exponents, it follows that :

474. PRINCIPLE. — *The logarithm of a root of a number is equal to the logarithm of the number divided by the index of the required root; that is,*

$$\text{To any base,} \quad \log \sqrt[n]{m} = \frac{\log m}{n}.$$

The above principle may be established as follows :

Let $\log_a m = x$, and let n be any number, a being any base.

It is to be proved that $\log_a \sqrt[n]{m} = x + n$.

$$\S 451, \quad a^x = m.$$

Taking the n th root of each member,

$$\text{Ax. 7 and } \S 250, \quad a^{x+n} = \sqrt[n]{m}.$$

$$\text{Hence, } \S 452, \quad \log_a \sqrt[n]{m} = x + n = \frac{\log_a m}{n}.$$

EXAMPLES

1. Find the square root of .1296 by logarithms.

SOLUTION

$$\begin{aligned} \text{Prin.,} \quad & \log \sqrt{.1296} = \frac{1}{2} \log .1296. \\ & \log .1296 = 9.1126 - 10. \\ & \quad \quad \quad \begin{array}{r} 2 \overline{) 19.1126 - 20} \\ \underline{9.5563 - 10} \end{array} \\ & 9.5563 - 10 = \log .360. \\ & \therefore \sqrt{.1296} = .36. \end{aligned}$$

Find by logarithms the value of

- | | | | |
|----------------------------|------------------------------|---------------------|------------------------------|
| 2. $225^{\frac{1}{2}}$. | 9. $1331^{\frac{1}{3}}$. | 16. $\sqrt{2}$. | 23. $\sqrt[3]{2}$. |
| 3. $12.25^{\frac{1}{2}}$. | 10. $1024^{\frac{1}{5}}$. | 17. $\sqrt{3}$. | 24. $\sqrt[3]{.027}$. |
| 4. $.2025^{\frac{1}{2}}$. | 11. $.6724^{\frac{1}{3}}$. | 18. $\sqrt{5}$. | 25. $\sqrt{30\frac{1}{4}}$. |
| 5. $324^{\frac{1}{2}}$. | 12. $5.929^{\frac{1}{3}}$. | 19. $\sqrt{6}$. | 26. $\sqrt{.90}$. |
| 6. $.512^{\frac{1}{2}}$. | 13. $.4624^{\frac{1}{3}}$. | 20. $\sqrt[3]{2}$. | 27. $\sqrt{.52}$. |
| 7. $.1181^{\frac{1}{2}}$. | 14. $1.4641^{\frac{1}{3}}$. | 21. $\sqrt[4]{4}$. | 28. $\sqrt[5]{.032}$. |
| 8. $3.375^{\frac{1}{2}}$. | 15. $.00032^{\frac{1}{5}}$. | 22. $\sqrt[3]{3}$. | 29. $\sqrt{.025}$. |

Simplify the following:

- | | |
|--|--|
| 30. $\frac{176}{15 \times 3.1416}$. | 35. $\frac{14.5\sqrt[3]{1.6}}{11}$. |
| 31. $\frac{100^2}{48 \times 64 \times 11}$. | 36. $\sqrt{\frac{.434 \times 96^4}{64 \times 1500}}$. |
| 32. $\frac{52^2 \times 300}{12 \times .31225 \times 400000}$. | 37. $\frac{.32 \times 5000 \times 18}{3.14 \times .1222 \times 8}$. |
| 33. $\sqrt{\frac{400}{55 \times 3.1416}}$. | 38. $\frac{11 \times 2.63 \times 4.263}{48 \times 3.263}$. |
| 34. $50 \times \frac{2^{2.5}}{8^{1.63}}$. | 39. $\sqrt{\frac{3500}{1.06^5}}$. |
| 40. $2^{\frac{1}{2}} \times (\frac{1}{2})^{\frac{1}{3}} \times \sqrt[3]{\frac{3}{2}} \times \sqrt{.1}$. | |

475. Logarithms applied to the solution of problems in compound interest and annuities.

1. What is the amount of \$1 for 1 year at 6%? By what must the amount for 1 year be multiplied to find the amount for 2 years at 6% compound interest?

2. By what must the amount for 2 years be multiplied to obtain the amount for 3 years, compound interest? By what must the amount for 3 years be multiplied to obtain the amount for 4 years, compound interest?

3. Since the amount of any principal at 6% compound interest for 1 year is 1.06 times the principal; for two years, 1.06×1.06 , or 1.06^2 times the principal; for 3 years, $1.06 \times 1.06 \times 1.06$, or 1.06^3 times the principal, etc., what will be the amount (A) of any principal (P) for n years at any rate per cent (r)?

FORMULA. $A = P(1 + r)^n.$

Expressing the formula by logarithms,

$$\log A = \log P + n \log(1 + r). \quad (1)$$

$$\therefore \log P = \log A - n \log(1 + r); \quad (2)$$

also $\log(1 + r) = \frac{\log A - \log P}{n}, \quad (3)$

and $n = \frac{\log A - \log P}{\log(1 + r)}. \quad (4)$

EXAMPLES

1. What is the amount of \$475 for 10 years at 6% compound interest?

SOLUTION

$$A = P(1 + r)^n.$$

$$\log 475 = 2.6767$$

$$\log 1.06^{10} = 0.2530$$

$$\log A = 2.9297$$

$$\therefore A = \$850.60.$$

2. What will be the amount of \$225 loaned for 5 years at 8% compound interest?

3. Find the amount of \$700 loaned for 5 years at 6% compound interest.

4. Find the amount of \$400 for 10 years at 3% compound interest.

5. Find the amount of \$1200 for 20 years at 4% compound interest.

6. What principal will amount to \$1000 in 10 years at 5% compound interest?

7. What sum of money invested at 4% compound interest, payable semiannually, will amount to \$ 743 in 10 years ?

8. What principal loaned at 4% compound interest will amount to \$ 1500 in 10 years ?

9. What sum of money invested at 4% compound interest from a child's birth until he is 21 years old will yield \$ 1000 ?

10. In what time will \$ 800 amount to \$ 1834.50, if put at compound interest at 5% ?

11. What is the rate per cent when \$ 300 loaned at compound interest for 6 years amounts to \$ 402 ?

12. A man agreed to loan \$ 1000 at 6% compound interest for a time long enough for the principal to double itself. How long was the money at interest ?

476. A sum of money to be paid periodically for a given number of years, during the life of a person, or forever, is called an **Annuity**.

The payments may be made once a year, or twice or four times a year, etc.

Interest is allowed upon deferred payments.

477. To find the amount of an annuity left unpaid for a given number of years, compound interest being allowed.

1. If an annuity of a dollars is not paid at the end of the first year, how much is then due ?

2. Upon what sum will compound interest be computed for the second year ? What will be the amount of that sum, if the rate is r ? What will be the whole sum due at the end of the *second* year ?

Ans. $a + a(1 + r)$.

3. Upon what sum will compound interest be computed for the third year ? What will be the amount of that sum at the given rate ?

Ans. $a(1 + r) + a(1 + r)^2$.

What will be the whole sum due at the end of the *third* year ?

Ans. $a + a(1 + r) + a(1 + r)^2$.

4. What will be the whole sum due at the end of the *fourth* year? What will be the whole sum due at the end of the *n*th year?

478. Let a represent the annuity, n the number of years, r the rate, and A the whole amount due at the end of the n th year.

Then,

$$\begin{aligned} A &= a + a(1+r) + a(1+r)^2 + a(1+r)^3 + \cdots a(1+r)^{n-1} \\ &= a \{1 + (1+r) + (1+r)^2 + (1+r)^3 + \cdots + (1+r)^{n-1}\}. \end{aligned}$$

Since the terms of A form a geometrical progression in which $1+r$ is the ratio, § 370, the sum of the series is

$$A = \frac{a}{r} [(1+r)^n - 1].$$

479. Sometimes annuities, drawing interest, are not payable until after a certain number of years. It is often necessary, therefore, to find the present value of such annuities.

480. A sum that will amount to the value of an annuity, if put at interest at the given rate for the given time, is called the **Present Value** of the annuity.

481. 1. If P denotes the present value of an annuity due in n years, allowing $r\%$ compound interest, to what sum will P be equal in that time at the given rate? *Ans.* $P(1+r)^n$.

2. Since the amount of the present value put at interest for the given time at the given rate is equal to the amount of the annuity for the same time and rate, equate the two sums and find the value of P .

$$P(1+r)^n = \frac{a}{r} [(1+r)^n - 1].$$

$$\begin{aligned} \therefore P &= \frac{a}{r} \cdot \frac{(1+r)^n - 1}{(1+r)^n} \\ &= \frac{a}{r} \left(1 - \frac{1}{(1+r)^n} \right). \end{aligned}$$

EXAMPLES

1. What will be the amount of an annuity of \$100 remaining unpaid for 10 years at 6% compound interest?

SOLUTION.
$$A = \frac{a}{r} [(1+r)^n - 1].$$

By logarithms,

$$\begin{aligned} 1.06^{10} &= 1.7904 \\ \therefore 1.06^{10} - 1 &= .7904 \\ \log 100 &= 2.0000 \\ \log .7904 &= 9.8978 - 10 \\ \text{colog } .06 &= 1.2218 \\ \hline \therefore \log A &= 3.1196 \end{aligned}$$

Hence, $A = \$1317$, the amount of the annuity.

2. What is the present value of an annuity of \$100 to continue 10 years at 6% compound interest?

SOLUTION.
$$P = \frac{a}{r} \cdot \frac{(1+r)^n - 1}{(1+r)^n}.$$

By logarithms,

$$\begin{aligned} 1.06^{10} &= 1.7904 \\ \therefore 1.06^{10} - 1 &= .7904 \\ \log 100 &= 2.0000 \\ \log .7904 &= 9.8978 - 10 \\ \text{colog } .06 &= 1.2218 \\ \text{colog } 1.06^{10} &= 9.7470 - 10 \\ \hline \therefore \log P &= 2.8666 \end{aligned}$$

Hence, $P = \$735.50$, the p. v. of the annuity.

3. To what sum will an annuity of \$25 amount in 20 years at 4% compound interest?

4. What is the present value of an annuity of \$300 for 5 years at 4% compound interest?

5. What will be the amount of an annuity of \$17.76 remaining unpaid for 25 years at $3\frac{1}{2}\%$ compound interest?

6. What is the present value of an annuity of \$1000 to continue 20 years, allowing compound interest at $4\frac{1}{2}\%$?

7. What annuity will amount to \$1000 in 10 years at 5% compound interest?

UNDETERMINED COEFFICIENTS

482. By division, $\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$.

1. If $x = \frac{1}{2}$, what is the value of the fraction? What are the successive terms of the series, when $x = \frac{1}{2}$? What is the sum of the first 2 terms? of the first 3 terms? of the first 4 terms? What value does the sum of the first n terms approach as a limit? How, then, does the sum of the series compare with 2, the value of the fraction, when $x = \frac{1}{2}$?

2. If $x = 1$, what are the successive terms of the series? How large can the sum of n terms be? How, then, does the sum of the series compare with the value of the fraction when $x = 1$?

3. How does the sum of the series compare with the value of the fraction when $x = 2$? when $x = 3$?

4. If $x = -1$, what is the value of the fraction? What is the sum of the first n terms of the series when n is even? the sum of the first n terms of the series when n is odd? How, then, does the sum of the series compare with the value of the fraction when $x = -1$?

483. When the sum of the first n terms of an infinite series approaches a finite number as a limit, as n is indefinitely increased, the limit is called the *sum* of the series, and the series is called a **Convergent Series**.

The infinite series $1 + x + x^2 + x^3 + \dots$, which arises from the fraction $\frac{1}{1-x}$, is convergent if $x = \frac{1}{2}$ or any number numerically less than 1; for the sum of the first n terms, as n increases, approaches the value of the fraction as a limit.

484. When the sum of the first n terms of an infinite series can be made numerically greater than any finite number by taking n sufficiently great, the series is called a **Divergent Series**.

The infinite series $1 + x + x^2 + x^3 + \dots$ is divergent if $x = 1$ or any number numerically greater than 1, as 2, or -2 , or 3, etc. ; for the sum of the first n terms, as n increases indefinitely, becomes larger than any finite number.

When x is numerically greater than 1, the divergent series $1 + x + x^2 + x^3 + \dots$ is not equal to the fraction from which it arises.

485. When the sum of the first n terms of an infinite series oscillates between certain fixed values, the series is called an **Oscillating Series**.

The infinite series $1 + x + x^2 + x^3 + \dots$ oscillates between the values 1 and 0 when $x = -1$; for the sum is 1 if the number of terms is odd, and 0 if the number of terms is even. There is no number of terms for which the series is equal to the fraction from which it arises.

It is evident that an oscillating series is neither convergent nor divergent.

It is evident from the definitions given above, that generally only convergent series can be used in demonstrations and discussions involving infinite series.

486. Coefficients assumed in the demonstration of a principle or the solution of a problem, whose values, not known at the outset, are to be determined by subsequent processes, are called **Undetermined Coefficients**.

487. PRINCIPLE OF UNDETERMINED COEFFICIENTS. — *If two series arranged according to the ascending powers of x are equal for every value of x that makes both series convergent, the coefficients of the like powers of x are equal, each to each.*

Let $A + Bx + Cx^2 + Dx^3 + \dots = A' + B'x + C'x^2 + D'x^3 + \dots$ for every value of x that makes both series convergent.

It is to be proved that $A = A'$, $B = B'$, $C = C'$, etc.

Since when $x = 0$, the first series is equal to A and the second is equal to A' , each series is convergent for $x = 0$.

Since the series are equal for every value of x that makes each of them convergent, they are equal for $x = 0$.

Substituting 0 for x , $A = A'$.

Since A and A' are constants and equal, $A = A'$, whatever the value of x .

Hence, **AX. 3.** $Bx + Cx^2 + Dx^3 + \dots = B'x + C'x^2 + D'x^3 + \dots$.

Dividing by x , $B + Cx + Dx^2 + \dots = B + C'x + D'x^2 + \dots$.

Reasoning as before, $B = B'$; $C = C'$; $D = D'$; etc.

488. An algebraic expression is said to be *developed* when it is transformed into a series the sum of whose terms is equal to the given expression.

The series is called the *development* of the given expression.

When x is numerically less than 1, the development of the fraction $\frac{1}{1-x}$ is the infinite converging series $1 + x + x^2 + x^3 + \dots$.

DEVELOPMENT OF FRACTIONS

EXAMPLES

489. 1. Develop the fraction $\frac{1+2x}{1+x+x^2}$.

SOLUTION

Assume $\frac{1+2x}{1+x+x^2} = A + Bx + Cx^2 + Dx^3 + Ex^4 + \dots$

Clearing of fractions and collecting terms,

$$1 + 2x = A + B \left| \begin{array}{c} x + C \\ + A \end{array} \right| x^2 + D \left| \begin{array}{c} x^2 + E \\ + C \end{array} \right| x^3 + \dots$$

Equating coefficients of like powers of x , § 487, and observing that

$$1 + 2x = 1 + 2x + 0x^2 + 0x^3 + 0x^4 + \dots,$$

$$A = 1.$$

$$B + A = 2; \therefore B = 1.$$

$$C + B + A = 0; \therefore C = -2.$$

$$D + C + B = 0; \therefore D = 1.$$

$$E + D + C = 0; \therefore E = 1.$$

$$\therefore \frac{1+2x}{1+x+x^2} = 1 + x - 2x^2 + x^3 + x^4 - \dots$$

The fraction may be developed also by division.

The exponent of x in the first term of the series may always be determined by dividing the term in the numerator having the lowest power of x by the term in the denominator having the lowest power of x . Beginning with this power of x , a series may be assumed proceeding according to the ascending powers of x .

2. Develop the fraction $\frac{2-x+2x^2}{x^2-2x^3}$.

SOLUTION

Since the first term of the quotient is evidently $\frac{2}{x^2}$, or $2x^{-2}$,

assume $\frac{2-x+2x^2}{x^2-2x^3} = Ax^{-2} + Bx^{-1} + C + Dx + Ex^2 + \dots$

Clearing of fractions,

$$2-x+2x^2 = A + \frac{B}{x} + \frac{C}{x^2} + \frac{D}{x^3} + \frac{E}{x^4} + \dots$$

Equating the coefficients of like powers of x , § 487,

$$A = 2.$$

$$B - 2A = -1; \quad \therefore B = 3.$$

$$C - 2B = 2; \quad \therefore C = 8.$$

$$D - 2C = 0; \quad \therefore D = 16.$$

$$E - 2D = 0; \quad \therefore E = 32.$$

$$\therefore \frac{2-x+2x^2}{x^2-2x^3} = 2x^{-2} + 3x^{-1} + 8 + 16x + 32x^2 + \dots$$

Develop to five terms:

3. $\frac{1+x}{1-x}$

10. $\frac{1-x-2x^2}{1-2x-x^2}$

17. $\frac{2-5x}{2x-x^2}$

4. $\frac{1+3x}{1+x}$

11. $\frac{x-x^2}{1+2x-x^2}$

18. $\frac{1+x+x^2}{x+x^3+x^4}$

5. $\frac{1}{1-2x}$

12. $\frac{1-x}{1-x+x^2}$

19. $\frac{1}{a-x}$

6. $\frac{3}{2-x}$

13. $\frac{1}{1-x-x^2}$

20. $\frac{1}{a+x}$

7. $\frac{1}{1-ax}$

14. $\frac{2+x-2x^2}{1-x+2x^2}$

21. $\frac{a}{1-x}$

8. $\frac{2+3x^2}{1-2x^2}$

15. $\frac{x^2+x^3}{1-2x+x^2}$

22. $\frac{a}{b-x}$

9. $\frac{4x-3x^2}{1+2x}$

16. $\frac{1-2x}{x^2+x^3+x^4}$

23. $\frac{c}{b-ax}$

DEVELOPMENT OF SURDS

EXAMPLES

490. 1. Develop the expression $\sqrt{a+x}$ by the use of undetermined coefficients.

SOLUTION

$$\begin{array}{ll} \text{Assume} & \sqrt{a+x} = A + Bx + Cx^2 + Dx^3 + Ex^4 + \dots \\ \text{Squaring,} & a+x = A^2 + 2ABx + B^2x^2 + 2ADx^3 + C^2x^4 + 2BCx^3 + 2AEx^4 + 2BDx^5 + \dots \end{array}$$

Equating the coefficients of like powers of x , § 487,

$$A^2 = a; \therefore A = \sqrt{a}.$$

$$2AB = 1; \therefore B = \frac{\sqrt{a}}{2a}.$$

$$B^2 + 2AC = 0; \therefore C = -\frac{\sqrt{a}}{8a^2}.$$

$$2AD + 2BC = 0; \therefore D = \frac{\sqrt{a}}{16a^3}.$$

$$C^2 + 2AE + 2BD = 0; \therefore E = -\frac{5\sqrt{a}}{128a^4}.$$

$$\therefore \sqrt{a+x} = \sqrt{a} \left(1 + \frac{x}{2a} - \frac{x^2}{8a^2} + \frac{x^3}{16a^3} - \frac{5x^4}{128a^4} + \dots \right).$$

The given surd may also be developed by the extraction of the root indicated or by the use of the binomial formula.

Develop the following to five terms by the use of undetermined coefficients:

2. $\sqrt{1-x}$.

8. $\sqrt{1+x}$.

3. $\sqrt{1+2x}$.

9. $\sqrt{1+x+x^2}$.

4. $\sqrt{1+4x}$.

10. $(1+4x+6x^2+4x^3+x^4)^{\frac{1}{2}}$.

5. $\sqrt{4+x}$.

11. $(1-3x+3x^2-x^3)^{\frac{1}{2}}$.

6. $\sqrt{a-x}$.

12. $(1+x)^{\frac{3}{2}}$.

7. $\sqrt{a^2-x^2}$.

13. $\sqrt{1+2x+3x^2+4x^3+\dots}$.

PARTIAL FRACTIONS

491. To resolve a fraction into its partial fractions is to separate it into fractions whose sum is equal to the given fraction. Since this is the reverse of the process of uniting fractions with different denominators into a single fraction having their lowest common denominator, it is evident that the denominators of the partial fractions must comprise all the various factors of the given denominator. The proper numerators may be found by the method of undetermined coefficients.

EXAMPLES

1. Resolve $\frac{3}{1-5x+6x^2}$ into its partial fractions.

SOLUTION

Assume that $\frac{3}{1-5x+6x^2} = \frac{A}{1-3x} + \frac{B}{1-2x}$ is an identity.

Clearing of fractions, $3 = A - 2Ax + B - 3Bx.$

That is, $3 + 0x = A + B - (2A + 3B)x.$

\therefore § 487, $A + B = 3$ and $2A + 3B = 0.$

Solving, $A = 9$ and $B = -6.$

Hence, $\frac{3}{1-5x+6x^2} = \frac{9}{1-3x} - \frac{6}{1-2x}.$

2. Resolve $\frac{5-6x}{1-4x+4x^2}$ into its partial fractions.

SOLUTION

Assume that $\frac{5-6x}{1-4x+4x^2} = \frac{A}{1-2x} + \frac{B}{(1-2x)^2}$ is an identity.

Reducing, $5-6x = A + B - 2Ax.$

\therefore § 487, $A + B = 5$ and $-2A = -6.$

Solving, $A = 3$ and $B = 2.$

Hence, $\frac{5-6x}{1-4x+4x^2} = \frac{3}{1-2x} + \frac{2}{(1-2x)^2}$

3. Resolve $\frac{7-11x+7x^2}{(1-x)^3(1+x+x^2)}$ into its partial fractions.

SOLUTION

By the previous solution it is evident that the fractions corresponding to the factor $(1-x)^3$ will be $\frac{A}{1-x}$, $\frac{B}{(1-x)^2}$, and $\frac{C}{(1-x)^3}$.

Since the factor $(1+x+x^2)$ is quadratic and has no rational simple factor, the numerator corresponding to the denominator $1+x+x^2$ may have two terms; therefore, assume that

$$\frac{7-11x+7x^2}{(1-x)^3(1+x+x^2)} = \frac{A}{1-x} + \frac{B}{(1-x)^2} + \frac{C}{(1-x)^3} + \frac{D+Ex}{1+x+x^2} \quad (1)$$

is an identity.

$$\begin{aligned} \text{Then, } 7-11x+7x^2 &= A(1-x)^2(1+x+x^2) + B(1-x)(1+x+x^2) \\ &\quad + C(1+x+x^2) + (D+Ex)(1-x)^3 \end{aligned} \quad (2)$$

is an identity; that is, is true for all values of x .

Since there are five coefficients, A , B , C , D , and E , to be determined, and since (2) is true for all values of x , by giving x in succession each of five different values, five independent equations involving the undetermined coefficients may be formed, and from these equations the coefficients may be determined.

Let $x = 1$; then, $(1-x)$, $(1-x)^2$, and $(1-x)^3$ reduce to 0, and (2) becomes

$$3 = 3C; \therefore C = 1. \quad (3)$$

$$\text{Let } x = 0; \text{ then, } 7 = A + B + C + D,$$

$$\text{or, since } C = 1, \quad A + B + D = 6. \quad (4)$$

$$\text{Let } x = -1; \text{ then, } 25 = 4A + 2B + 1 + 8D - 8E,$$

$$\text{or, dividing by 2, } 2A + B + 4D - 4E = 12. \quad (5)$$

$$\text{Let } x = 2; \text{ then, } 13 = 7A - 7B + 7 - D - 2E,$$

$$\text{or } 7A - 7B - D - 2E = 6. \quad (6)$$

$$\text{Let } x = -2; \text{ then, } 57 = 27A + 9B + 3 + 27D - 54E,$$

$$\text{or, dividing by 9, } 3A + B + 3D - 6E = 6. \quad (7)$$

Solving the equations, (4), (5), (6), and (7), we have, together with (3),

$$A = 2, B = 0, C = 1, D = 4, E = 2.$$

$$\text{Hence, } \frac{7-11x+7x^2}{(1-x)^3(1+x+x^2)} = \frac{2}{1-x} + \frac{1}{(1-x)^3} + \frac{4+2x}{1+x+x^2}$$

4. Resolve $\frac{x+2x^2}{1-x^3}$ into its partial fractions.

SUGGESTION. — Assume $\frac{x+2x^2}{1-x^3} = \frac{A}{1-x} + \frac{B+Cx}{1+x+x^2}$.

5. Resolve $\frac{x^3+x-4}{x^2-1}$ into its partial fractions.

SUGGESTION. — When the numerator is not of lower degree than the denominator, the numerator should be divided by the denominator until the remainder is of lower degree than the denominator. The fractional part of the resulting mixed expression may then be resolved into partial fractions, and these may be annexed to the integral part.

Resolve each of the following into its partial fractions:

- | | |
|---|---|
| 6. $\frac{2x}{8-6x+x^2}$. | 12. $\frac{x^2-6x}{(x-5)^3}$. |
| 7. $\frac{3+4x}{1+8x+16x^2}$. | 13. $\frac{x^2-5}{x^2-1}$. |
| 8. $\frac{3x}{1+x^3}$. | 14. $\frac{2x^2+9x+11}{x^2+4x+4}$. |
| 9. $\frac{1-x-6x^2}{x-x^3}$. | 15. $\frac{2-6x+6x^2}{1-6x+11x^2-6x^3}$. |
| 10. $\frac{1-2x+2x^2}{(1-x)(1-2x)^2}$. | 16. $\frac{49}{(2-3x)^2(3-x)}$. |
| 11. $\frac{3x-2}{(x-3)^2}$. | 17. $\frac{1+2x+3x^2+2x^3}{x-x^5}$. |

REVERSION OF SERIES

492. To revert a series of ascending powers of x is to express the value of x by means of a series of ascending powers of some other letter, as y .

Let it be required to revert the series

$$y = ax + bx^2 + cx^3 + dx^4 + \dots, \quad (1)$$

in which x has any value that will render the series convergent.

Assume $x = Ay + By^2 + Cy^3 + Dy^4 + \dots$ (2)

Substituting this value of x in (1), and dropping all terms involving a higher power of y than the fourth,

$$y = aAy + aB \left| \begin{array}{c} y^2 + aC \\ + bA^2 \end{array} \right| y^3 + aD \left| \begin{array}{c} + bB^2 \\ + 2bAC \\ + 3cA^2B \\ + dA^4 \end{array} \right| y^4 + \dots$$

Since (1) is an identity, § 487,

$$aA = 1; \quad \therefore A = \frac{1}{a}.$$

$$aB + bA^2 = 0; \quad \therefore B = -\frac{bA^2}{a} = -\frac{b}{a^3}.$$

$$aC + 2bAB + cA^3 = 0; \quad \therefore C = \frac{-2bAB - cA^3}{a} = \frac{2b^2 - ac}{a^5}.$$

$$aD + bB^2 + 2bAC + 3cA^2B + dA^4 = 0; \quad \therefore D = -\frac{a^2d - 5abc + 5b^3}{a^7}.$$

Hence,

$$x = \frac{1}{a}y - \frac{b}{a^3}y^2 + \frac{2b^2 - ac}{a^5}y^3 - \frac{a^2d - 5abc + 5b^3}{a^7}y^4 + \dots \quad (3)$$

EXAMPLES

1. Revert the series $y = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$

SOLUTION.—Substituting in (3) 1 for a , $-\frac{1}{2}$ for b , $\frac{1}{3}$ for c , $-\frac{1}{4}$ for d , the values of the undetermined coefficients in (2) are found;

whence, $x = y + \frac{1}{2}y^2 + \frac{1}{3}y^3 + \frac{1}{24}y^4 + \dots$

Revert the series in the following equations:

$$2. y = x + x^2 + x^3 + x^4 + \dots$$

$$3. y = x - 3x^2 + 5x^3 - 7x^4 + \dots$$

$$4. y = x + 2x^2 + 3x^3 + 4x^4 + \dots$$

$$5. y = 2x + 3x^2 + 4x^3 + 5x^4 + \dots$$

$$6. y = x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \dots$$

$$7. y = x - 3x^2 + 5x^3.$$

$$8. y = x - 2x^2 + 2x^3 - 2x^4 + \dots$$

9. Find the approximate value of x to four terms in the series

$$\frac{1}{2} = \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{30} + \frac{x^4}{56} + \dots$$

SOLUTION

$$\begin{aligned}\text{Reverting, } x &= 2\left(\frac{1}{2}\right) - \frac{1}{3}\left(\frac{1}{2}\right)^2 - \frac{1}{45}\left(\frac{1}{2}\right)^3 - \frac{1}{1575}\left(\frac{1}{2}\right)^4 - \dots \\ &= 1 - \frac{1}{6} - \frac{1}{90} - \frac{1}{1575} - \dots \\ &= .8189 +.\end{aligned}$$

Find the approximate value of x in the following:

$$10. \quad \frac{1}{2} = x + \frac{x^2}{6} + \frac{x^3}{24} + \frac{x^4}{60} + \dots$$

$$11. \quad \frac{1}{5} = x - \frac{x^2}{3} + \frac{3x^3}{10} - \frac{2x^4}{7} + \dots$$

**BINOMIAL THEOREM—FRACTIONAL AND
NEGATIVE EXPONENTS**

493. It has been shown (§ 445, formula IV) that when n is a positive integer,

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{1 \cdot 2}x^2 + \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3}x^3 + \dots \quad (1)$$

It is yet to be proved that this formula is true when n is a positive fraction, a negative integer, or a negative fraction.

I. *When n is a positive fraction.*

Let $n = \frac{p}{q}$, in which p and q are positive integers.

$$\text{Then, § 246,} \quad (1+x)^{\frac{p}{q}} = \sqrt[q]{(1+x)^p}$$

$$\text{§ 444,} \quad = \sqrt[q]{1+px+\dots} \quad (2)$$

$$\text{Assume, § 490, } \sqrt[q]{1+px+\dots} = A + Bx + Cx^2 + \dots \quad (3)$$

where x may have any value that will make both series *convergent*.

Raising both members to the q th power,

$$\begin{aligned}1+px+\dots &= [A + (Bx + Cx^2 + \dots)]^q \\ &= A^q + qA^{q-1}(Bx + Cx^2 + \dots) + \dots\end{aligned}$$

Equating the coefficients of like powers of x in two terms,

$$1 = A^r \text{ and } p = qA^{r-1}B;$$

whence,
$$A = 1, \text{ and } B = \frac{p}{q}.$$

Substituting these values in (2) and (3),

$$(1+x)^{\frac{p}{q}} = 1 + \frac{p}{q}x + \dots$$

That is, the formula is true for *two* terms, when n is a positive fraction.

II. When n is negative, and either integral or fractional.

§ 244,
$$(1+x)^{-n} = \frac{1}{(1+x)^n}$$

§ 444 and Case I,
$$= \frac{1}{1+nx+\dots}$$

Dividing the numerator by the denominator,

$$(1+x)^{-n} = 1 - nx + \dots$$

That is, the formula is true for *two* terms when n is negative and either integral or fractional.

Therefore, the formula is true for *two* terms, whether the exponent is positive or negative, integral or fractional, and the coefficient of the second term is n .

494. To find the coefficients of terms after the second, assume

$$(1+x)^n = 1 + nx + Ax^2 + Bx^3 + Cx^4 + \dots \quad (1)$$

Since only two terms of the expansion of the first member are *known*, the coefficients A , B , etc., cannot be determined from (1) in its present form.

To find the value of the undetermined coefficients A , B , etc., we involve them in an identical equation, so that the coefficients of like powers of the same variable may be equated (§ 487).

Since in (1) x represents any number, put $1 + (x+z)$ for $1+x$.

Then, $(1 + x + z)^n = [1 + (x + z)]^n$

$$\begin{aligned} \text{by (1),} \quad &= 1 + n(x + z) + A(x + z)^2 + B(x + z)^3 + \dots \\ &= 1 + nx + Ax^2 + Bx^3 + \dots \\ &\quad + (n + 2Ax + 3Bx^2 + \dots)z + \dots. \end{aligned} \quad (2)$$

Since in (1) x represents any number, and since $1 + (x + z)$

$$= (1 + x) + z = (1 + x) \left(1 + \frac{z}{1 + x} \right), \text{ put } 1 + \frac{z}{1 + x} \text{ for } 1 + x.$$

Then,

$$\begin{aligned} (1 + x + z)^n &= (1 + x)^n \left(1 + \frac{z}{1 + x} \right)^n \\ \text{by (1),} \quad &= (1 + x)^n \left(1 + n \frac{z}{1 + x} + A \frac{z^2}{(1 + x)^2} + B \frac{z^3}{(1 + x)^3} + \dots \right) \\ &= (1 + x)^n + n(1 + x)^{n-1}z \\ &\quad + A(1 + x)^{n-2}z^2 + B(1 + x)^{n-3}z^3 + \dots. \end{aligned} \quad (3)$$

Therefore, from (3) and (2), Ax. 1,

$$\begin{aligned} (1 + x)^n + n(1 + x)^{n-1}z + A(1 + x)^{n-2}z^2 + B(1 + x)^{n-3}z^3 + \dots \\ = 1 + nx + Ax^2 + Bx^3 + \dots + (n + 2Ax + 3Bx^2 + \dots)z + \dots \end{aligned}$$

when both members are convergent.

Equating the coefficients of z ,

$$\S 487, \quad n(1 + x)^{n-1} = n + 2Ax + 3Bx^2 + \dots.$$

Multiplying each member by $1 + x$,

$$n(1 + x)^n = n + (2A + n)x + (3B + 2A)x^2 + \dots.$$

$$(1) \times n, \quad n(1 + x)^n = n + n^2x + nAx^2 + \dots.$$

Equating coefficients, $2A + n = n^2$,

$$\text{and} \quad 3B + 2A = nA;$$

$$\text{whence,} \quad A = \frac{n(n-1)}{1 \cdot 2},$$

$$\text{and} \quad B = \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3}.$$

In like manner any number of successive coefficients may be found.

Substituting these values in (1),

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{1 \cdot 2} x^2 + \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3} x^3 + \dots \quad (4)$$

The expansion of $(1+x)^n$ is not the expansion for the most general form of a binomial, since the first term is 1; consequently, the equation must be changed so as to express the expansion of $(a+x)^n$ for the most general form.

Putting $\frac{x}{a}$ for x ,

$$\left(1 + \frac{x}{a}\right)^n = 1 + n \frac{x}{a} + \frac{n(n-1)}{1 \cdot 2} \frac{x^2}{a^2} + \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3} \frac{x^3}{a^3} + \dots, \quad (5)$$

or $\left(\frac{a+x}{a}\right)^n = \frac{1}{a^n} \left(a^n + na^{n-1}x + \frac{n(n-1)}{1 \cdot 2} a^{n-2}x^2 + \dots\right).$

Multiplying by a^n ,

$$(a+x)^n = a^n + na^{n-1}x + \frac{n(n-1)}{1 \cdot 2} a^{n-2}x^2 + \dots \quad (6)$$

The binomial formula has thus been proved true when n is any positive integer, any positive fraction, any negative integer, or any negative fraction, that is, when n is any *commensurable* exponent, provided the second member of the formula is convergent when it is an infinite series.

495. By reference to (5), § 494, it will be seen that when $x > a$ the series developed by the binomial formula is divergent, but when $x < a$ the series is convergent, if the series is infinite. Hence, the expansion of $(a+x)^n$, when n is fractional or negative, is not feasible when $x > a$ numerically. However, when $x > a$, the n th power of $(a+x)$ may be obtained by expanding $(x+a)^n$, because the series will then be convergent.

Exercises for practice in the use of the binomial formula will be found on pages 401 to 404.

Inasmuch as the preceding discussion has been given for the purpose of establishing the truth of a theorem already assumed to be true, it is unnecessary to give examples for practice here.

PERMUTATIONS AND COMBINATIONS

496. All the different orders in which it is possible to arrange a given number of things, taking either some or all of them at a time, are called the **Permutations** of the things.

Thus, the permutations of the letters a and b are ab, ba ; the permutations of three letters, two at a time, are ab, ac, ba, bc, ca, cb .

497. All the different selections that it is possible to make from a given number of things taking either some or all of them at a time, without regard to the order in which they are placed, are called the **Combinations** of the things.

Thus, while the permutations of three letters, two at a time, are ab and ba, bc and cb , and ca and ac , their *combinations*, two at a time, are ab (or ba), bc (or cb), and ac (or ca); again, the six permutations of three letters among themselves, viz., $abc, acb, bca, bac, cab, \text{ and } cba$, form but one combination, abc (or acb , or bca , or bac , or cab , or cba).

It is evident that there can be but one combination of any number of things taken all at a time.

498. Notation. — The symbol for the number of permutations of n different things, taken r at a time, is P_r^n ; of n different things, taken n at a time, or all together, P_n^n .

The symbol for the number of combinations of n different things, taken r at a time, is C_r^n ; of n different things, taken n at a time, or all together, is C_n^n .

499. The product of the successive natural numbers from 1 to n , or from n to 1, inclusive, is called **factorial n** , written $[n$.

$$[5 = 1 \times 2 \times 3 \times 4 \times 5, \text{ or } 5 \times 4 \times 3 \times 2 \times 1;$$

$$[n = 1 \cdot 2 \cdot 3 \cdots (n-2)(n-1)n, \text{ or } n(n-1)(n-2)(n-3) \cdots 3 \cdot 2 \cdot 1.$$

$$[n \text{ is sometimes written } n!.$$

500. To find the number of permutations of n different things taken r at a time.

Since the permutations of the letters a, b , and c , taken 2 at a time, are ab and ac, ba and bc, ca and cb , formed by writing after each of the letters a, b , and c , each of the other letters in turn, the number of permutations of 3 different things taken 2 at a time is 3×2 .

The number of permutations of n letters taken 2 at a time may be found by associating with each of the n letters each of the $n - 1$ other letters. Consequently, the number of permutations of n different things taken 2 at a time is $n(n - 1)$.

Since the number of permutations of n letters 2 at a time is $n(n - 1)$, if the letters are taken 3 at a time there will be $n - 2$ letters each of which may be associated with each of the $n(n - 1)$ permutations of letters taken 2 at a time. Hence, the number of permutations of n different things taken 3 at a time is

$$n(n - 1)(n - 2).$$

PRINCIPLE 1. — *The number of permutations of n different things taken r at a time is equal to the continued product of the natural numbers from n to $n - (r - 1)$ inclusive. The number of factors is r . That is,*

$$\begin{aligned} P_r^n &= n(n - 1)(n - 2) \dots \text{to } r \text{ factors} \\ &= n(n - 1)(n - 2) \dots (n - r + 1). \end{aligned} \quad (\text{I})$$

Multiplying and dividing the second member of (I) by

$(n - r)(n - r - 1)(n - r - 2) \dots 2 \cdot 1$; that is, by $\underline{n - r}$,

$$P_r^n = \frac{n}{\underline{n - r}}. \quad (\text{II})$$

It will usually be more convenient to employ formula (I) in solving numerical examples; but when simply algebraic results are desired, formula (II) will be preferable.

501. When $r = n$; that is, when the things are taken all together, the last, or n th, factor in (I) is 1. Consequently,

PRINCIPLE 2. — *The number of permutations of n different things taken all at a time is equal to \underline{n} . That is,*

$$P_n^n = n(n - 1)(n - 2) \dots \text{to } n \text{ factors} = \underline{n}. \quad (\text{III})$$

EXAMPLES

1. Three boys enter a car in which there are 5 empty seats. In how many ways may they choose seats?

SOLUTION. — Since the first boy may choose any one of 5 seats; and since, after he has chosen one of them, for each seat that he may choose, the second boy may choose any one of the 4 seats remaining, the greatest possible number of ways in which two of the boys may be seated is 5×4 .

Again, since after each choice of seats made by two of the boys there will be left to the third boy a choice of one of the 3 seats remaining, the number of ways in which all may choose seats is $5 \times 4 \times 3$, or 60.

Or, by (I), $P_r^n = n(n-1)(n-2) \dots (n-r+1)$,

$$P_3^5 = 5 \times 4 \times 3 = 60.$$

2. How many numbers between 100 and 1000 can be expressed by the figures 1, 3, 5?

SOLUTION. — Since the numbers lie between 100 and 1000, each must be expressed by three figures. Hence, the number of numbers between 100 and 1000 that can be expressed by the figures 1, 3, and 5 is the same as the number of permutations of these 3 figures taken 3 at a time.

Since, Prin. 2, $P_3^3 = \underline{3} = 3 \cdot 2 \cdot 1 = 6$,
there are six such numbers. They are 135, 153, 351, 315, 513, and 531.

3. How many permutations can be made of the letters in the word *Albany*, each beginning with capital A?

SOLUTION. — Since A is to be prefixed to each permutation of the 5 other letters, the required number is

$$P_5^5 = 5 \times 4 \times 3 \times 2 \times 1 = 120.$$

4. In how many orders may 4 persons sit on a bench?

5. How many permutations may be made of the letters in the word *steam*?

6. If 10 athletes run a race, in how many ways may the first and second prizes be awarded?

7. In how many different orders may the colors violet, indigo, blue, green, yellow, orange, and red be arranged?

8. There are five routes to the top of a mountain. In how many ways may a person go up and return by a different way?

502. To find the number of combinations of n different things taken r at a time.

Since two letters, as a and b , have two permutations, ab and ba , but form only one combination, the number of combinations of n letters taken 2 at a time is one half the number of permutations of n letters taken 2 at a time.

Since three letters taken 3 at a time have 3×2 permutations, but form only one combination, the number of combinations of n letters taken 3 at a time is obtained by dividing the number of permutations of n letters taken 3 at a time by 3×2 .

Since four letters taken 4 at a time have $\lfloor 4$ permutations but form only one combination, to obtain the number of combinations of n letters taken 4 at a time, the number of permutations of n letters taken 4 at a time must be divided by $\lfloor 4$.

Hence it follows that:

PRINCIPLE 3. — *The number of combinations of n different things taken r at a time is equal to the number of permutations of n different things taken r at a time, divided by the number of permutations of r different things taken all together.* That is,

$$\begin{aligned} C_r^n &= P_r^n \div P_r^r = \frac{n(n-1)(n-2) \dots \text{to } r \text{ factors}}{r(r-1)(r-2) \dots \text{to } r \text{ factors}} \\ &= \frac{n(n-1)(n-2) \dots (n-r+1)}{1 \cdot 2 \cdot 3 \dots r} \quad (\text{IV}) \end{aligned}$$

or, by (II),

$$\begin{aligned} &= \frac{\lfloor n}{n-r} \div \lfloor r \\ &= \frac{\lfloor n}{\lfloor r \lfloor n-r} \quad (\text{V}) \end{aligned}$$

503. Since for every combination of r things out of n different things there is left a combination of $n-r$ things, it follows that:

PRINCIPLE 4. — *The number of combinations of n different things is the same when taken $n-r$ at a time as when taken r at a time.* That is,

$$C_{n-r}^n = C_r^n = \frac{\lfloor n}{\lfloor r \lfloor n-r} \quad (\text{VI})$$

The above principle may be established as follows :

$$\text{By (V),} \quad C_r^n = \frac{|n|}{|r| |n-r|}. \quad (1)$$

$$\begin{aligned} \text{Substituting } n-r \text{ for } r, \quad C_{n-r}^n &= \frac{|n|}{|n-r| |n-(n-r)|} \\ &= \frac{|n|}{|n-r| |r|}. \end{aligned} \quad (2)$$

Since the second members of (1) and (2) are identical, $C_{n-r}^n = C_r^n$.

The above principle is useful in abridging numerical computations.

Thus, the number of combinations of 18 things taken 16 at a time is computed by Prin. 3 as follows :

$$C_{16}^{18} = \frac{18 \cdot 17 \cdot 16 \cdot 15 \cdot 14 \cdot 13 \cdot 12 \cdot 11 \cdot 10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8 \cdot 9 \cdot 10 \cdot 11 \cdot 12 \cdot 13 \cdot 14 \cdot 15 \cdot 16} = 153.$$

By Prin. 4, the computation is abridged as follows :

$$C_{16}^{18} = C_2^{18} = \frac{18 \cdot 17}{1 \cdot 2} = 153.$$

EXAMPLES

1. A man has 6 friends and wishes to invite 4 of them to dinner. In how many ways may he select his guests ?

SOLUTION.—Since each party, or combination, of 4 guests could be arranged, or permuted, in $|4|$ ways, the number of combinations must be $\frac{1}{|4|}$ of the number of permutations of 6 things taken 4 at a time.

Hence, the number of ways is

$$C_4^6 = P_4^6 \div P_4^4 = \frac{6 \times 5 \times 4 \times 3}{1 \times 2 \times 3 \times 4} = 15.$$

2. A man and his wife wish to invite 11 of their friends, 6 men and 5 women, to dinner, but find that they can entertain only 8 guests. In how many ways may they invite 4 men and 4 women ?

SOLUTION.—As in the previous example, 4 men may be selected from 6 men in 15 ways, and in a similar manner 4 women may be selected from 5 women in 5 ways.

Since when any set of 4 men has been invited, the party of 8 may be completed by inviting any one of 5 sets of 4 women, the whole number of different parties that it is possible to invite is 15×5 , or 75. That is,

$$C_4^6 \times C_4^5 = \frac{6 \cdot 5 \cdot 4 \cdot 3}{1 \cdot 2 \cdot 3 \cdot 4} \times \frac{5 \cdot 4 \cdot 3 \cdot 2}{1 \cdot 2 \cdot 3 \cdot 4} = 75.$$

3. In how many ways may a baseball nine be selected from 12 candidates?

4. Find the value of C_2^{10} ; of C_3^{12} ; of C_4^{25} .

5. How many different combinations of 5 cards can be formed from 52 cards?

6. Which is the greater, C_3^{10} or C_3^{10} ? C_4^{10} or C_5^{10} ?

7. From 11 Republicans and 10 Democrats how many different committees can be selected composed of 6 Republicans and 5 Democrats?

8. A man forgets the combination of figures and letters by which his safe is opened. If they are arranged on the circumferences of three wheels, one bearing the numbers 0 to 9 inclusive, another the letters A to M inclusive, and the third the letters N to Z inclusive, what is the greatest number of trials he may have to make to open the safe?

9. From 6 consonants and 4 vowels how many words may be formed each consisting of 4 consonants and 2 vowels, if any arrangement of the letters is considered a word?

SOLUTION. — The number of combinations is $C_4^6 \times C_2^4$; and since by permuting the letters of each combination $\underline{6}$ words can be formed, the number of words is $C_4^6 \times C_2^4 \times \underline{6}$.

10. In an omnibus that will seat 8 persons on a side there are seated 4 persons, 3 on one side and 1 on the other. In how many ways may 12 more persons be seated?

SOLUTION. — Since 5 persons must take seats on one side and 7 persons on the other, 12 persons are to be divided into two classes, 5 and 7. The number of these combinations, formula (V), is

$$C_5^{12}, \text{ or } C_7^{12}, = \frac{\underline{12}}{\underline{5} \underline{7}}.$$

Since each combination of 5 may have $\underline{5}$ permutations of the 5 that compose it, and each combination of 7 may have $\underline{7}$ permutations each of which may be associated with each of the $\underline{5}$ permutations, the required number of ways is $C_5^{12} \times P_5^5 \times P_7^7$,

$$\text{or} \quad \frac{\underline{12}}{\underline{5} \underline{7}} \times \underline{5} \times \underline{7} = \underline{12}.$$

Or, since there are 12 persons to be seated in 12 seats, the number of ways is $P_{12}^{12} = \underline{12}$.

11. Out of 20 consonants and 5 vowels how many words containing 3 consonants and 3 vowels can be formed, if any arrangement of the letters is considered a word?

12. How many different sums can be paid with a cent, a half-dime, a dime, a quarter, and a dollar?

13. From 5 boys and 5 girls how many committees of 6 can be selected so as to contain at least 2 boys?

14. A company of a soldiers is joined by another company of b soldiers. In how many ways is it possible to leave c of them to garrison the fort, dividing the rest into two scouting parties, one of m , the other of n soldiers?

15. If $C_5^n = 2 C_2^n$, find the number of things.

SOLUTION.—By formula (V), $C_5^n = \frac{|n|}{|5| |n-5|}$ and $C_2^n = \frac{|n|}{|2| |n-2|}$.

Since $C_5^n = 2 C_2^n$,

$$\frac{|n|}{|5| |n-5|} = \frac{2 |n|}{|2| |n-2|}$$

$$\frac{1}{|5| |n-5|} = \frac{1}{|n-2|}$$

$$\therefore |n-2| = |5| |n-5|.$$

$$\frac{|n-2|}{|n-5|}, \text{ or } (n-2)(n-3)(n-4) = 5 \times 4 \times 3 \times 2 \times 1,$$

or

$$(n-2)(n-3)(n-4) = 6 \times 5 \times 4.$$


$$\therefore n = 8.$$

16. If $3 C_3^n = 2 C_4^{n+1}$, find n , C_3^n , and C_4^{n+1} .

504. To find the number of circular permutations of n different things taken n at a time.

Suppose four letters a, b, c, d placed in a fixed position around a circle in the order $abcd$. Since the arrangement may be read $abcd$, $bcda$, $cdab$, or $dabc$, without changing the direction in which the letters are read, it is evident that each circular permutation of 4 letters taken all together takes the place of 4 permutations of the letters all together.

That is, the number of circular permutations of 4 things taken all together is one fourth of the number of permutations of 4 things taken all together.



The whole number of permutations of n things taken all together is $|n$. But if the n things are arranged around a circle, n of these permutations may be obtained from any circular permutation without disturbing the relative positions of the things.

Hence,

PRINCIPLE 5. — *The number of circular permutations of n things taken all together is equal to $\frac{1}{n}$ th of the whole number of their permutations taken all together.* That is,

$$P_n^n (\text{circular}) = \frac{|n|}{n} = |n-1|. \quad (\text{VII})$$

EXAMPLES

1. In how many orders may 6 persons seat themselves around a table?

2. In how many orders may 4 gentlemen and their wives seat themselves around a table?

3. In how many orders may 4 gentlemen and their wives seat themselves around a table so that each gentleman sits opposite his wife?

4. In how many orders may 4 gentlemen and their wives seat themselves around a table so that each gentleman sits opposite a lady?

5. In how many ways may the colors violet, indigo, blue, green, yellow, orange, and red be arranged on a disk, the colors radiating from the center?

505. To find the number of permutations of n things taken n at a time when they are not all different.

If, in the permutation (a, b, c, d, e, f, g) , the letters b, d , and g are permuted while the other letters remain fixed in position, the resulting number of permutations will be the same as the number of permutations of b, d , and g . If b, d , and g are different things, the number of permutations resulting will be $|3|$; but if b, d , and g become alike, there will be but 1 permutation.

That is, the number of permutations of any number of things

when three of them are alike is equal to the number of permutations of the things, considered as all different, divided by $\underline{3}$; if 4 of the things are alike, by $\underline{4}$; if p of the things are alike, by \underline{p} .

Hence, it follows that:

PRINCIPLE 6. — *The number of permutations of n things, taken all together, when p of them are alike, is $\frac{n}{p}$.*

Since, if q of the remaining $n - p$ different things become alike, but different from the p like things, the number of permutations must be divided by \underline{q} ; if r others become alike, by \underline{r} ; etc.: it follows that:

PRINCIPLE 7. — *The number of permutations of n things, taken all together, when p of them are of one kind, q of another, r of another, etc., is $\frac{n}{p \underline{q} \underline{r} \dots}$.*

EXAMPLES

1. How many permutations may be made with the letters of the word *Mississippi* taken all together?

SOLUTION. — The number is $\frac{11}{\underline{4} \underline{4} \underline{2}} = 34650$.

2. How many permutations may be made with the letters of each of the following words, all at a time in each case: *zoölogy*, *coefficient*, *ecclesiastical*, *divisibility*.

3. How many permutations may be made with the letters represented in the product $a^4b^5c^2$ written out in full?

506. To find the total number of combinations of n different things.

The number of combinations of n different things taken successively 1, 2, 3, ... n at a time is called the *total* number of combinations of n things.

The total number of combinations of 2 things is

$$C_1^2 + C_2^2 = 2 + 1 = 3, \text{ or } 2^2 - 1.$$

The total number of combinations of 3 things is

$$C_1^3 + C_2^3 + C_3^3 = 3 + 3 + 1 = 7, \text{ or } 2^3 - 1.$$

The total number of combinations of 4 things is

$$C_1^4 + C_2^4 + C_3^4 + C_4^4 = 4 + 6 + 4 + 1 = 15, \text{ or } 2^4 - 1.$$

PRINCIPLE 8. — *The total number of combinations of n different things is $2^n - 1$.*

The above principle may be established as follows:

§ 445, when n is a positive integer,

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{1 \cdot 2} x^2 + \dots + \frac{n(n-1)(n-2) \dots 1}{1 \cdot 2 \cdot 3 \dots n} x^n.$$

$$\text{If } x = 1, \quad 2^n = 1 + n + \frac{n(n-1)}{1 \cdot 2} + \dots + \frac{n(n-1)(n-2) \dots 1}{1 \cdot 2 \cdot 3 \dots n}.$$

$$\text{Prin. 3,} \quad = 1 + C_1^n + C_2^n + \dots + C_n^n = 1 + C_{\text{total}}^n.$$

$$\therefore C_{\text{total}}^n = 2^n - 1.$$

EXAMPLES

1. How many different sums can be paid with a cent, a 3-cent piece, a half-dime, a dime, a quarter, and a dollar?

SOLUTION. — Total $C^6 = 2^6 - 1 = 63$.

2. A man has 10 friends. In how many ways may he invite one or more of them to dinner?

3. How many different quantities can be weighed by weights of 1 oz., 1 lb., $\frac{1}{2}$ lb., 5 lb., and 10 lb.?

4. How many signals can be made with 7 flags?

5. By permuting the letters of the word *counter*, how many permutations can be formed

(a) ending in *er*?

(b) with *n* as the middle letter?

(c) without changing the position of any vowel?

(d) beginning with a consonant?

6. How many numbers can be formed with the digits 1, 2, 3, 4, 3, 2, 1, so that the odd digits always occupy the odd places?

7. If the number of permutations of n different things taken 5 at a time is equal to 24 times the number of permutations of the same number of things taken 2 at a time, find n .

Higher Mathematics

Crockett's Plane and Spherical Trigonometry

With Tables	\$1.25
The Same. Without Tables	1 00
Logarithmic and Trigonometric Tables (separate)	1.00

A clear analytic treatment of the elements of Plane and Spherical Trigonometry and their practical applications to Surveying, Geodesy, and Astronomy, with convenient and accurate "five place" tables for the use of the student, engineer, and surveyor. Designed for High Schools, Colleges, and Technical Institutions.

Raymond's Plane Surveying

Cloth, 8vo, 485 pages. With Tables and Illustrations	\$3.00
--	--------

A modern text-book for the study and practice of Land, Topographical, Hydrographical, and Mine Surveying. Special attention is given to labor-saving devices, to coördinate methods, and to the explanation of difficulties encountered by young surveyors. The appendix contains a large number of original problems, and full tables for class and field work.

Murray's Integral Calculus

Cloth, 8vo, 302 pages	\$2.00
---------------------------------	--------

An elementary course designed for beginners, and for students in Engineering whose purpose in studying the Integral Calculus is to acquire facility in performing easy integrations, and the power of making applications in practical work.

McMahon and Snyder's Elements of the Differential Calculus

Cloth, 12mo, 336 pages	\$2.00
----------------------------------	--------

A systematic treatise on the Differential Calculus based on the theory of limits and infinitesimals and introducing the latest methods of treatment.

Tanner and Allen's Elementary Course in Analytic Geometry

Cloth, 12mo, 400 pages	\$2.00
----------------------------------	--------

Designed for the general student of Higher Mathematics as well as for the special student in the various departments of engineering, architecture, etc.

Copies of any of the above books will be sent prepaid to any address, on receipt of the price, by the Publishers :

American Book Company

New York
(88)

Cincinnati

Chicago

Text-Books in Trigonometry

CROCKETT'S ELEMENTS OF PLANE AND SPHERICAL TRIGONOMETRY AND TABLES. By C. W. CROCKETT, Professor of Mathematics and Astronomy in Rensselaer Polytechnic Institute	\$1.25
ELEMENTS OF TRIGONOMETRY—without Tables	1.00
LOGARITHMIC AND TRIGONOMETRIC TABLES	1.00

In this work the treatment of the subject is adapted to the needs of beginners while it is at the same time sufficient to meet the requirements of advanced technical institutions and colleges.

So far as possible each article of the text is supplemented by examples showing its applications, and a large number of practical problems, with appropriate diagrams, is introduced to give interest to the study and to show its value. Many of these are problems in Surveying and applications of Spherical Trigonometry to Geodesy and Astronomy. In addition to the analytical proofs, used throughout the book, geometrical proofs are employed in many cases to assist the student to a clearer understanding of the subject.

The Logarithmic and Trigonometric Tables are printed on tinted paper from large, clear, differentiated type to facilitate their use.

PHILLIPS AND STRONG'S ELEMENTS OF PLANE AND SPHERICAL TRIGONOMETRY AND TABLES. By ANDREW W. PHILLIPS, Professor of Mathematics, and WENDELL M. STRONG, Tutor in Mathematics, Yale University	\$1.40
ELEMENTS OF TRIGONOMETRY—without Tables	90 cents
LOGARITHMIC AND TRIGONOMETRIC TABLES	\$1.00
KEY TO PLANE AND SPHERICAL GEOMETRY	1 25

The aim in this work has been to place the essentials of the subject before the student in a simple and lucid form, giving especial emphasis to the things which are of the most importance. Some of its noteworthy features are:—graphic solution of spherical triangles; logical combination of the ratio and line methods; simplicity and directness of treatment; use of photo-engravings of models in the Spherical Trigonometry; emphasis given to the formulas essential to the solution of triangles and other essential points; carefully selected exercises given at frequent intervals and a large number of miscellaneous exercises given in a separate chapter, etc.

The Tables include, besides the ordinary five-place tables, a complete set of four-place tables, a table of Napierian logarithms, tables of the exponential and hyperbolic functions, a table of constants, etc.

Copies of these books will be sent, prepaid, on receipt of the price.

American Book Company

New York
(70)

Cincinnati

Chicago

Text-Books on Surveying

RAYMOND'S PLANE SURVEYING

By WILLIAM G. RAYMOND, C.E., Member American Society of Civil Engineers; Professor of Geodesy, Road Engineering, and Topographical Drawing in Rensselaer Polytechnic Institute \$3.00

This work has been prepared as a manual for the study and practice of surveying. The long experience of the author as a teacher in a leading technical school and as a practicing engineer has enabled him to make the subject clear and comprehensible for the student and young practitioner. It is in every respect a book of modern methods, logical in its arrangement, concise in its statements, and definite in its directions. In addition to the matter usual to a full treatment of Land, Topographical, Hydrographical, and Mine Surveying, particular attention is given to system in office work, to labor-saving devices, the planimeter, slide rule, diagrams, etc., to co-ordinate methods, and to clearing up the practical difficulties encountered by the young surveyor. An appendix gives a large number of original problems and illustrative examples.

Other Text-Books on Surveying

DAVIES'S ELEMENTS OF SURVEYING (Van Amringe) . . .	\$1.75
ROBINSON'S SURVEYING AND NAVIGATION (Root) . . .	1.60
SCHUYLER'S SURVEYING AND NAVIGATION	1.20

Copies will be sent, prepaid, to any address on receipt of the price.

American Book Company

New York
(76)

• Cincinnati •

Chicago

Milne's Plane and Solid Geometry

Half Leather, 12mo, 384 pages. Price, \$1.25

Milne's Plane Geometry—Separate

Half Leather, 12mo, 242 pages. Price, \$0.75

By WILLIAM J. MILNE, Ph.D., LL.D.

President State Normal College, Albany, N. Y.

This new text-book in geometry embodies many original methods of treatment, which while novel and attractive, are based on sound mathematical and pedagogical principles. The author's aim has been to combine the valuable features of inventional and concrete geometry with the forms of rigid deductive reasoning, which is one of the chief objects sought in this branch of study.

In this work the student is introduced to geometry through the employment of inventional steps, supplemented by demonstrations of the fundamental propositions of the science. Every theorem is introduced by questions designed to lead the student to discover the geometrical concept clearly and fully before attempting a demonstration. The laboratory method is introduced at the beginning and is continued throughout the book. A great abundance of undemonstrated theorems and of unsolved problems is supplied for supplementary and original work. Summaries showing the subjects covered and the truths established in the chapters to which they are appended are an original feature of the book. These will be found of great practical value to the student, both in reviewing his work and in preparing him for advanced steps. The diagrams and other figures used in the demonstrations are all that could be desired in a text-book on geometry.

Copies of Milne's Geometries will be sent, prepaid, to any address on receipt of the price by the Publishers :

AMERICAN BOOK COMPANY

NEW YORK . CINCINNATI . CHICAGO


y

, \$1.25

\$0.75

origi-
ttrac-
ogical
the
metry
one

etry
nted
the
de-
ical
ra-
be-
eat
ed
k.
is
e
of
s
-
!





To avoid fine, this book should be returned on
or before the date last stamped below

SON-9.40

--	--	--

